

Which psychosocial interventions improve sex worker well-being? A systematic review of evidence from resource-rich countries

Kevin Turner , Jane Meyrick , Danny Miller , Laura Stopgate

► Additional supplemental material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjsh-2021-201028>).

Department of Health and Social Sciences, University of the West of England, Bristol, UK

Correspondence to

Kevin Turner, Department of Health and Social Sciences, University of the West of England, Bristol BS16 1QY, UK; kevin.turner@live.uwe.ac.uk

Received 4 January 2021

Revised 17 March 2021

Accepted 26 March 2021

Published Online First

29 April 2021



Check for updates

© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Turner K, Meyrick J, Miller D, et al. *BMJ Sex Reprod Health* 2022;**48**:e88–e100.

ABSTRACT

Objective To establish the state of the evidence base around psychosocial interventions that support well-being in sex workers in order to inform policy and practice within a resource-rich geographical context.

Methods Published and unpublished studies were identified through electronic databases (PsychINFO, CINHAL Plus, MEDLINE, EMBASE, The Cochrane Library and Open Grey), hand searching and contacting relevant organisations and experts in the field. Studies were included if they were conducted in high-income settings with sex workers or people engaging in exchange or transactional sex, and evaluated the effect of a psychosocial intervention with validated psychological or well-being measures or through qualitative evaluation.

Results A total of 19 202 studies were identified of which 10 studies met the eligibility criteria. The heterogeneity found dictated a narrative synthesis across studies. Overall, there was very little evidence of good quality to make clear evidence-based recommendations. Despite methodological limitations, the evidence as it stands suggests that peer health initiatives improve well-being in female street-based sex workers. Use of ecological momentary assessment (EMA), a diary-based method of collecting real-life behavioural data through the use of twice-daily questionnaires via a smartphone, increased self-esteem and behaviour change intentions.

Conclusions Work with sex workers should be based on an evidence-based approach. Limitations to the existing evidence and the constraints of this work with vulnerable groups are recognised and discussed.

INTRODUCTION

Work on the well-being of sex workers¹ has traditionally either focused on the

Key messages

- There is a gap in the evidence around the effectiveness of psychosocial interventions aiming to improve sex worker well-being.
- Weak evidence exists to support the benefits of ecological momentary assessment in reducing anxiety and depression and improving opportunities for behaviour change.
- Research on sex working communities focuses on female street-based sex workers and underrepresents the experiences of male sex workers.
- Participatory methodologies are recommended to ensure that future research is grounded in the actual rather than perceived needs of sex working communities.

ways in which legal and human rights issues affect sex worker vulnerability,² or on access to sexual health screening opportunities, in attempts to reduce the acquisition and transmission of sexually transmitted infections (STIs).³

These criminal justice or public health approaches present a limited narrative of sex work as something that is criminal or virologically dangerous. This is further compounded by a focus from sexual health services on biomedical interventions such as condom use,⁴ HIV pre- and post-exposure prophylaxis (PrEP/PEP)⁵ and hepatitis B vaccinations⁶ to support the perceived health needs of sex workers.

The stigma,⁷ labour and complex routes within sex work place unique demands on the coping resources of sex workers⁸ highlighting additional well-being needs. Previous studies based on street-based




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) 2009 flow diagram.

female sex workers evidences some of these psychosocial issues, which include substance misuse, mental health problems, violence, and homelessness.⁹

This systematic review aims to gather and marshal evidence on the range and effectiveness of psychosocial interventions aimed at improving the well-being of sex workers within resource-rich countries with the aim of producing recommendations to inform policy and practice within the UK.

METHODS

Search strategy

PsychINFO, CINAHL Plus, MEDLINE, EMBASE, Open Grey and The Cochrane Library databases were searched throughout January 2020 for peer-reviewed articles published in English between January 2000 and January 2020. Truncated keywords and relevant medical subject headings (MeSH) related to the study's PICO: 'sex workers' (Population), 'psychosocial interventions' (Intervention), 'well-being' (Outcome) were used and linked using Boolean operators (online supplemental appendix 1). The reference list of included articles was also searched in addition to contacting experts in the field and sex worker organisations to further identify any additional eligible articles.¹⁰

Study selection process

The titles of all articles identified from the search were screened by one reviewer (KT). Two reviewers screened the abstracts of the remaining articles (KT and DM). Articles were included if they evaluated a psychosocial intervention using either validated well-being measures or qualitative methodologies, included sex workers or people engaging in exchange or transactional sex,¹¹ and were conducted in high-resource countries (online supplemental appendix 2). Discrepancies were resolved through further discussion with a third reviewer (LS). A protocol for this review was peer reviewed and is registered with the International Prospective Register of Systematic Reviews (PROSPERO) CRD42020204592.

Data extraction

Extracted data inclusive of sample characteristics, intervention type, study methods, outcome measures and findings were extracted by two reviewers (KT and DM) using a standardised form developed by the review team (online supplemental appendix 3).

Quality assessment

A quality assessment was carried out by two reviewers (KT and DM) using the Effective Public Health Practice Project (EPHPP) Quality Assessment Tool for Quantitative Data¹² (online supplemental appendix 4) and the National Institute for Health and Care Excellence (NICE) Quality Appraisal Checklist¹³ for qualitative data (online supplemental appendix 5). A third reviewer (LS) reviewed any discrepancies.

Analysis

The heterogeneity of the intervention's aims, study design, outcome measures and sample populations precluded a meta-analysis of their results. A narrative synthesis across qualitative and quantitative data is presented by intervention type. Scientific quality is used to frame the validity of study effect findings, common methodological flaws, risk of bias, and how well studies were conceptualised.

RESULTS

A total of 19 202 articles were identified from the literature search (figure 1). Seventy articles included studies on psychosocial interventions aimed at improving sex worker well-being. Ten articles were selected after full-text review. Reasons for exclusion are documented using a PRISMA flow¹⁴ diagram (figure 1). An overview of study characteristics and interventions for qualitative, quantitative and mixed-method studies are presented (table 1).

Study characteristics

Studies from all resource-rich countries were eligible for inclusion but only studies from America (n=6),^{15–19} followed by Canada (n=2)^{20 21} and the

Table 1 Summary of study characteristics, data extraction table and quality appraisal outcomes

Study and aim	Geographical location, sample size, characteristics and recruitment method	Intervention	Study design	Outcomes measures	Data collection	Data analysis	Findings	Quality score
Quantitative Data								
Gunn et al. (2016) To describe the psychological benefits associated with participation in a daily diary and interview study ¹⁵	USA, n=24. Female. Median age: 42.5 years. Ethnicity: Black (75%), non-Latina (87%). 58% engaged in transactional sex at least once a week 90 days prior to enrollment. 25% consumed four or more alcoholic drinks per day, 25% used marijuana daily, 42% used cocaine weekly. All participants had been arrested at least once. 58% experienced CSA before the age of 13 years. 42% initiated sex work as minors (before the age of 18). 37% reported receiving little social support from family or friends at baseline. Baseline mean self-esteem score 27/40, depression 9/27. 33% reported high anxiety. Recruitment: Targeted outreach, venue-based recruitment and incentivised peer referral	EMA Twice-daily cell phone diaries capturing information on mood; alcohol craving and use; partnered sexual behaviour and condom use; partner characteristics and locations of sexual events	Quasi-experimental, single group pre-test post-test	Self-esteem: Rosenberg Self-Esteem scale (RSE) Depression: Patient Health Questionnaire (PHQ-9) Anxiety: Brief Symptom Inventory-Anxiety (BSI-A) Taken at baseline and exit (4 weeks later)	(1) 12-Hourly digital diary entries via a smartphone provided by the study. Assessing mood, alcohol/drug craving and use. Partnered sexual behaviours including condom use, partner characteristics and locations of sexual events (2) Weekly face-to-face interviews with study personnel focusing on context of sexual lives supported by diary entries	Paired t-tests to compare self-esteem, anxiety and depression at baseline and exit	Self-esteem improved significantly increasing 4.08 points from baseline to exit (p<0.001). Mean scores for depression and anxiety decreased from baseline to exit but were overall not statistically significant. Women who initiated sex work as minors reported decreased depression between baseline and exit (4.1 points, p=0.05). Anxiety decreased in women who drank less than 4 alcoholic drinks per day (1.9 points, p=0.03) and those who used marijuana daily (3.7 points, p=0.05).	Weak (EPHPP)
Litchfield et al. (2010) Outcomes of a primary care drugs treatment intervention for sex workers who use heroin, using PMT and psychosocial support ²²	UK, n=34. Female. Street-based sex workers with a physiological addiction to heroin. Recruitment: Incoming referrals, clinical attendance to evaluated service provision	PMT for heroin use in conjunction with undefined intensive health and psychosocial support	Quasi-experimental, single group pre-test post-test	Christo Inventory (QoL). Urine samples for opiate screening at the point of entry and at 1 year. Self-reported involvement in sex work	Urine samples, Christo Inventory	Paired t-tests comparing pre-test and post-test Christo scores	Reduction in mean Christo score from 12.05 at entry to 8.97 after 1 year (p<0.001). 67% reduction in sex work involvement. 87% of urine samples tested positive to heroin at baseline compared with 72% at follow-up.	Weak (EPHPP)
Qualitative Data								

Continued

Table 1 Continued

Study and aim	Geographical location, sample size, characteristics and recruitment method	Intervention	Study design	Outcomes measures	Data collection	Data analysis	Findings	Quality score
Arnold et al. (2000) To explore the psychosocial treatment needs of female street-walking prostitutes by drawing on a recent programme evaluation of a CMP that targets female prostitutes who have involvement in the criminal justice system ¹⁶	USA, n=23. Programme staff (n=4), community professionals (n=9), street-walking female sex workers (n=10). Sex worker average age: 32 years (range 23–38 years). Ethnicity: African American (n=2), White (n=8). Drug/alcohol use: crack cocaine users (n=8), alcohol (n=2), heroin (n=1). Referred while in jail (n=9). 8 diagnosed with mental health disorder: bipolar disorder (n=3), depression (n=3), ADHD (n=1), anxiety (n=1). Average (mean) time in the programme at the point of interview 5.9 months	CMPs In CMPs a named case manager acts as a fixed point of contact for a patient during the co-ordination of their care including referrals to relevant support agencies	Phenomenological		Observation of a support group for programme participants and face-to-face (n=7) or telephone (n=2) interviews with clients, staff and community professionals	Not reported	(1) Substance misuse treatment programmes should be specific to prostitution. (2) Challenges experienced by staff helping clients to achieve stability with mental health. (3) Child custody expressed as a realistic short-term goal by sex workers and were anxious to access relevant support. (4) Access to child custody support was felt to be outside of the scope of CMP by support staff – who promoted the need for respect and support in client's rights to self-determine. (5) Women do not view themselves as victims.	(–) (NICE)
Benoit et al. (2017) Evaluation of a pilot study facilitating peer-based community leadership by designing and evaluating a small STBBI prevention strategy led by sex workers as peer educators ²⁰	Canada, n=5. Sex workers (independent indoor/webcam/escort agency/independent outdoor), varied in age, gender, sexual orientation, indigenous background, socioeconomic status and sex work history – no details provided. Recruitment: Purposive criterion sampling. Cash honorarium \$40 for weekly interventions and \$40 for participating in debriefing sessions	An 8-week STBBI peer health education programme. Developed in consultation with sex workers delivering sessions on empowerment, health and safety, harm reduction, diversity awareness and overdose prevention	Phenomenological		(1) Qualitative semi-structured interviews with participants prior to the training, after the training and at the end of an 8-week intervention phase. (2) Journals kept by the participants and project co-ordinator after each training and debrief session	Thematic analysis NVivo	(1) Reduced internalised stigma and increased self-esteem in participants. (2) Heightened critical consciousness relating to diversity within the sex industry. (3) Improved knowledge of local services. (4) Increased confidence in challenging stigma from front-line services.	(++) (NICE)

Continued

Table 1 Continued

Study and aim	Geographical location, sample size, characteristics and recruitment method	Intervention	Study design	Outcomes measures	Data collection	Data analysis	Findings	Quality score
Bodkin et al. (2015) To highlight the effectiveness of a collaboration between a healthcare physician and police officer in accessing street level sex worker populations in London, Ontario, Canada ²¹	Canada, n=14. Street-level female sex workers. Age range: 23–49 years. Time in sex work: 2 months to 34 years. Ethnicity: Caucasian (n=11), Native (n=3). Sex work status: active (n=5), semi-retired (n=1), exited (n=8). Recruitment: Snowball sampling. Travel reimbursement, coffee shop voucher	PAR Outreach programme provided by a female police officer and GP to improve access to law enforcement and healthcare services	Phenomenological		(1) 15–60 minute 1-2-1 semi-structured interviews with female sex workers enrolled in the PAR programme (n=14). (2) Semi-structured interviews with health and law enforcement professionals	Qualitative Descriptive Analytical approach	(1) PAR programme highlighted as essential in improved assess and continued engagement with healthcare/law enforcement services. (2) Treatment needs for mental health and addiction prioritised. (3) Location and flexible opening hours in addition to female gender of service workers favoured.	(–) (NICE)
Felsher et al. (2018) How participation in EMA – a self-report diary intervention – improved the mental health of women engaging in transactional sex ²³	USA, n=25. Female. Median age: 42.5 years. Ethnicity: 75% Black. 58% engaged in transactional sex at least once per week. 25% consumed +4 alcoholic drinks per day, 25% used marijuana daily, 42% used cocaine weekly. 42% initiated sex work before the age of 18 years. 37% reported receiving little social support from friends or family. Recruitment: Targeted outreach, venue-based recruitment, incentivised peer referral	EMA (1) 12-Hourly digital diary entries via a smartphone provided by the study assessed mood, alcohol/drug craving and use. Partnered sexual behaviours – including condom use, partner characteristics and locations of sexual events, over a period of 4 weeks. (2) Weekly face-to-face interviews with study personnel focusing on context of sexual lives supported by diary entries	Phenomenological		Interviews were conducted in a private room at community-based organisations	Thematic analysis	Participants reported that EMA through twice-daily electronic diaries increased self-reflection and heightened self-awareness of their emotions and behaviour. Resulting in either actual or intended changes in behaviour including: decreased engagement in sex work, sobriety, procurement of condoms and addressing negative behavioural triggers. F	(+) (NICE)
Jeal et al. (2017) To explore street sex workers views and experiences of drug treatment ¹	UK, n=24. Female street-based sex workers. Age: 26–54 years. 24 disclosed experience of street-based sex work and dependency on crack cocaine/heroin in the previous 5 years, 14 had injected drugs at most recent use, 6 reported using illicit drugs daily. Recruitment: Flyers displayed in venues of sex worker support organisations. £02 shopping voucher	Drug treatment groups	Phenomenological		20–90-min in-depth interviews undertaken at the site of recruitment, university setting or in the participants own home. Topic guide used for consistency	Framework analysis	(1) Inability to discuss sex work within drug treatment groups undermined engagement in treatment processes. (2) Disclosure of sex work often resulted in stigma or unwanted attention. (3) Recommendations made for 1-2-1 therapy with female therapists and sex worker-only treatment groups.	(+) (NICE)

Continued

Table 1 Continued

Study and aim	Geographical location, sample size, characteristics and recruitment method	Intervention	Study design	Outcomes measures	Data collection	Data analysis	Findings	Quality score
Preble et al. (2016) Describe the experiences of female sex workers receiving services to support sustained exit from the sex industry ¹⁷	USA, n=13. Female. Prostitution (n=5), exotic dancer (n=5), exotic dancing/prostitution (n=2), prostitution/erotic massage (n=1). Industry years ranged between 3.5 and 20+ years. 9 reported a history of substance misuse. Participants divided into three groups depending on length and level of participation within the programme: Beginners from 3 months to 12 months (n=7), middle participants had been with the agency for more than 1 year but had not graduated (n=6) and graduates who had completed all requirements of the agency's programming (n=7). Four participants from each group were randomly selected to make up the final sample. A 13th participant was selected to serve as a "pilot case" to test the interview questions.	Faith-based 'Fresh Start Programme' providing legal and medical assistance referrals, employment services, counselling, educational opportunities, financial literacy, life skills training and computer literacy	Phenomenological		1.5-hour semi-structured, face-to-face interviews. Conducted in private rooms chosen by participants located within the church. Focus groups to sense-check initial data analysis	Thematic analysis	(1) Financial assistance for basic needs was powerful for beginners. (2) Opportunities for peer support promoted community cohesion and reinvestiture. (3) Support with budgeting was experienced as patriarchal and disempowering. (4) More support could be offered around housing assistance. (5) CMPs were beneficial; however, these were criticised for being grounded in assumption rather than the actual needs of service users.	(++) (NICE)

Continued

Table 1 Continued

Study and aim	Geographical location, sample size, characteristics and recruitment method	Intervention	Study design	Outcomes measures	Data collection	Data analysis	Findings	Quality score
Wahab (2006) Evaluate the usefulness of a PDP ¹⁸	USA, n=31. Female. Current or ex-sex workers (n=12) and non-sex workers-service providers (n=19). Age range: 22–43 years. Age when starting sex work: 15–22 years. Industry years range: 1 month to 27 years. Ethnicity: Hispanic (n=3), African American (n=1), White/Caucasian (n=7), Bi-racial (n=1). Education: 1–3 years of high school to a college degree. Recruitment: Sex workers referred to the PDP at the time of the study received a generic recruitment letter. Honorarium at \$20 per hour	PDP was designed and operationalised as a three-phase programme lasting for 40 weeks. Each phase required attendance at weekly group sessions of facilitated harm reduction workshops, individual counselling and engagement with outside treatment providers	Phenomenological		1–2-hour semi-structured qualitative interviews, extended on-site observations, field notes and copies of written programme materials	Open coding Atlas-TI	Sex workers: felt supported by counsellors and felt they were able to teach PDP stakeholders about sex work, reported engaging in harm reduction behaviours and consequently believed they were protecting themselves from HIV. Personal benefits include individual therapy, resource referrals and help with sobriety.	(+) (NICE)
Mixed-Methods Data								

Continued

Table 1 Continued

Study and aim	Geographical location, sample size, characteristics and recruitment method	Intervention	Study design	Outcomes measures	Data collection	Data analysis	Findings	Quality score
Decker et al. (2018) Evaluate a brief trauma-informed intervention to improve safety and reduce HIV risk among female sex workers ¹⁹	USA, n=60. Female, 73 % street-based sex workers, 27% venue-based. Average age: 35.3 years. Ethnicity: 72% White, 16% Black. 41% relied on sex work as their sole income, 86% current IVDU. Recruitment: Participants were recruited from a street-based needle exchange outreach van in a location with high levels of sex trading activity	INSPIRE (Integrating Safety Promotion with HIV Risk Reduction) Intervention consists of a brief semi-structured dialogue, reinforced with a safety card. Content includes trauma-informed support, validation, safety promotion and links to services	Quasi-experimental, single group pre-test post-test	Revised Conflict Tactics Scale (CTS) adapted for sex work, six-item Perceptions of Abuse Likert Scale, Sex Work-specific Rape Myths Scale adapted from general population instruments, Sex Work Safety Behaviour Scale adapted from the Safety Promoting Behaviour Checklist, Condom Confidence Scale, Depression: CESD, PTSD 17-item PTSD checklist, Intervention acceptability Likert Scale	Pre-test self-administered survey. Post-intervention exit survey. 10–12-week follow-up survey and participation in a 25–45-min in-depth interview. Interventions and data collection were conducted in private locations in the outreach van or adjacent vehicles	Descriptive analysis was calculated for baseline characteristics. Attrition analysis compared baseline characteristics to those retained using t-tests and chi-square analysis. Baseline and follow-up data were evaluated using paired t-tests. Iterative thematic analysis was undertaken on qualitative data from interviews	Increase in Safety Behaviour Scores (51.2 vs 58.1, $p<0.0001$) and use of support programmes responding to intimate partner violence increased from 10.5% to 28.9% ($p<0.01$) between baseline and follow up. Avoidance of condom negotiation decreased between baseline and follow-up (2.0 vs 1.4, $p=0.04$). Average frequency of sex with clients while under the influence of drugs or alcohol decreased (mean=4.4 vs 4.0, $p=0.04$). PTSD and depression symptomology were high at baseline (PTSD mean=51.4; CESD mean=19.2), no changes were observed from baseline to follow-up.	Weak (EPHPP) (+) (NICE)

ADHD, attention-deficit hyperactivity disorder; CESD, Center for Epidemiological Studies-Depression Scale; CMP, case management programme; CSA, childhood sexual abuse; EMA, ecologic momentary assessment; EPHPP, Effective Public Health Practice Project; GP, general practitioner; IVDU, intravenous drug user; NICE, National Institute for Health and Care Excellence; PAR, Persons at Risk Programme; PDP, Prostitution Diversion Project; PMT, prescribed maintenance treatment; PTSD, post-traumatic stress disorder; QoL, quality of life; STBB1, sexually transmitted and blood-borne infection.

UK (n=2)^{1 22} were identified. Five studies exclusively focused on street-based sex workers.^{1 16 19 21 22} Three studies included sex workers from a variety of contexts including exotic dancing, erotic massage and escorting,^{17 18 20} while two studies investigated participants engaging in transactional sex.^{15 23}

Intervention characteristics

Interventions included ecologic momentary assessment (EMA) (n=2),^{15 23} drug treatment services (n=2),^{1 22} exiting and diversion programmes (n=2),^{17 18} trauma-informed interventions (n=2),^{19 21} peer health initiatives (n=1)²⁰ and case management programmes (CMPs) (n=1).¹⁶

Evaluation design

Well-being outcomes were measured in quantitative studies by assessing quality of life (QoL), anxiety, depression and post-traumatic stress disorder (PTSD) using validated measures. One study used the Christchurch Inventory,²² a widely used clinical tool validated as a QoL measure in people who use drugs. Two studies measured depression using either the Patient Health Questionnaire (PHQ-9)¹⁵ or the Center for Epidemiological Studies-Depression (CES-D) Scale.¹⁹ Anxiety was measured in one study using the Brief Symptom Inventory-Anxiety (BSI-A).¹⁵ One study measured PTSD symptom severity using the PTSD checklist.¹⁹ All quantitative studies used pre- and post-intervention measures.^{15 19 22} The timeframe between measures ranged between 4 weeks^{15 23} and 1 year.²²

Semi-structured interviews, observations, journal entries and field notes were used across qualitative studies^{1 16–21 23} to evaluate the effectiveness of psychosocial interventions.

Summary of findings

Ecologic momentary assessment

EMAs study people's thoughts and behaviour in their daily lives by repeatedly collecting questionnaire data in their normal environment, at or close to the time they carry out that behaviour. This is achieved through regular self-report diary entries covering key information around risks logged by mobile phone.

In a quantitative study investigating the benefits of EMA through smartphone-enabled diary entries every 12 hours and weekly face-to-face interviews, levels of self-esteem increased from 4.08 points from baseline to exit ($p<0.001$) over a period of 4 weeks among a sample of 24 women engaging in transactional sex. While mean scores for anxiety and depression decreased from baseline, they were not statistically significant. Women who initiated sex work as minors reported decreased depression between baseline and exit (4.1 points, $p=0.05$). Anxiety also decreased in women who drank less than four alcoholic drinks per day (1.9 points, $p=0.03$) and those who used marijuana daily (3.7 points, $p=0.05$).¹⁵

Statistical analysis through the use of t-tests failed to stratify which element of the intervention, EMA or weekly interviews, had the greatest effect on the otherwise combined outcome measures reported. The short study time documented for participating in EMA impacts on the ability to assess sustainability of intervention success outside of the documented 4 weeks. The small sample size of this study further limits the power of the findings to detect differences across wider populations of sex workers.

In a separate study, qualitative evaluations of women participating in transactional sex who engaged in the same intervention experienced a heightened awareness of their emotions and behaviour, resulting in either actual or intended changes in behaviour, including decreased engagement in sex work, sobriety, procurement of condoms, and addressing negative behavioural triggers.²³ While the exploratory nature of the study, absence of theoretical framework and small sample (n=25) limits the generalisability of the findings, EMA shows some utility in its ability to facilitate behaviour change to further support sex worker well-being.

Exiting programmes

Exiting programmes address the causes and consequences of sex work to encourage industry exit. Group counselling was experienced as being helpful to women participating in qualitative evaluations of a Prostitution Diversion Programme (PDP), specifically in its utility to facilitate conversations around addiction, abuse, trauma, mental health and relationships.¹⁸ These were considered beneficial to both participants and stakeholders who were able to learn directly from lived experiences to further inform and develop group sessions.

Programmes that offered financial assistance to women who were in the early stages of exiting sex work (between 3 months and 1 year) were positively evaluated in a qualitative review of a faith-based exiting programme developed to support sustained exit from the sex industry. Peer support was encouraged by women participating in sex worker support services. Engaging with peers promoted a sense of community belonging and cohesion in addition to presenting opportunities for reinvesting help and support to other sex working women.¹⁷

The lack of quantitative-based research inclusive of validated measures to assess intervention success contributes to the weak evidence base for diversion and exiting programmes. Available evidence is grounded in selection bias due to the fact that programme enrolment is largely offered as an alternative to jail time or dependent on a prerequisite to have already exited from sex work.

Drug support

Prescribed maintenance therapy (PMT) in the form of a regulated and controlled prescription for heroin

to support drug addiction, along with psychosocial support for female street sex workers from a specialist general practice setting in the UK, reported significant improvements in QoL between pre- and post-test measures recorded 1 year apart (12.05 at entry to 8.97 after 1 year, $p < 0.001$).²²

The use of paired t-tests to look for changes in pre- and post-test scores fails to distinguish between the separate effects of PMT and psychosocial interventions. Poor reporting fails to provide a definition as to what psychosocial interventions were offered and how these were accounted for in response to confounding factors. Despite being free from attrition bias, these findings are vulnerable to bias, given that the setting was based in a general practitioner (GP) practice, where over-reporting of healthy behaviour is likely and reporting success could also be perceived by participants as a requisite for securing ongoing prescriptions.

Qualitative evaluations of female street sex workers' experiences of drug treatment services ($n=24$) highlight the importance of providing opportunities for sex working women to openly discuss their drug use free from the unwanted attention of male service users. Across the interview data, participants described how feeling unable to discuss their sex work in drug treatment groups undermined their engagement in treatment processes. Non-disclosure meant that they could not discuss unresolved issues around trauma which emerged or increased when reducing their drug use.¹

Recommendations were made for sex worker-only services to be delivered by female staff. The provision of one-to-one counselling was felt to provide the opportunities for people to explore personal issues in more depth, which was not possible within group settings. However, these claims lack transferability to male or transgendered sex workers given that the findings reflect the voices and experiences of women.²⁴

Trauma-informed interventions

The development of a safety card, developed in consultation with sex workers, providing harm reduction, safety information and support for accessing violence-related services for sex workers attending an outreach needle exchange service, reported an increase in safety behaviour scores (51.2 to 58.1, $p < 0.0001$) and use of support programmes responding to intimate partner violence (10.5% to 28.9%, $p < 0.01$) between baseline and follow-up (10–12 weeks).¹⁹

While no changes were observed at follow-up from high baseline levels of PTSD (mean=51.4) or depression (mean=19.2), avoidance of condom negotiation (2.0 to 1.4, $p=0.04$) and the average frequency of sex with clients while under the influence of drugs or alcohol (mean=4.4 to 4.0, $p=0.04$) decreased. The generalisability of these findings is restricted due to rates of data attrition (39/60) and the population being limited to street-based female sex workers already engaging in relevant risk reduction interventions.

Within qualitative evaluations, new knowledge of support organisations included on the safety card prompted and enabled women to offer peer support to friends and colleagues. An enhanced confidence was experienced by women through open discussions, enabled through use of the card, around topics rarely discussed including coercive barriers to condom use and safety.

The Persons at Risk Programme (PAR), a harm reduction service which aimed to improve access to health-care and essential services for street-level sex workers through outreach work undertaken by a GP and police officer, was qualitatively evaluated in a second trauma-informed intervention study.²¹

The PAR was valued by sex workers for the streamlined and focused nature of care provision in overcoming barriers to services otherwise avoided due to fear of stigma from frontline service staff, including drug abuse, infectious diseases and mental health assistance. However, findings are limited to a sample of women who choose not to access frontline services, who had stopped using drugs and successfully exited sex work.

Peer health initiatives

Findings from qualitative interviews show that peer advocacy in the delivery of a sexually transmitted and blood-borne infection (STBBI) prevention strategy, developed and delivered by sex workers as peer educators,²⁰ led to reduced internalised stigma and increased self-esteem and confidence across participants ($n=5$). Improved critical consciousness and resource mobilisation were attributed to the inclusion of training materials that promoted diversity within sex working communities and awareness of local support agencies.

Small numbers of participants limits the generalisability of findings in this study and inclusion of broader sex working experiences across wider demographics and geographical contexts. The short duration of the study also restricts our understanding of the long-term sustainability and ownership of peer health initiatives. Future studies incorporating quantitative measures of internalised stigma and self-esteem would help to strengthen the evidence base for peer health interventions. However, the study provides proof of concept that local sex working communities are receptive and willing to participate in peer-led health initiatives.

Case management programmes

In CMPs, a named case manager acts as a fixed point of contact for a patient during the co-ordination of care. One study qualitatively evaluated a community-based CMP for street-walking prostitutes in Florida.¹⁶ Across a purposive sample, access to sex worker-specific treatment programmes for substance misuse and support with child custody were identified as important services among sex working women ($n=10$), while support with engaging in mental services was highlighted by

programme staff (n=4) and community professionals (n=9).

The inability to compare outcomes from these services independently of CMP referrals weakens the evidence base for the effectiveness of CMPs. Bias exists in the sample, as the majority of participants had been referred into the service while in jail. Attendance to the programme is likely to be linked to conditions of their parole and therefore not representative of those freely engaging in service provision.

DISCUSSION

Sex workers present with specific health and well-being needs²⁵ beyond the scope of sexual health screening. Despite this, an identifiable gap exists in the current evidence around how to respond to the additional psychosocial needs experienced by sex workers or those engaging in transactional sex.

The results from this review highlight the utility of a range of interventions which aim to improve sex worker well-being including peer health initiatives, EMA (smartphone-based diary intervention), drug support services and trauma-informed interventions. However, the limited information around the study characteristics and small sample sizes reflects low levels of participation beyond street-based female sex workers, limiting the power of studies to detect differences across more diverse and less researched populations including male, transgender and migrant sex workers and those using a range of platforms to engage in wider arenas of sex work.

The implications of these findings are, first, that the field would benefit from broadening definitions of sex work by including wider and more contemporary outlets of sex work such as adult content creators operating on subscription-only platforms. A broader definition of sex work will help to adequately acknowledge and represent the diversification of sex work, helping to challenge perpetuated stereotypes of sex worker identities and their needs.

Poor study design contributed to the weak evidence base for psychosocial interventions aiming to improve sex worker well-being. Across quantitative studies the opportunities for comparison of findings against control groups and an inability to discriminate between intervention effects in statistical analysis impacts our ability to make clear evidence-based recommendations to inform policy and practice within geographical resource-rich countries.

While the evidence for EMA (smartphone-prompted diary approach)¹⁵ remains weak and unsubstantiated, participation in regular diary entries enabled by smartphone technology improved self-esteem while encouraging intention for, or actual, behaviour change.²³ Further research grounded in behaviour change theory would inform the development of EMA and its ability to identify and support individual psychosocial well-being needs.

The small number of methodologically rigorous studies reflects the challenges of studying this population,²⁶ including the ethical issues such as compensation of sex worker time or researcher standpoint on decriminalisation. Furthermore, barriers to sex worker identification and availability include exacerbation of minority stress, given that sex working practices are perceived to differ from the majority of surrounding society²⁷ and potential breaches in confidentiality which may expose sex workers to public disclosure of highly stigmatised and criminalised identities.^{28 29}

In addressing these issues, some studies included in the review recruited sex workers to patient and public involvement (PPI) roles, to assist in the administration of surveys and questionnaires or facilitation of focus groups.³⁰ Very few authors discussed the pro and cons of this approach, particularly with consideration to any impact on the data collected. Similarly, the use of incentives such as gift cards, travel coupons or money to enable participation remained largely undiscussed.

The use of incentives to recruit research participants remains a controversial issue,³¹ but was featured in several of the studies included in this review. Sanders³² argues that paying sex workers for gaining access to information about their life experiences is similar to the situation of a client paying the sex worker for gaining access to their body and is highly exploitative. However, Maher³³ contends that providing modest remuneration is only fair practice and one that encourages participation. In previous research with socially and economically marginalised communities, uncompensated studies of sex workers often bias the sample towards more privileged and more politically engaged individuals than studies which offer recompense. This is important as much sex work is driven by economic survival.³⁴

Alongside fear of exploitation, community participation in research can be constrained by scepticism that the research will not result in any direct benefit.³⁵ The underuse of participatory methodologies often means that available studies are often conceptualised without relevant engagement with sex working communities or organisations, resulting in the production of research which targets the perceived rather than actual needs of sex workers.

Participatory methodologies along with PPI help to address some of these ethical issues in their ability to develop research that adequately addresses the needs of sex workers while also safeguarding participants from exploitation.^{35 36} Attempts to engage sex workers in the design of research should not be tokenistic or used to legitimise research, but instead should focus on developing methodologies and equitable partnerships that meet the needs of sex working communities.

The evidence for peer health initiatives²⁰ further highlights the importance of community-based responses that prioritise the engagement of target

populations in the development and delivery of support programmes.³⁷ Collective processes, initiated by this engagement, are experienced as helping to create a community voice capable of social and behavioural changes, including improved awareness and access to support services.²⁰ The findings presented in this review provide some proof of concept that peer support is effective in hidden and stigmatised populations³⁸ who do not otherwise engage well with healthcare providers. Peer education may provide the opportunity for sex workers to become authentic educators,³⁹ not only in their community but across public service organisations.

Potential biases of the review process

The scope of this review focused on psychosocial interventions which were evaluated using only validated health and well-being outcome measures, which are considered to be at a reduced risk of bias compared with self-reported measures.⁴⁰ This may have led to the exclusion of some studies where interventions were less rigorously evaluated, but still experienced as beneficial to overall well-being by participants. Disaggregation of data will have also contributed to the exclusion of studies where sex work is reported within sample characteristics, but not presented separately within research findings.

CONCLUSIONS

This systematic review identifies a gap in the evidence base around the effectiveness of psychosocial interventions to support the well-being of sex workers. Available studies are weak in their design and lack generalisability beyond female street-based sex workers. Smartphone-based diaries such as EMA provides some evidence of promoting intentions for behaviour change but, as with other approaches, would benefit from having some focus around how such interventions create change. Finally, peer health initiatives developed in consultation with sex workers offer promise but warrant further investigation.

Additional Educational Resources

- ▶ The nature and prevalence of prostitution and sex work in England and Wales today. Professor Marianne Hester *Prostitution_and_Sex_Work_Report.pdf* (publishing.service.gov.uk)
- ▶ Sex work and mental health: reviewing the occupational risks of sex workers in comparison to other 'risky' professions. Professor Teela Sanders *Microsoft Word - briefing paper final.docx* (le.ac.uk)
- ▶ Guiding principles for best practice research with sex workers. Samuel Brookfield *Barriers to Accessing Sexual Health Services for Transgender and Male Sex Workers: A Systematic Qualitative Meta-summary* | SpringerLink

Twitter Jane Meyrick @DrJaneMeyrick

Acknowledgements The authors would like to acknowledge the assistance of Hannah Poore, specialist subject librarian at the University of the West of England Bristol, and Dr Lucy Platt from the London School of Hygiene and Tropical Medicine.

Contributors KT and JM initiated the study and defined the research question. All authors contributed to developing the methodology. KT carried out all literature reviews. KT, DM and LS acted as data reviewers and extractors. JM provided reconciliation in the event of any discrepancies. KT drafted the protocol and the manuscript that was later edited by JM.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Disclaimer The findings and conclusions of this review are those of the authors and do not represent the official position of the University of the West of England Bristol or other institutions with which the authors are affiliated.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iDs

Kevin Turner <http://orcid.org/0000-0002-6657-1477>
Jane Meyrick <http://orcid.org/0000-0002-2510-9251>
Danny Miller <http://orcid.org/0000-0003-0707-4830>

REFERENCES

- 1 Jeal N, Macleod J, Salisbury C, *et al.* Identifying possible reasons why female street sex workers have poor drug treatment outcomes: a qualitative study. *BMJ Open* 2017;7:e013018.
- 2 Overs C, Hawkins K. Can rights stop the wrongs? Exploring the connections between framings of sex workers' rights and sexual and reproductive health. *BMC Int Health Hum Rights* 2011;11:S3–6.
- 3 Nitschke H, Oliveira F. P2.165 10 Years of experience: comprehensive sexual health services for female sex workers (FSW) in Cologne. *Sex Transm Infect* 2013;89:A138.2–A138.
- 4 Callender D, Bell S, Minichiello V. P712 "You need a cash buffer": male sex work and condom use in the era of HIV pre-exposure prophylaxis. *Sex Transm Infect* 2019;2019:A309–10.
- 5 Mukandavire Z, Vickerman P. P3.400 Comparing the utility of condom and HIV pre-exposure prophylaxis (PreP) use among FSWs. *Sex Transm Infect* 2013;89:A274.2–A274.
- 6 Evans S, Malek R, King G. Hepatitis B vaccination in male sex workers. *HIV Med* 2014;15:148.
- 7 Benoit C, Jansson SM, Smith M, *et al.* Prostitution stigma and its effect on the working conditions, personal lives, and health of sex workers. *J Sex Res* 2018;55:457–71.
- 8 Koken JA. Independent female escort's strategies for coping with sex work related stigma. *Sex Cult* 2012;16:209–29.
- 9 Buttram ME, Surratt HL, Kurtz SP. Resilience and syndemic risk factors among African-American female sex workers. *Psychol Health Med* 2014;19:442–52.
- 10 Chan A-W. Out of sight but not out of mind: how to search for unpublished clinical trial evidence. *BMJ* 2012;344:d8013.
- 11 Harcourt C, Donovan B. The many faces of sex work. *Sex Transm Infect* 2005;81:201–6.

- 12 Thomas BH, Ciliska D, Dobbins M, *et al.* A process for systematically reviewing the literature: providing the research evidence for public health nursing interventions. *Worldviews Evid Based Nurs* 2004;1:176–84.
- 13 National Institute for Health and Care Excellence (NICE). *Methods for the development of NICE public health guidance*. London: NICE, 2012.
- 14 Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009;62:1006–12.
- 15 Gunn JKL, Roth AM, Center KE, *et al.* The unanticipated benefits of behavioral assessments and interviews on anxiety, self-esteem and depression among women engaging in transactional sex. *Community Ment Health J* 2016;52:1064–9.
- 16 Arnold EA, Stewart JC, McNeece CA. The psychosocial treatment needs of street-walking prostitutes: perspectives from a case management program. *J Offender Rehabil* 2000;30:117–32.
- 17 Preble KM, Praetorius RT, Cimino A. Supportive exits: a best practices report for a sex worker intervention. *J Hum Behav Soc Environ* 2016;26:162–78.
- 18 Wahab S. Evaluating the usefulness of a prostitution diversion project. *Qualitative Social Work* 2006;5:67–92.
- 19 Decker MR, Tomko C, Wingo E, *et al.* A brief, trauma-informed intervention increases safety behavior and reduces HIV risk for drug-involved women who trade sex. *BMC Public Health* 2018;18:75.
- 20 Benoit C, Belle-Isle L, Smith M. Sex workers as peer health advocates: community empowerment and transformative learning through a Canadian pilot program. *Int J Equity Health* 2017;2017:160.
- 21 Bodkin K, Delahunty-Pike A, O'Shea T. Reducing stigma in healthcare and law enforcement: a novel approach to service provision for street level sex workers. *Int J Equity Health* 2015;14:35.
- 22 Litchfield J, Maronge A, Rigg T, *et al.* Can a targeted GP-led clinic improve outcomes for street sex workers who use heroin? *Br J Gen Pract* 2010;60:514–6.
- 23 Felsher M, Wiehe SE, Gunn JKL, *et al.* 'I got it off my chest': an examination of how research participation improved the mental health of women engaging in transactional sex. *Community Ment Health J* 2018;54:171–9.
- 24 Meyrick J. What is good qualitative research? A first step towards a comprehensive approach to judging rigour/quality. *J Health Psychol* 2006;11:799–808.
- 25 Spice W. Management of sex workers and other high-risk groups. *Occup Med* 2007;57:322–8.
- 26 Arnold EM, Stewart JC, McNeece CA, *et al.* Understanding violence by female street walking prostitutes. *Violence Vict* 2001;16:145–59.
- 27 Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence. *Psychol Bull* 2003;129:674–97.
- 28 Daly W. Law enforcement in Times Square, 1970's - 1990's. In: McNamara R, ed. *Sex, scams, and street life: The sociology of New York City's Times Square*. Westport: Praeger, 1995: 97–106.
- 29 Davis N. *Prostitution: an international handbook on trends, problems, and policies*. Westport: Greenwood Press, 1993.
- 30 Kloek M, Dijkstra M, Jonker M. THAD0103 Violence towards sex workers in the Netherlands. *Journal of the International AIDS society* 2018;21:e252148.
- 31 Bauermeister JA, Pingel E, Zimmerman M. Data quality in HIV/AIDS web-based surveys: handling invalid and suspicious data. *Field Methods* 2012;24:272–91.
- 32 Sanders T. Researching sex work: dynamics, difficulties, and decisions. In: Hobbs D, Wright R, eds. *The SAGE handbook of fieldwork*. London: Sage, 2006: 202–21.
- 33 Maher L. *Sexed work: gender, race and resistance in a Brooklyn drug market*. Oxford, UK: Oxford University Press, 2000.
- 34 Brents BG, Sanders T. Mainstreaming the sex industry: economic inclusion and social ambivalence. *J Law Soc* 2010;37:40–60.
- 35 Jeffreys E. Sex worker-driven research: best practice ethics. challenging politics: critical voices, 2010. Available: <https://www.nswp.org/resource/sex-worker-drivenresearch-best-practice-ethics> [Accessed 14 Mar 2021].
- 36 Shaver FM. Sex work research: methodological and ethical challenges. *J Interpers Violence* 2005;20:296–319.
- 37 Benoit C, Carroll D, Chaudhry M. In search of a Healing Place: Aboriginal women in Vancouver's Downtown Eastside. *Soc Sci Med* 2003;56:821–33.
- 38 Broadhead R, Heckathorn D, Weakliem D. Harnessing peer networks as an instrument for AIDS prevention: results from a peer-driven intervention. *Public Health Rep* 1998;113:42–57.
- 39 Moore L, Chersich MF, Steen R, *et al.* Community empowerment and involvement of female sex workers in targeted sexual and reproductive health interventions in Africa: a systematic review. *Global Health* 2014;10:47.
- 40 Blank L, Baxter SK, Payne N, *et al.* Systematic review and narrative synthesis of the effectiveness of contraceptive service interventions for young people, delivered in health care settings. *Health Educ Res* 2012;27:1102–19.