

Intracervical local anaesthesia for IUD fitting

I am grateful for the endorsement in respect of intracervical local anaesthesia (LA) for intrauterine device (IUD) fitting that has been offered by both medical and nursing colleagues.^{1,2}

Intracervical LA is easy to do, easy to teach and can make a very significant difference to both the procedure and the patient experience. While it may not be necessary for all, my contention is that it is a technique that should be both available and offered to all women contemplating an IUD fitting. I would agree with Ann Eady² that this can be available to nurse fitters just as to doctors.

I remain uncomfortable with the ethics of not being prepared or able to offer an option that will reduce pain if this can

be anticipated. I propose that LA should become an integral element to the practical aspect of training. I do however feel that the e-SRH module 18 pertaining to this needs some modification and would ask that this be considered.

Sarah Gray, MRCGP, DFRH

General Practitioner, Primary Care Lead in Women's Health, NHS Cornwall and Isles of Scilly, Callington Health Centre, Callington, UK;
sarah.gray@ciospct.cornwall.nhs.uk

Competing interests None

J Fam Plann Reprod Health Care 2011;**37**:254–255.
doi:10.1136/jfprhc-2011-100198

REFERENCES

- 1 **Hutt S.** Option of local anaesthetic for IUD fittings: author's response [Letter]. *J Fam Plann Reprod Health Care* 2011;**37**:190.
- 2 **Eady A.** Local anaesthetic administration for IUD/IUS insertion [Letter]. *J Fam Plann Reprod Health Care* 2011;**37**:254.