

F

A

C

T

Faculty

Aid to

CPD

Topics

A CPD Self-Assessment Test

QUESTION SHEET

Review No. 2001/03

To be reviewed not later than 31 May 2006

Resuscitation in the family planning and reproductive health care setting

Indicate your answer by ticking the appropriate box for each question	True	False
1. Cervical stimulation produces an anaphylactic reaction.	<input type="checkbox"/>	<input type="checkbox"/>
2. Bradycardia is defined as a ventricular rate below 60 b/min.	<input type="checkbox"/>	<input type="checkbox"/>
3. Atropine is the drug of choice for reversing bradycardia.	<input type="checkbox"/>	<input type="checkbox"/>
4. Signs of ‘cervical shock’ include pallor, sweating, and tachycardia.	<input type="checkbox"/>	<input type="checkbox"/>
5. If ‘cervical shock’ occurs, the first line of action is to raise the legs and lower the head.	<input type="checkbox"/>	<input type="checkbox"/>
6. Basic life support should commence immediately after cessation of breathing.	<input type="checkbox"/>	<input type="checkbox"/>
7. HIV has been transmitted to person giving rescue breathing.	<input type="checkbox"/>	<input type="checkbox"/>
8. Survival is rare if defibrillation and/or drug therapy is unavailable within 30 minutes of cardiac arrest.	<input type="checkbox"/>	<input type="checkbox"/>
9. IV adrenaline is the drug of choice for anaphylaxis.	<input type="checkbox"/>	<input type="checkbox"/>
10. Current adult basic life support guidelines recommends a compression to ventilation ratio of 15:2.	<input type="checkbox"/>	<input type="checkbox"/>

Turn to page 173 for answers