Low-calorie sweeteners and the risk of preterm delivery: results from two studies and a meta-analysis

Carlo La Vecchia

Head, Department of Epidemiology, Istituto di Ricerche Farmacologiche Mario Negri, Milan and Professor, Department of Clinical Sciences and Public Health, Università degli Studi di Milano, Milan, Italy

Correspondence to

Professor Carlo La Vecchia, Department of Epidemiology, Istituto di Ricerche Farmacologiche Mario Negri, Via Giuseppe La Masa, Milan 19-20156, Italy; carlo.lavecchia@marionegri.it

Received 1 November 2012

Revised 6 November 2012

Accepted 7 November 2012

BACKGROUND

The possible relation between low-calorie sweeteners (as well as sugar-sweetened beverages) and unfavourable pregnancy outcomes, in particular preterm delivery and low birth weight, has recently become of considerable interest, given the widespread use of low-calorie sweeteners, and the high frequency of preterm deliveries in high-income countries.

NORWEGIAN COHORT DATA

In a study from Norway, Englund-Ogge *et al.*¹ considered the possible relation between low-calorie-sweetened and sugar-sweetened beverages and preterm delivery using data from 60 761 women in the Norwegian Mother and Child Study cohort. Overall, 3281 preterm deliveries (<37 weeks) were observed, corresponding to 5.4% of all deliveries. Of these, 3.9% were late preterm deliveries (34 to <37 weeks), 0.8% moderately preterm deliveries (32 to <34 weeks) and 0.7% early preterm (<32 weeks) deliveries.

There was a moderate association between frequency of use of sugar-sweetened beverages during pregnancy and the risk of preterm deliveries. As compared to women who did not drink sugar-sweetened beverages, the relative risk (RR) was 1.15 for <1 serving/week, 1.15 for 1–6 servings/week, 1.25 for 1 serving/day, 1.19 for 2–3 servings/day and 1.41 for ≥4 servings/day. Allowance was made for age, education, obstetric history and tobacco but not for alcohol. Women with diabetes were excluded. Although the test for trend was significant ($p=0.008$), a statistically significant association was already observed for sporadic use (<1 serving/week), and there was no linear dose–risk relation, since the RR was similar for <1 serving/week (1.15) and 2–3 servings/day (1.19).

This shed doubts on the possibility of a real association between sugar-sweetened beverages and the risk of preterm delivery.

The association with preterm deliveries was, if anything, even less consistent for low-calorie or ‘artificially sweetened’ beverages. Compared to non-users of artificially sweetened beverages, the RR was 1.01 for <1 serving/week, 1.09 for 1–6 servings/week, 1.20 for 1 serving/day, 1.01 for 2–3 servings/day and 1.12 for ≥4 servings/day. The test for trend was formally non-significant ($p=0.053$), and there was no linear dose–risk relation; that is, the RR was 1.01 for both women reporting sporadic use (<1 serving/week) and for those who exhibited regular use of 2–3 drinks/day.

DANISH COHORT DATA

The issue of potential side effects of low-calorie sweeteners in pregnancy, with specific focus on the risk of preterm delivery (<37 weeks), was originally raised in 2010 by Halldorsson *et al.*² from a study based on the Danish National Birth Cohort (1996–2002), including 59 334 women and a total of 2739 cases of preterm births. The original report presented the results stratified by carbonated and non-carbonated drinks only, and gave a RR of 1.78 [95% confidence interval (CI) 1.19–2.66] for drinkers of ≥4 servings/day of ‘artificially sweetened carbonated soft drinks’. This RR estimate was, however, based on 27 cases only (i.e. less than 1% of the cases), a likely selected sub-population. This subgroup may well have other baseline characteristics that influence its risk of preterm birth, and some residual confounding is likely to be present also after multivariate analysis. There was moreover an appreciable difference between carbonated and non-carbonated drinks. The RR, in fact, was

To cite: La Vecchia C. *Journal of Family Planning and Reproductive Health Care* 2013; **39**, 12–13.

appreciably lower (1.29) among heavy drinkers of 'artificially sweetened non-carbonated soft drinks' (105 cases), and not materially different from that of heavy drinkers of 'sugar-sweetened non-carbonated soft drinks' (RR 1.16, 102 cases). When carbonated and non-carbonated low-calorie drinks were combined, there was no material excess risk up to 2 drinks/day (RR 1.09).³

Any possible effect of 'sweetened' versus 'artificially sweetened' drinks – as well as of carbonated versus non-carbonated drinks – on low birth weight in that study was therefore limited and inconsistent, confounded by the two types of drinks (carbonated and non-carbonated),⁴ as well as by residual confounding due to other likely differences in baseline characteristics of pregnant women drinking low-calorie versus sweetened, and carbonated versus non-carbonated drinks.

There is, moreover, a lack of biological consistency for any possible role of low-calorie sweeteners on the risk of low birth weight. Halldorsson *et al.*² refer to a possible role of caffeine in hypertension and consequently in preterm delivery,⁵ but – if real – any effect of caffeine should be shared by sweetened and low-calorie drinks. Several other standard criteria for inference in observational studies (the Hill criteria), including strength of the association, dose–risk relation, time–risk relation and consistency with other reports, also remain open to discussion.^{6–7} A critical discussion of the application of the Hill criteria^{6–7} can be put forward with reference to the Danish cohort, as for most observational studies.

META-ANALYSIS

We pooled the main findings of the two studies described above using standard meta-analytic techniques.⁸ The Danish data were introduced in two separate strata of carbonated and non-carbonated beverages, since any possible association was apparently stronger for carbonated beverages, both sugar-sweetened and low-calorie ones.

The results of the meta-analysis are given in Table 1. The pooled RRs in all levels of consumption were close to unity both for sugar-sweetened and for low-calorie beverages and, most important, there was no difference in the risk estimates of sugar and low-calorie beverages, since all the CIs largely overlap.

There is therefore no evidence that low-calorie beverages, which are US Food and Drug Administration approved for consumption during pregnancy, have an

Table 1 Combined analysis (meta-analysis) of the main findings of the studies by Halldorsson *et al.*² and Englund-Ogge *et al.*¹ on the association between sugar-sweetened, low-calorie-sweetened beverages and the risk of preterm delivery

Beverage consumption	Relative risk (95% CI)	
	Sugar-sweetened	Low-calorie-sweetened
Non-users	1*	1*
<1 serving/week	0.97 (0.91–1.05)	1.05 (0.98–1.12)
1–6 servings/week	0.93 (0.87–0.99)	1.09 (1.02–1.16)
1 serving/day	1.03 (0.93–1.14)	1.18 (1.06–1.32)
2–3 servings/day	1.06 (0.96–1.16)	1.14 (1.02–1.26)
≥4 servings/day	1.23 (1.06–1.42)	1.25 (1.09–1.43)

The Halldorsson *et al.* study examined carbonated and non-carbonated drinks pooled from separate strata.

*Reference category.

CI, confidence interval.

impact on preterm delivery at any variance from that of sugar-sweetened beverages.

Funding None.

Competing interests The author has received in the past unconditional grant support from the International Sweeteners Association, Brussels, Belgium for addressing the issue of sweeteners and cancer risk.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Englund-Ogge L, Brantsaeter AL, Haugen M, *et al.* Association between intake of artificially sweetened and sugar-sweetened beverages and preterm delivery: a large prospective cohort study. *Am J Clin Nutr* 2012;96:552–529.
- 2 Halldorsson TI, Strom M, Petersen SB, *et al.* Intake of artificially sweetened soft drinks and risk of preterm delivery: a prospective cohort study in 59,334 Danish pregnant women. *Am J Clin Nutr* 2010;92:626–633.
- 3 Halldorsson TI. Intake of artificially sweetened soft drinks and risk of preterm delivery. *Am J Clin Nutr* 2010;92:1540; author reply 1540–1542.
- 4 La Vecchia C. Intake of artificially sweetened soft drinks and risk of preterm delivery. *Am J Clin Nutr* 2010;92:1540.
- 5 Winkelmayr WC, Stampfer MJ, Willett WC, *et al.* Habitual caffeine intake and the risk of hypertension in women. *JAMA* 2005;294:2330–2335.
- 6 Bursey RG, Watson ML. Intake of artificially sweetened soft drinks and risk of preterm delivery. *Am J Clin Nutr* 2010;92:1277–1278; author reply 1278–1280.
- 7 Hill AB. The environment and disease: association or causation? *Proc R Soc Med* 1965;58:295–300.
- 8 Greenland S. Quantitative methods in the review of epidemiologic literature. *Epidemiol Rev* 1987;9:1–30.