An alternative method of anaesthesia for implant insertion: description of a clinical initiative in contraceptive care

Julia Shefras,¹ Annabel Forsythe,² Fatima Nathani,³ Donna Hoffman⁴

¹Consultant in Contraception and Sexual Health, C&SH Oxford Health NHS Foundation Trust, The Alec Turnbull Centre, Oxford, UK

²Specialist Trainee in Community Sexual & Reproductive Health, Department of Contraception & Sexual Health, Oxford Health NHS Foundation Trust, Oxford, UK

³Consultant in Contraception and Sexual Health, Russells Hall Hospital, Dudley Group of Hospitals, Dudley, UK

⁴Clinical Nurse Specialist in Contraception and Sexual Health, Oxford Health NHS Foundation Trust, Oxford, UK

Correspondence to

Dr Julia Shefras, Department of Contraception & Sexual Health, Oxford Health NHS Foundation Trust, The Alec Turnbull Centre, 23 Between Towns Road, Oxford OX4 3JH, UK; julia.shefras@oxfordhealth.nhs.uk

Received 16 July 2013

Revised 10 December 2013

Accepted 12 December 2013

Published Online First

5 February 2014

WHAT IS THE CURRENT ANAESTHETIC PRACTICE FOR IMPLANT INSERTIONS?

Adequate anaesthesia is an important procedural step when inserting contraceptive implants. Whilst the product information for Nexplanon[®] describes a choice of anaesthetic spray or injection for insertion,¹ the Faculty of Sexual & Reproductive Healthcare (FSRH) does not mention spray, solely recommending use of lidocaine injection.² Therefore anaesthetic spray such as ethyl chloride, despite being a recognised technique for implant insertions, is not widely used.

WHAT IS ETHYL CHLORIDE SPRAY?

Ethyl chloride spray acts as a local anaesthetic when sprayed onto the skin. It is a fast acting vapo-coolant with minimal side effects and it can be safely used for sterile procedures with no additional risk of bacterial contamination.³ It is a long-established analgesic technique for insertion of cannulas and needles, particularly in paediatric care.

WHY IS AN ALTERNATIVE ANAESTHESIA NEEDED?

Lidocaine injection provides effective anaesthesia, but may cause some discomfort during injection. It involves the cost, storage and management of needles, syringes, sharps disposal and local anaesthetic. Furthermore, the time taken to gather and prepare these items and then dismantle the trolley represents the greater part of the overall process.

The aim of this project was to evaluate whether use of ethyl chloride spray as an alternative anaesthetic for insertion of contraceptive implants is an innovation offering improvement for patients, increased productivity for the service and

potentially cost savings in comparison to use of lidocaine.

HOW WAS THE CHANGE IMPLEMENTED?

During a 6-month period commencing January 2012, 4 of 16 implant fitters in the Oxford Health C&SH service offered a choice of spray or lidocaine injection for implant insertions. During this time, an evaluation of patients' experiences was performed. Alongside this a comparative evaluation of the two methods by all staff fitting implants was also undertaken in a non-clinical environment. Following the results, training in the technique was cascaded to all implant fitters and spray is now offered by all fitters within Oxford Health C&SH services.

HOW HAVE PATIENTS FOUND THE USE OF ETHYL CHLORIDE SPRAY?

Four clinicians trained in use of ethyl chloride spray saw 25 clients for implant insertions. They offered clients the choice of spray or injection for the insertion. Afterwards the clients were invited to complete a questionnaire.

The clients were in the age range 14–48 (median 22) years. Ten had used an implant in the past, inserted with local anaesthetic injection. Fifteen were having an implant fitted for the first time. Four clients chose to have the injection (two first-time users and two previous users).

Twenty-three clients felt the procedure was comfortable, two felt it was uncomfortable (one spray, the other injection). Comfort ratings ranged from 2 to 10 (average, 7.6) with 10 being the best health state, 0 being the worst health state. Twenty-three clients said they would recommend the procedure to a

To cite: Shefras J, Forsythe A, Nathani F, et al. *J Fam Plann Reprod Health Care* 2014;**40**:226–228.

Comments from clients included “felt fine”, “fantastic”, “painless and quick” and “use spray!”. There was one negative comment “it hurt” by a client who had spray and was one of the two who said the procedure was uncomfortable.

Before training was cascaded throughout the service all fitters were invited to participate in a comparative evaluation of anaesthesia in a non-clinical environment. Each participant was shown two photographs (Figures 1 and 2) that demonstrated the items

Following the initial staff evaluation and patient survey, a procedure was developed and the technique was taught to all staff fitting implants within the service.

Based on current practice within the service, we estimate a minimum saving of £117.80/1000 patients using ethyl chloride spray in place of lidocaine injection.

The relative speed of using spray may encourage more quick-starting to take place, facilitating one-stop care and fewer second visits. Although initial savings based on equipment are modest, savings on a grander scale may be predicted if it helps to reduce follow-up appointments and possibly unintended pregnancies.

Table 1 Staff comments, concerns and suggestions about use of ethyl chloride spray

Positive comments	Negative comments/concerns	Suggestions
Easier	Anxious about having to rush insertion before spray wears off	Would like guidance from FSRH
Time efficient	Worried about not giving enough spray	Would like spray to be readily available
Cost efficient	Can frozen skin restrict glide?	Would like a protocol
Less risk of needlestick	Lidocaine gives longer to chat to patient and type up notes	
Appears less painful for patient		
Good for the needle phobic		

FSRH, Faculty of Sexual & Reproductive Healthcare.

WHAT COULD THE FUTURE HOLD?

Use of ethyl chloride spray has potential to improve both quality of care, patient choice and productivity. It may offer additional clinical advantages and cost savings.

Lidocaine remains the anaesthetic of choice for implant removals due to the need for a longer duration of action for this procedure.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Organon Laboratories Ltd. Nexplanon – Information for healthcare professionals. Last approved July 2012.
- 2 Faculty of Sexual & Reproductive Healthcare Clinical Effectiveness Unit. *Progestogen-only Implants*. April 2008 (updated January 2009). <http://www.fsrh.org/pdfs/CEUGuidanceProgestogenOnlyImplantsApril08.pdf> [accessed 2 June 2013].
- 3 Acorus Therapeutics Ltd. Ethyl Chloride EEC Safety Data Sheet. 2001 (updated 2007).