Faculty Aid to CPD Topics

A CPD Self-Assessment Test

REVIEW

This review is intended as an educational exercise and reports the personal views of the authors

Review No. 2000/03

To be reviewed not later than 31 March 2005

Subdermal contraceptive implants

Emeka Oloto, FWACS, MRCOG, MFFP

*Specialist Registrar in Obstetrics and Gynaecology, St. James's University Hospital, Beckett Street, Leeds LS9 7TF, UK.
SCMO, Leeds Community and Mental Health NHS Trust, Leeds, UK*

Lawrence Mascarenhas, MD, MRCOG, MFFP, M.Ed. (Cantab)

Consultant Obstetrician and Gynaecologist, Department of Obstetrics and Gynaecology, C Floor, East Block, Queens Medical Centre, Nottingham NG7 2UH, UK

(Accepted April 22 2000)

How to use a FACT

A FACT is an up-to-date review of a subject relevant to the speciality, intended to help you fulfil your CPD requirements in your home or place of work. Whilst FACTs are edited and reviewed at various levels within the Faculty, the actual contents and views expressed are those of the authors and not the Faculty. More specifically, these reviews are not guidelines. The CEC is producing clinical guidelines separately.

FACTs have three sections: a review, a true/false test, and discussion points. To use a FACT to earn CPD credits you should do the following:

1. **Working alone:** Read the review and do the test. The answers are provided on page 183 so you can mark yourself. If there are points you are unsure about, disagree with, or need further clarification on, make a note of these for use at a later date. This should take you no more than 1 hour. Keep a record of having done this in your CPD diary and, unless indicated otherwise on the FACT, this will earn you 1 hour (DFFP), 1 credit (MFFP).
2. **Working as a group:** arrange a meeting of at least 1 hour with colleagues to discuss the discussion points given in the FACT (page 174) and any issues the participants have come up with as a result of reading the FACT. Keep a record of having done this in your CPD diary and, unless indicated otherwise on the FACT, this will earn you 1 hour (DFFP), 1 credit (MFFP).

Introduction

Subdermal contraceptive implants are an additional approach to meeting the world-wide need for more effective and acceptable birth control. They provide long-acting, highly effective and reversible contraception by achieving low and stable concentrations of synthetic progestogens and require little user compliance, thereby achieving user failure rates that are similar to the method failure rates.^{1,2}

History and development

Subdermal contraceptive implants may be divided into two main groups: non-biodegradable, some of which are now widely used, and biodegradable, which remain under development in clinical trials. These are summarised in Table 1. This review will focus on systems that are in use in the UK (Norplant® and Implanon®) or nearing general availability (Norplant II®).

The development of subdermal contraceptive implant systems usually involves a wide range of materials and approaches. Nearly all of the synthetic progestogens have been used in these systems, but levonorgestrel is the most widely used because of its high potency per volume.²

Non-biodegradable systems make use of a suitably inert carrier for the contraceptive steroid, from which it diffuses over time e.g. polydimethylsiloxane (Silastic®) or ethylene vinyl acetate (EVA). Norplant® is a first generation implant system that was developed as a result of the studies by Segal and Croxatto showing that different steroids were released from subdermally inserted Silastic® capsules at different rates, and that the rate of release increased proportionately to the surface area of the capsules, and decreased with increasing wall thickness.³ Norplant II®, which consists of two silastic-covered rods and Implanon®, which consists of one EVA rod, are classed as second

Table 1 Subdermal contraceptive implants currently in use or under development

Implant	Number	Progestogen (per capsule or rod or pellet)	Use (years)	Phase of development
Non-biodegradable:				
Norplant®	6 C	Levonorgestrel 36 mg	5	Marketed*
Norplant II® (Jadelle®)	2 R	Levonorgestrel 70 mg	5	Marketed
Sino implant I®	6 C	Levonorgestrel 36 mg	5	Marketed
Sino implant II®	2 R	Levonorgestrel 75 mg	3–5	Marketed
Implanon®	1 R	Etonogestrel 68 mg	3	Marketed**
Nestorone (ST-1435)	1 C	Nor-progesterone 40/78 mg	2	I-II
Uniplant or Surplant	1 R	Nomegestrel acetate 38 mg	1	I
Biodegradable:				
Capronor	1 C	Levonorgestrel (12 or 21.6 mg)	1	II
Annuelle	4 P	Norethindrone 90%; Cholesterol 10%	1	I

C=Capsule; R=Rod; P=Pellet; *=Marketed in the UK **=Currently marketed in the UK

generation subdermal contraceptive implant systems. They were developed in an attempt to reduce the problems of insertion and removal associated with the six-implant system (Norplant®). Other reasons that motivated further development of implants included the need for them to be less obvious under the skin; the need for implants to degrade and not require removal at all; the need to use progestogens that are less androgenic than levonorgestrel so that acne, weight gain, and perhaps other side effects would be less common; and the need to use simple, disposable devices to speed insertion and reduce the risk of blood-borne disease transmission.^{4,5} Norplant II® received Federal Drug Agency (FDA) approval in 1996 and is marketed in Finland, while Implanon® has been approved by the European Drug Licensing Agency (EDLA) and is currently marketed in Australia, Indonesia and 11 European countries.

Scientific basis for sustained-release contraceptives

The major aims of non-oral routes of contraceptive steroids include long duration of action, continuous and steady steroid blood levels and avoidance of the ‘first pass’ peak effect through the hepatic portal system, thus avoiding the peak plasma levels of steroids associated with oral dosing without compromising contraceptive efficacy.¹ The higher bioavailability leads to lower doses of steroids being administered, with a possible parallel reduction in the occurrence of adverse effects.

Mechanism of action

The mechanisms of action include: cervical mucus blockade and prevention of sperm penetration;⁶ suppression of oestradiol-induced cyclic maturation of the endometrial lining, causing hypotrophic changes;⁶ and ovarian effects ranging from disordered luteal phase through complete anovulation to persistent follicles.⁶ However, Implanon® was designed to achieve complete inhibition of ovulation during its period of action.⁷

Advantages and disadvantages

The efficacy of these methods is high: Norplant® has a cumulative failure rate of 1 per 100 users after 5 years of use (0.26 per 100 woman years);⁸ Norplant II® has a similar cumulative failure rate after 3 and 5 years of use of 0.8 per 100 users⁹ and combined data from 13 studies with Implanon® indicated that no pregnancies occurred in 1716 women treated - 53 530 cycles (4103 women years), resulting in a Pearl Index of 0.0 (95% CI, 0.00 - 0.09).⁷ Based on the principle of informed choice and ability to

discontinue method at any time for whatever reason, they require little user compliance or motivation, resulting in use-effectiveness rates which closely approximate theoretical effectiveness. In addition it may be used by women who have contraindications to oestrogens.² Furthermore, fecundity returns immediately following removal.²

On the other hand, insertion and removal of contraceptive implants are minor surgical procedures that require trained personnel and with risks similar to any minor surgical procedure (infection, bleeding or haematoma). Women cannot initiate or discontinue the method themselves, and may encounter difficulty in finding personnel trained in removal techniques. Finally, the disruption of menstrual pattern frequently associated with contraceptive implant use, and the visibility of the capsules in some cases, may be unacceptable to women or their partners.^{2,10}

Insertion and removal

Contraceptive implants are inserted subdermally, superficially enough to be palpated. Users of non-biodegradable implants must be able to have them removed whenever they want, for whatever reason, or at the end of their duration of contraceptive action, with minimal inconvenience or discomfort.

Insertion and removal of implants is performed under local anaesthesia, which is uncomplicated in the majority of cases. However, some problems associated with insertion and removal have been reported.^{11,12} Incorrect insertion makes removal much more difficult, which stresses the importance of training.² Difficult Norplant® removals, using the standard technique as recommended by the manufacturers, have prompted some researchers to develop new methods for removal.^{13,14}

The Faculty of Family Planning and Reproductive Health Care offers a training framework, which covers both the theoretical and practical aspects of subdermal contraceptive implant insertion and removal. Further details of this framework can be obtained from the Faculty.

Implanon® has several advantages over Norplant® with respect to insertion and removal: it is a single rod; the EVA carrier has a semi-rigid consistency facilitating removal, and finally it is conveniently preloaded within a sterile, disposable applicator.¹⁵ A meta-analysis of the data from seven prospective randomised studies in 1378 women showed that it was approximately four times quicker to insert and remove Implanon® as compared to Norplant® (mean insertion times 1. 1 Vs 4.3 minutes, respectively;

mean removal times 2.65 Vs 10.2 minutes, respectively).¹⁵ Furthermore, Implanon® was associated with a significantly lower incidence of removal complications (0.2% Vs 4.8% respectively; $P < 0.001$).¹⁵ Finally, the mean removal time for Norplant II® reported in another study was 4.9 minutes, i.e. twice that reported with Implanon® and half that required for Norplant®.¹⁶

Implanon® is licensed for 3 years of continuous usage. The manufacturer's datasheet suggests that Implanon® may be less effective during the third year of use in women weighing more than 70 kg. As there is no scientific evidence to support this, it is our normal practice not to treat women weighing more than 70 kg any differently from other women using this method.

Side effects and management

Subdermal contraceptive implants share common side effects, the most common of which is menstrual disruption^{1,2} which is also the most common reason for discontinuation of the method and presents a major obstacle to its widespread use.¹⁷ Because the changes in menstrual bleeding pattern vary widely, it is not possible to predict the change a particular client may experience.

Over a period of 2 years, Implanon® produced a general picture of bleeding pattern incidence that was not dissimilar to that for Norplant®, though amenorrhoea occurred more frequently, at about 30-40% compared with 20-30% for Norplant® between 12-24 months of use.¹⁸ Infrequent bleeding occurred in about 50% of women in the first 3 months, but declined to about 30% after 6 months. The incidence of prolonged bleeding was also higher in the first 3 months than subsequently, when it had an incidence of 10-20%. Frequent bleeding was uncommon, usually less than 10% at any time over 2 years. However, it is important to remember that in untreated women of reproductive age,¹⁹ amenorrhoea occurred in about 1% of women aged about 30 years; infrequent bleeding in about 8%, and frequent and prolonged bleeding in less than 0.1%.

Other reported side effects such as weight changes, headaches, mood changes, abdominal bloating and discomfort, loss of libido, nausea, acne, alopecia etc. affect less than 10% of users.⁶ No significant difference has been reported in the side effect profile between Norplant® and Implanon®.⁷

The pathophysiological mechanisms of the menstrual disturbances are poorly understood. While there has been some progress in the understanding of the mechanisms of normal and abnormal menstruation,²⁰ there is a relative lack of progress in the actual prevention and treatment of the menstrual disruption associated with progestogen-only contraceptives.

The only consensus of opinion appears to be thorough pre-treatment counselling about probable alterations to the menstrual cycle, and subsequent strong reassurance at follow-up visits.^{8,21} Inevitably, some patients will require specific treatment, but surveys of such treatments by experts around the World²² revealed absence of universal treatment criteria and variation in the type, dose and duration of use of medications such as combined oral contraceptives, oestrogens (synthetic or conjugated), progestogens (levonorgestrel, norethisterone, etc.), and non-steroidal anti-inflammatory drugs. Evidence-based management is not available for the other side effects associated with subdermal contraceptive implants.

An important positive message, however, is that continuation rates and user satisfaction levels are high among users who have undergone detailed pre-insertion

counselling, in spite of the fact that these users also report a significant incidence of side effects, probably because the perceived advantages greatly outweigh the nuisance effects.^{2,6,10} With good counselling, 83.4% and 53% continuation rates at 1 year²³ and 5 years,²⁴ respectively, have been reported with Norplant®. The 2-year continuation rate of Implanon® was reported to be 69%.²⁵

The UK experience

Norplant® was licensed in October 1993 and was the only subdermal contraceptive implant system in general use in the UK until September 1999. It is estimated to have been used by about 55 000 UK women.²⁶ World-wide, about six million women have used this implant, which is registered in 60 countries.⁸ The initial clinical experience with the method^{23,27,28} was promising, although negative reports about Norplant® have appeared in the lay and medical press,^{29,30} mostly due to unrealistic method expectations and problems with removal. For economic reasons, the main distributor discontinued marketing Norplant® in the UK in November 1999, although it is still available from other sources.

The single-rod implant, Implanon®, was licensed for use in the UK in September 1999. It is anticipated that this system should have significant advantages in terms of ease of insertion and removal.

Conclusion

In spite of the very low failure rates and high continuation rates, subdermal contraceptive implants have low usage within Europe. They have been found to be effective and acceptable methods of contraception in other countries. Detailed counselling about their advantages and disadvantages should precede insertion of the implants. Post-insertion counselling may also be necessary. Only trained professionals should insert and remove the implants. The main challenge for the future is to find appropriate therapy to alleviate bleeding problems, which are the main reason for the discontinuation of the method.

References

- Mascarenhas L. Long-acting methods of contraception. *British Medical Journal* 1994; **308**: 991-992.
- Darney PD, Klaisle CM, Tanner S, et al. Sustained-release contraceptives. *Current Problems in Obstetrics, Gynaecology and Fertility* 1990; **XIII**(3): 85-125.
- Segal SJ, Croxatto HB. Single administration of hormones for long term control of reproductive function. Presentation at the XXIII Meeting of American Fertility Society, April 14-16, Washington D, C. 1967.
- Darney PD. Hormonal implants: Contraception for a new century. *American Journal of Obstetrics and Gynaecology* 1994; **170**(5): 1536-1543.
- Chez RA. Clinical aspects of three new progestogens: Desogestrel, gestodene, and norgestimate. *American Journal of Obstetrics and Gynaecology* 1989; **160**: 1296-1300.
- The Population Council, Decisions for Norplant® Programs. *Population Report* (K) Series(4), 1992.
- Croxatto HB, Makarainen L. The pharmacodynamics and efficacy of Implanon®: An overview of the data. *Contraception* 1998; **58**(Suppl. 6): 91S-97S.
- Fraser IS, Tiitinen A, Agandi B, et al. Norplant® Consensus Statement and Background Review. *Contraception* 1998; **57**: 1-9.
- Sivin I, Alvarez F, Mishell DRJ, et al. Contraception with two levonorgestrel rod implants. A 5-year study in the United States and Dominican Republic. *Contraception* 1998; **58**(5): 275-282.
- Mascarenhas L, McLean J, Newton P, et al. Acceptability of contraceptive implants (Norplant® and Implanon®) in Birmingham, U.K. *Fertility and Sterility* 2000. (In Press).
- Chang A, Edlich RF, Himel HN. Local reactions at the insertion site of Norplant® contraceptive system. *Journal of Long-term Effects of Medical Implants* 1993; **3**(4): 305-312.
- Crist T, Barnes MR, Whitehurst WC. Difficulty finding and removing a Norplant® system capsule. *North Carolina Medical Journal* 1994; **55**(2): 76.
- Praptoharjo U, Wibowo S. The "U" technique: a new method for Norplant® implants removal. *Contraception* 1993; **48**: 526-536.
- Sarma SP, Hatcher R. The Emory Method: a modified approach to Norplant® implants removal. *Contraception* 1994; **49**: 551-556.
- Mascarenhas L. Insertion and removal of Implanon®. *Contraception* 1998; **58**(Suppl. 6): 79S-83S.
- Sivin I, Viagas O, Compton IE. Clinical performance of a new two rod levonorgestrel implant: a three year randomised study with Norplant® implants as controls. *Contraception* 1997; **55**: 73-80.
- Baird DT, Glasier AF. Menstrual bleeding patterns and contraception. *IPPF Medical Bulletin* 1991; **25**(4): 1-2.
- Edwards JE, Moore A. Implanon®: A review of clinical studies. *The British Journal of Family Planning* 1999; **24**(Suppl. 4): 3-16.
- Belsey EM, Machin D, d'Arcangues C. Task Force on Long-acting Systemic Agents for Fertility Regulation. Menstrual bleeding patterns in untreated women. *Contraception* 1997; **55**: 57-65.
- Fraser IS, Hickey M, Song J. A comparison of mechanisms underlying disturbances of bleeding caused by spontaneous dysfunctional uterine bleeding or hormonal contraception. *Human Reproduction* 1996; **11**(Suppl. 2): 165-178.
- Olotto EJ, Bromham DR. Treatment of menstrual side effects in Norplant® users, recommendations of expert group. *The British Journal of Family Planning* 1995; **21**(Suppl.): 3-5.

22 Nutley T, Dunson TR. Treatment of bleeding problems associated with progestin-only contraceptives: survey results. *Advances in Contraception* 1997; **13**: 419–428.

23 Kirkman RJE, Bromham DR, O'Connor TP, et al. Prospective multicentre study comparing levonorgestrel implants with a combined contraceptive pill: final results. *The British Journal of Family Planning* 1999; **25**: 36–40.

24 Sivin I, Campodonico I, Kiriwat O, et al. The performance of levonorgestrel rod and Norplant® contraceptive implants: a 5 year randomised study. *Human Reproduction* 1998; **13**(12): 3371–3378.

25 Croxatto HB, Urbancsek J, Massai R, et al. A Multicentre efficacy and safety study of the single contraceptive implant Implanon®. Implanon® Study Group. *Human Reproduction* 1999; **14**(4): 976–981.

26 Eaton T, Hoechst Marion Roussel: The use of Norplant® in the UK. September 1998. Personal communication.

27 Mascarenhas L, Lee S, Newton P, et al. First clinical experience with contraceptive implants (Norplant® and Implanon®) in the UK. *British Journal of Obstetrics and Gynaecology* 1996; **102**: 573–574.

28 Peers T, Stevens JE, Graham J, et al. Norplant® implants in the UK: first year continuation and removals. *Contraception* 1996; **53**(6): 345–351.

29 Rogers L. Contraceptive implants face legal challenge. *Sunday Times* 3rd July, 1994.

30 Roberts J. Women in US sue makers of Norplant®. *British Medical Journal* 1994; **309**: 145.

Discussion points

1. Funding of training and service provision for contraceptive implants: whose responsibility is it?

2. Although the ‘Norplant® Action Group’ collapsed, what are the likely consequences on the future provision and utilisation of other contraceptive implants?

3. Management of menstrual disruption associated with the use of contraceptive implants.

4. The ‘migrating implant’: real or imagined phenomenon?

5. Should patients be informed at the end of the life span of the implants?

F
A
C
T

Faculty
Aid to
CPD
Topics

A CPD Self-Assessment Test

QUESTION SHEET

Subdermal contraceptive implants

Indicate your answer by ticking the appropriate box for each question	True	False
1. Norethisterone is the most widely used progestogen for contraceptive implants.	☐	☐
2. Counselling is the best treatment option for the menstrual disturbances associated with contraceptive implants.	☐	☐
3. The rate of release of steroids from contraceptive implants decreases proportionately to the surface area of the capsules, and increases with increasing wall thickness.	☐	☐
4. Norplant® is a six capsule implant system with each capsule containing 36 mg of levonorgestrel.	☐	☐
5. Implanon® contains norgestrel acetate, a second generation progestogen.	☐	☐
6. Mood swings are the most frequent side effect of subdermal contraceptive implants and the most common reason for discontinuation of the method.	☐	☐
7. It is easier to remove the implants than to insert them.	☐	☐
8. The two-rod contraceptive implant system is licensed for use in at least two countries.	☐	☐
9. Capronor, a biodegradable implant, is now in the phase II stage of development.	☐	☐
10. Implanon® uses a semi-rigid EVA carrier.	☐	☐

Turn to Page 183 for answers