

**Statements on funding and competing interests**

*Funding.* None identified.

*Competing interests.* None identified.

**References**

- 1 Kesby GJ, Korda AR. Migration of a Filshie clip into the urinary bladder seven years after laparoscopic sterilisation. *Br J Obstet Gynaecol* 1997; **104**: 379–382.
- 2 Filshie GM, Casey D, Pogmore JR, Dutton AG, Symonds EM, Peake AB. The titanium/silicon rubber clip for female sterilization. *Br J Obstet Gynaecol* 1981; **88**: 655–656.
- 3 Graf AH, Staudach A, Steiner H, Spitzer D, Martin A. An evaluation of the Filshie clip for postpartum sterilization in Austria. *Contraception* 1996; **54**: 309–311.
- 4 Robson S, Kerin J. Recurrence of pelvic abscess associated with a detached Filshie clip. *Aust N Z J Obstet Gynaecol* 1993; **33**: 446–448.
- 5 Tan BL, Chong C, Tay EH. Migrating Filshie clip. *Aust N Z J Obstet Gynaecol* 2004; **44**: 583–584.
- 6 Gupta AN, Chakravarti RN, Majumdar S, Mapa MK, Dhall GI. Pathological changes in Fallopian tubes following three different kinds of occlusive techniques in primates. *Contraception* 1986; **33**: 245–255.

## JOURNAL REVIEW: Contraceptive issues and obesity: a review of three recent publications

Reviewed by **Claire I Alexander**, MPhil, MRCOG

Consultant Obstetrician and Gynaecologist, Simpson Centre for Reproductive Health, Royal Infirmary of Edinburgh, 51 Little France Crescent, Old Dalkeith Road, Edinburgh EH16 4SA, UK.  
E-mail: [Claire.Alexander@luht.scot.nhs.uk](mailto:Claire.Alexander@luht.scot.nhs.uk)

**Introduction**

Obesity is a big problem getting bigger. The prevalence of obesity now exceeds the critical threshold of 15%, as defined by the World Health Organization (WHO), to be described as an epidemic.<sup>1</sup> The WHO defines normal weight as a body mass index (BMI) <25.0 kg/m<sup>2</sup>, overweight as 25.0–29.9 kg/m<sup>2</sup> and obese as a BMI >30.0 kg/m<sup>2</sup>. In the UK House of Commons Health report on obesity in 2004, it was predicted that obesity would soon overtake smoking as the leading health problem in the UK.<sup>2</sup>

A pregnancy associated with obesity is at increased risk of most major pregnancy complications. In a large cohort study in the North Thames Region, Sebire *et al.* found that gestational diabetes, pre-eclampsia, induction of labour, emergency Caesarean section, postpartum haemorrhage, genital tract infection, wound infection, birth weight above the 90th centile and intrauterine fetal death are all significantly more likely to occur in the obese parturient than her normal-weight counterpart.<sup>3</sup> Furthermore, it is well known that the incidence of obesity is increasing in pregnancy, with investigators in Scotland and in the USA recently recording increasing BMI in women in early pregnancy over a 10-year period and an up to two-fold increase in the number of obese pregnant women in the same time period.<sup>4,5</sup>

The '2005 American Committee on Obstetrics and Gynecology Opinion on Obesity in Pregnancy' stated that one-third of pregnant women in the USA are obese and recognised that these women are at increased risk of the complications previously mentioned.<sup>6</sup> It emphasised the need for obstetricians to provide pre-pregnancy counselling for such women and encourage weight reduction programmes prior to pregnancy.

Clearly, contraception in obese women is an important area for health professionals and, indeed, health care providers and politicians. Good contraception can give obese women the opportunity to optimise their health prior to pregnancy. It can allow time for the health professional to encourage weight loss and stabilise any other co-morbidities. It is also crucial from a health economic point of view. As a result, there have been an increasing number of studies addressing contraception in obese women. The following three recent publications have been chosen to illustrate aspects of this health issue.

### Contraceptive use by diabetic and obese women.

Chuang CH, Chase GA, Bensyl DM, Weisman CS. *Womens Health Issues* 2005; **15**: 167–173

Chuang *et al.* from Pennsylvania performed a cross-sectional, retrospective study using data from 11 states in 2000. They analysed contraceptive use in 7943 sexually active non-sterilised women and compared that with BMI, age, ethnicity, marital status, education and socioeconomic status. They found a significant increase in contraceptive non-use in overweight and obese women when compared with normal-weight women. When they

controlled for the co-variables mentioned, obesity remained a significant predictor of contraceptive non-use (OR 1.34, 95% CI 1.16–1.55). The authors postulate that obese women perceive themselves as less fertile and as having a negative perception of contraception in the context of being overweight. This under-use may also illustrate the difficulties that some obese women have in accessing health care. Whilst possible weaknesses in the study may have occurred because the BMI was calculated from self-reported height and weight, and bias could have been introduced as data were only gathered from 11 states, it is an interesting paper and highlights the need for advice and action in relation to preventative health care and contraception in this group.

### The association between body weight, unintended pregnancy resulting in a livebirth, and contraception at the time of conception.

Brunner Huber LR, Hogue CJ. *Matern Child Health J* 2005; **9**: 413–420

In this research article, the same multi-state database was accessed, but from 1999, and information on pregnancy intention, BMI and contraceptive use at the time of conception was analysed. Unintended pregnancy was defined as an 'unwanted' or 'mis-timed' pregnancy. The BMI data were again self-reported and the method of contraception unfortunately not determined. The authors recognised that not all unintended pregnancies represent contraceptive failures and not all contraceptive failures are unintended. They state that of the 6 million pregnancies in the USA each year, 3 million are unintended. Half of them, however, occur in the 90% of women who use some form of contraception. The other half occur in women who are not using contraception despite an intention not to become pregnant. The women were analysed in two groups: those using and those not using contraception, and within those groups the authors determined which women had unintended pregnancies. Following multivariable logistic regression analysis, the authors found an association between BMI and unintended pregnancy in the group using contraception in overweight and obese women when compared to normal-weight women. Obese women who were non-smokers were more likely to have unintended pregnancies than lighter women who did not smoke. The authors hypothesise that as non-smokers were more likely to be using the combined oral contraceptive pill (COC) than smokers, the obese non-smokers were at greater risk of unintended pregnancy as the COC was more likely to fail due to problems with absorption and increased levels of free oestrogen affecting negative feedback mechanisms. The method of contraception was, however, not determined. Unfortunately the database only included women with live births and so no data were available about BMI, contraception and pregnancy intention in women who underwent induced abortion.

### Weight-related issues and high-risk sexual behaviours among college students.

Eisenberg ME, Neumark-Sztainer D, Lust KD. *J Am Coll Health* 2005; **54**: 95–101

This group from Minneapolis analysed weight-related issues and 'high-risk' sexual behaviours in a group of college students completing a questionnaire. The questionnaire assessed sexual risk-taking behaviour, BMI, body image and unhealthy weight-modifying behaviours such as inducing vomiting, binge eating, use of laxatives, and so on. The response rate was 39%. Of the respondents, 20% were overweight and 7% obese. Some 42% of female respondents were never or rarely satisfied with their body image and one-third exhibited unhealthy weight control behaviours. There was a positive association, in female students, with high BMI and casual sex and intoxication at the time of most recent intercourse. Unhealthy weight-modifying behaviour was significantly associated with casual sex, non-use of condoms and with intoxication. Interestingly, the differences were not demonstrated in the males studied. The authors conclude that whilst their findings might simply represent clustering of risk-taking behaviours previously described in adolescent health literature, it may reflect a situation where young women with increased BMI are engaging in high-risk sexual behaviours in order to feel better about themselves by demonstrating the ability to attract a partner.

**Conclusions**

Few anti-obesity interventions including drugs, surgery, diet and behavioural therapies have been shown to be effective in the short term for the treatment of obesity.<sup>7</sup> Therefore, obstetricians and gynaecologists need to develop strategies in order to care for women with obesity and related problems in order to maximise health and minimise complications.

Unfortunately, the above studies suggest that obese women may be more likely to demonstrate 'high-risk' sexual behaviours, are at greater risk of contraceptive failure and are more likely to report contraceptive non-use. Whilst there is a need for education and health promotion to tackle the rise in obesity, there is also clearly need for targeted education about contraception in addition to improved access to contraception for overweight and obese women. Further study of contraceptive use and outcomes of obesity intervention in this group would be of value.

**References**

- 1 International Obesity Task Force. Task Force About Obesity. <http://www.iof.org> [Accessed 12 May 2006].
- 2 House of Commons Health Committee. *Obesity*. London, UK: Stationery Office, 2004. <http://www.publications.parliament.uk/pa/cm200304/cmselect/cmhealth/23/23.pdf> [Accessed 12 May 2006].
- 3 Sebire NJ, Jolly M, Harris JP, Wadsworth J, Joffe M, Beard RW, *et al.* Maternal obesity and pregnancy outcome: a study of 287 213 pregnancies in London. *Int J Obesity* 2001; **25**: 1175–1182.
- 4 Kanagalingam MG, Forouhi NG, Greer IA, Sattar N. Changes in body mass index over a decade: retrospective analysis from a Glasgow Maternity Hospital. *Br J Obstet Gynaecol* 2005; **112**: 1431–1433.
- 5 Yeh J, Shelton MS. Increasing pre-pregnancy body mass index: analysis of trends and contributing variables. *Am J Obstet Gynecol* 2005; **193**: 1994–1998.
- 6 American College of Obstetricians and Gynecologists. ACOG Committee Opinion number 315, September 2005. Obesity in pregnancy. *Obstet Gynecol* 2005; **106**: 671–675.
- 7 McTigue KM, Harris R, Hemphill B, Lux L, Sutton S, Buntun AJ, Lohr KN. Screening and interventions for obesity in adults: summary of the evidence of the US Preventive Services Task Force. *Ann Intern Med* 2003; **139**: 939–949.