

References

- 1 United Nations (UN). *United Nations Millennium Declaration: Resolution Adopted by the General Assembly, 55/2*. New York, NY: UN, 2000.
- 2 World Health Organization (WHO). *The Prevention and Management of Unsafe Abortion. Report of a Technical Working Group*. Geneva, Switzerland: WHO, 1992.
- 3 World Health Organization (WHO). *Safe Abortion: Technical and Policy Guidance for Health Care Systems*. Geneva, Switzerland: WHO, 2003.
- 4 World Health Organization (WHO). *Unsafe Abortion: Global and Regional Estimates of the Incidence of Unsafe Abortion and Associated Mortality in 2000*. Geneva, Switzerland: WHO, 2004.
- 5 Alan Guttmacher Institute. *Sharing Responsibility: Women, Society and Abortion Worldwide*. New York, NY: The Alan Guttmacher Institute, 1999.
- 6 United Nations (UN). *Programme of Action Adopted at the International Conference on Population and Development, Cairo 5-13 September 1994*. New York, NY: UN, 1994.
- 7 Salter CL, Johnston HB, Hengen N. Care for postabortion complications: saving women's lives. *Popul Rep L* 1997; **10**: 1.
- 8 Billings DL, Benson J. Postabortion care in Latin America: policy and service recommendations from a decade of operations research. *Health Policy Plan* 2005; **20**: 158-166.
- 9 Koontz SL, Molina de Perez O, Leon K, Foster-Rosales A. Treating incomplete abortion in El Salvador: cost savings with manual vacuum aspiration. *Contraception* 2003; **68**: 345-351.
- 10 Dickson-Tetteh K, Billings DL. Abortion care services provided by registered midwives in South Africa. *Int Fam Plan Perspect* 2002; **28**: 144-150.
- 11 Ipas, Division of International Health (IHCAR). *Deciding Women's Lives are Worth Saving: Expanding the Role of Midlevel Providers in Safe Abortion Care*. Chapel Hill, NC: Ipas, 2002.
- 12 Dayaratna V, Winfrey W, McGreevy W, Hardee K, Smith J, Mumford E, et al. *Reproductive Health Interventions: Which Ones Work and What Do They Cost?* Washington, DC: The Futures Group International, 2000.
- 13 Jowett M. Safe motherhood interventions in low-income countries: an economic justification and evidence of cost effectiveness. *Health Policy* 2000; **53**: 201-228.
- 14 Huntington D, Nawar L, Hassan EO, Youssef H, Abdel-Tawab N. The postabortion caseload in Egyptian hospitals: a descriptive study. *Int Fam Plan Perspect* 1998; **24**: 25-31.
- 15 de Pinho H, McIntyre D. *Cost Analysis of Abortions Performed in the Public Sector*. Cape Town, South Africa: University of Cape Town Department of Health, 1997.
- 16 King T, Benson J, Stein K. *Comparing the Cost of Postabortion Care in Africa and Latin America: The DataPAC Project*. New York, NY: Population Council, 1997.
- 17 Blumenthal PD, Remsburg RE. A time and cost analysis of the management of incomplete abortion with manual vacuum aspiration. *Int J Gynaecol Obstet* 1994; **45**: 261-267.
- 18 Bradley J, Rogo K, Johnson R, Okoko L, Healy J, Benson J. A comparison of the costs of manual vacuum aspiration (MVA) and evacuation and curettage (E and C) in the treatment of early incomplete abortions in Kenya. *J Obstet Gynaecol East Cent Africa* 1993; **11**: 12-19.
- 19 Johnson BR, Benson J, Bradley J, Ordoñez AR. Costs and resource utilization for the treatment of incomplete abortion in Kenya and Mexico. *Soc Sci Med* 1993; **36**: 1443-1453.
- 20 Management Sciences for Health (MSH). *CORE: A Tool for Cost and Revenue Analysis, User's Guide: Version 1.0*. Boston, MA: MSH, 1998.
- 21 Management Sciences for Health (MSH), John Snow, Inc., United States Pharmacopeia. *CES: Cost-Estimate Strategy for Reproductive Health Commodity Management, User's Guide*. Arlington, VA: MSH, 2000.
- 22 The Policy II Project. Washington, DC: Policy Project, 2007. <http://www.policyproject.com/software.cfm?page=Software&ID=Spectrum> [Accessed 10 April 2007].
- 23 World Health Organization (WHO). *Mother-Baby Package Costing Spreadsheet*. Geneva, Switzerland: WHO Reproductive Health and Research Unit, 1999.
- 24 Janowitz B, Bratt JH. *Methods for Costing Family Planning Services*. New York, NY: United Nations Populations Fund and Family Health International, 1999.
- 25 PRIME II. *Cost and Results Analysis: Practical Applications in International Health from the PRIME II Project. Better Practices in Performance Improvement*. Chapel Hill, NC: PRIME II, 2003.
- 26 Weissman E, Sentumbwe-Mugisa O, Mbonye AK, Kayaga E, Kihuguru SM, Lissner C. *Uganda Safe Motherhood Programme Costing Study*. Geneva, Switzerland: World Health Organization, 1999.
- 27 Management Sciences for Health (MSH). *The International Drug Price Indicator Guide, 2003*. Boston, MA: MSH, 2004.
- 28 Nawar L, Huntington D, Fattah MNA. Cost analysis of postabortion care in Egypt. In: Huntington D, Piet-Pelon NJ (eds), *Postabortion Care: Lessons Learned from Operations Research*. New York, NY: Population Council, 1999.
- 29 Uganda Bureau of Statistics (UBOS), ORC Macro. *Uganda Demographic and Health Survey 2000-2001*. Calverton, MD: UBOS and ORC Macro, 2001.
- 30 Gebreselassie H, Gallo MF, Monyo A, Johnson BR. The magnitude of abortion complications in Kenya. *Br J Obstet Gynaecol* 2005; **112**: 1229-1235.
- 31 Mumford EA, Dayaratna V, Winfrey W, Since J, McGreevy WP. *Reproductive Health Costs: Literature Review. Policy Project, Working Paper Series No. 3, 1998*. <http://www.policyproject.com/abstract.cfm/240> [Accessed 13 March 2006].
- 32 Johnson BR, Ndhlovu S, Farr SL, Chipato T. Reducing unplanned pregnancy and abortion in Zimbabwe through postabortion contraception. *Stud Fam Plann* 2002; **33**: 195-202.
- 33 Solo J, Billings DL, Aloo-Obunga C, Ominde A, Makumi M. Creating linkages between incomplete abortion treatment and family planning services in Kenya. *Stud Fam Plann* 1999; **30**: 17-27.
- 34 Warriner IK, Meiric O, Hoffman M, Morroni C, Harries J, My Huong NT, et al. Rates of complication in first trimester manual vacuum aspiration abortion done by doctors and mid-level providers in South African and Vietnam: a randomized controlled equivalence trial. *Lancet* 2006; **368**: 1965-1972.
- 35 World Health Organization (WHO). *World Health Report 2004: Changing History*. Geneva, Switzerland: WHO, 2004.
- 36 Cook RJ, Dickens BM, Fathalla MF. *Reproductive Health and Human Rights: Integrating Medicine, Ethics, and Law*. New York, NY: Oxford University Press, 2003.

Celebrating 20 years of Positively Women

The HIV charity, Positively Women, has launched a special edition of its magazine to mark the organisation's 20th anniversary. The commemorative edition features the stories of 24 women diagnosed over the last 20 years. The different stories illustrate the changing attitudes and treatment of women living with HIV over the last two decades.

For further information contact Lucy Osman on 020 7713 0444 or email losman@positivelywomen.org.uk.

Independent Advisory Group on Sexual Health and HIV

Strong links between alcohol, drugs and risky sexual behaviour are providing fuel for a "sexual health crisis" in Britain. Sexual health among the

young has deteriorated seriously in the last 12 years, and media coverage of celebrities' hard-partying lifestyles may make the problem worse, are key findings of the Independent Advisory Group on Sexual Health. "Young people, from their early teens, are defining a lifestyle that involves alcohol, drugs and sex", said the group's Chair, Baroness Joyce Gould.

The report found that Britain had the highest rate of sexually transmitted infections (STIs) and teenage pregnancy in Europe. There has been a "disturbing" increase in STIs - over the last 12 years, chlamydia and HIV are up 300% and syphilis 2000%. The study found a "strong correlation" between STIs, sex and drug use and added that alcohol "can increase the risk of having unprotected sex". The report also highlighted the impact that advertising and celebrities can have from a young age, citing the

"publicity and coverage of sex, drugs and excessive drinking by 'celebrities' and 'pop idols' and by characters in 'soaps', making it an acceptable lifestyle". The report's authors recommended a more collaborative and less "puritanical" official approach to the issues and said condoms should be more accessible to young people and sex education mandatory in schools. Factors including young people having parents who were ambitious for them plus a confidante as well as achievement at school were all helpful.

Source: http://www.dh.gov.uk/en/Policyandguidance/Healthandsocialcaretopics/Sexualhealth/Sexualhealthgeneralinformation/DH_4079794

Reported by **Henrietta Hughes**, MRCPG, DFFP
General Practitioner, London, UK