

**Tabel X 1 Sexuality education ever in life among non-migrant, migrant, and second-generation migrant women seeking abortion care in Stockholm County**

	<b>Non-migrant n (%)</b>	<b>Migrant n (%)</b>	<b>Second- generation migrant n (%)</b>	<b>Total n (%)</b>	<b>p-value</b>
Yes	414 (97.6)	93 (62.8)	60 (93.8)	567 (89.2)	<0.001
No	10 (2.4)	55 (37.2)	4 (6.3)	69 (10.8)	

**Tabel X 2 Contraception counselling/information ever in life among non-migrant, migrant, and second-generation migrant women seeking abortion care in Stockholm County**

	<b>Non- migrant n (%)</b>	<b>Migrant n (%)</b>	<b>Second- generation migrant n (%)</b>	<b>Total n (%)</b>	<b>p-value</b>
Yes	418 (98.8)	128 (86.5)	62 (96.9)	608 (95.7)	<0.001 <sup>b</sup>
No	4 (0.9)	19 (12.8)	2 (3.1)	25 (3.9)	
Don't know	1 (0.2)	1 (0.7)	0 (0)	2 (0.3)	

<sup>b</sup>Fischer Exact test

**Tabel X 3 Reasons for contraceptive failure among non-migrant, migrant, and second-generation migrant women seeking abortion care in Stockholm County**

	<b>Non- migrant n (%)</b>	<b>Migrant n (%)</b>	<b>Second- generation migrant n (%)</b>	<b>Total n (%)</b>	<b>p-value</b>
Miscalculated days	14 (9.8)	7 (20.0)	2 (11.1)	23 (11.7)	0.706 <sup>a</sup>
Condom failure	31 (21.7)	8 (22.9)	5 (27.8)	44 (22.4)	
IUC misplaced/expelled	1 (0.7)	0 (0)	1 (5.6)	2 (1.0)	
No withdrawal	11 (7.7)	3 (8.6)	0 (0)	14 (7.1)	
Not used acc. to prescriptions	36 (25.2)	5 (14.3)	5 (27.8)	46 (23.5)	
Drug interaction	1 (0.7)	1 (2.9)	0 (0)	2 (1.0)	
Vomited/diarrhea	6 (4.2)	2 (5.7)	1 (5.6)	9 (4.6)	
Don't know	27 (18.9)	6 (17.1)	3 (16.7)	36 (18.4)	
Other	16 (11.2)	3 (8.6)	1 (5.6)	20 (10.2)	

<sup>a</sup>Fischer Exact test obtained through a Monte Carlo simulation

**Tabel X 4 Important factors for choosing contraception except become pregnant among non-migrant, migrant, and second-generation migrant women seeking abortion care in Stockholm County**

	<b>Non-migrant</b>	<b>Migrant</b>	<b>Second-generation migrant</b>	<b>Total</b>	<b>p-value</b>
	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>	
Regular bleeding	24 (6.3)	4 (3.1)	2 (3.8)	30 (5.3)	0.023 <sup>a</sup>
No bleeding	26 (6.8)	5 (3.9)	0 (0)	31 (5.5)	
Long-acting/don't need to remember	30 (7.9)	16 (12.4)	5 (9.4)	51 (9.1)	
Start immediately	2 (0.5)	1 (0.8)	1 (1.9)	4 (0.7)	
Partner satisfied/accepts the method	1 (0.3)	2 (1.6)	0 (0)	3 (0.5)	
The cost	0 (0)	0 (0)	0 (0)	0 (0)	
STI protection	1 (0.3)	4 (3.1)	1 (1.9)	6 (1.1)	
The effectiveness	40 (10.5)	23 (17.8)	2 (3.8)	65 (11.5)	
No hormones	33 (8.7)	5 (3.9)	5 (9.4)	43 (7.6)	
Less hormones	35 (9.2)	9 (7.0)	5 (9.4)	49 (8.7)	
Invisible	1 (0.3)	0 (0)	0 (0)	1 (0.2)	
Well-known/friends recommended	6 (1.6)	1 (0.8)	1 (1.9)	8 (1.4)	
No side effects	122 (32.0)	35 (27.1)	20 (37.7)	177 (31.4)	
Other	60 (15.7)	24 (18.6)	11 (20.8)	95 (16.9)	

<sup>a</sup>Fischer Exact test obtained through a Monte Carlo simulation