

Supplementary Table 1 Sensitivity analyses											
		0		2-4		5-9		10-19		≥20	
		OR [95% CI] ¹	<i>p</i>	OR [95% CI]	<i>p</i>	OR [95% CI]	<i>p</i>	OR [95% CI]	<i>p</i>	OR [95% CI]	<i>p</i>
Men											
	Fair/poor self-rated health	0.52 [0.25-1.07]	.077	0.86 [0.25-1.07]	.289	0.94 [0.67-1.31]	.714	1.36 [0.92-2.02]	.123	1.01 [0.67-1.53]	.970
	Limiting long-standing illness	0.76 [0.39-1.48]	.414	0.82 [0.64-1.07]	.142	1.15 [0.85-1.55]	.369	1.03 [0.71-1.49]	.883	1.09 [0.75-1.59]	.664
	Cancer	1.39 [0.46-4.18]	.562	1.62 [1.04-2.52]	.033	1.54 [0.91-2.61]	.110	1.83 [0.99-3.39]	.054	1.64 [0.85-3.17]	.139
	Coronary heart disease	0.68 [0.31-1.52]	.347	0.99 [0.71-1.38]	.954	0.87 [0.57-1.33]	.524	0.58 [0.33-1.03]	.063	1.15 [0.69-1.91]	.587
	Stroke	1.00 [0.30-3.33]	.995	0.62 [0.36-1.07]	.088	0.85 [0.45-1.63]	.634	0.92 [0.41-2.03]	.830	1.13 [0.51-2.53]	.760
Women											
	Fair/poor self-rated health	1.76 [1.01-3.06]	.048	1.17 [0.92-1.49]	.201	1.32 [0.95-1.83]	.101	1.19 [0.71-1.99]	.503	0.78 [0.41-1.46]	.435
	Limiting long-standing illness	1.33 [0.77-2.29]	.304	1.19 [0.96-1.48]	.104	1.57 [1.18-2.09]	.002	1.89 [1.23-2.92]	.004	1.41 [0.83-2.39]	.204
	Cancer	1.81 [0.75-4.37]	.188	1.42 [0.95-2.12]	.088	1.28 [0.74-2.21]	.377	2.06 [0.99-4.27]	.052	2.03 [0.84-4.89]	.115
	Coronary heart disease	1.31 [0.63-2.74]	.474	1.31 [0.91-1.89]	.142	1.09 [0.60-1.98]	.772	0.33 [0.08-1.43]	.137	0.50 [0.11-2.30]	.372
	Stroke	2.52 [0.97-6.54]	.058	1.29 [0.78-2.14]	.317	1.65 [0.81-3.37]	.165	2.03 [0.71-5.78]	.185	0.36 [0.03-4.64]	.431
¹ Adjusted odds ratios (OR) and 95% confidence intervals (CI) for reporting health problem relative to the group who had had 1 sexual partner. Notes: All percentages and odds ratios are adjusted for age, partnership status, ethnicity, wealth, smoking status, alcohol intake, physical activity and depressive symptoms, and weighted for sampling probabilities and differential nonresponse.											

