

Table S2. Non-randomised, observational comparison of experiences of women who received an IUD by 6 weeks, either immediate or delayed insertion after medical abortion at 17-20 weeks gestation.

	Immediate insertion n=45	Delayed insertion n=25	p-value ^a
IUD-related complications			
Signs of localized infection			
<i>At 6 weeks (n)^b</i>	n=37	n=11	
Yes	4 (11)	2 (18)	0.42
Cramping over the last 6 weeks			
<i>At 3 months (n)</i>	n=34	n=21	0.56
None / Mild	31 (91)	19 (90)	
Moderate	3 (9)	1 (5)	
High	0 (0)	1 (5)	
Satisfaction	n (%)	n (%)	
<i>At 6 weeks (n)^b</i>	n=37	n=9	0.64
Very satisfied	35 (95)	9 (100)	
Somewhat satisfied	2 (5)	0 (0.0)	
Somewhat / very dissatisfied	0 (0)	0 (0.0)	
<i>At 3 months (n)</i>	n=34	n=22	0.15
Very satisfied	30 (88)	22 (100)	
Somewhat satisfied	4 (12)	0 (0.0)	
Somewhat / very dissatisfied	0 (0.0)	0 (0.0)	
<i>At 6 months (n)</i>	n=26	n=16	0.28
Very satisfied	23 (89)	16 (100)	
Somewhat satisfied	3 (12)	0 (0)	
Somewhat / very dissatisfied	0 (0)	0 (0)	
Would recommend to a friend having TOP			
<i>At 3 months (n)</i>	n=36	n=22	~1.00
Highly agree	35 (97)	22 (100)	
Somewhat agree	0 (0)	0 (0)	
Somewhat / highly disagree	1 (3)	0 (0)	

^a Chi-squared or Fisher's exact test for proportions

^b Excludes participants getting the IUD for the first time at 6 weeks but includes participants that had IUD replaced at 6 weeks