



## Supplementary Material

### Supplementary Figure S1: Participant flow diagram showing selection and inclusion in study



Adapted from Dema, et al., 2022: **Methodology of Natsal-COVID Wave 2: A large, quasi-representative, longitudinal survey measuring the impact of COVID-19 on sexual and reproductive health in Britain** [version 1; peer review: awaiting peer review] (<https://wellcomeopenresearch.org/articles/7-166/v1>), Figure 1

## Supplementary Figure S2: Accessing contraception services



## Supplementary Box 1 – Classifications of contraceptive types

Participants' responses to questions of contraceptives used were classified into 'more effective' and 'less effective' types by typical use failure rates below and above 10% respectively. Emergency methods were categorised separately.

### More effective methods:

Intrauterine device (IUD)	Oral contraceptive pill
Intrauterine system (IUS)	Injection
Implant	Transdermal patch
Vaginal ring	

### Less effective methods:

Condoms	Withdrawal
Spermicides	'Other' methods
Rhythm method	

### Emergency methods:

Emergency contraceptive pills	Emergency IUD insertion
-------------------------------	-------------------------

## Supplementary Table S1: Socio-demographic characteristics of sexually active participants aged 18-44 years, described as female at birth who reported sex with a man in the past year

Age group	18-24	25-29	30-34	35-44	All ages
<b>Total</b>					
Distribution across age categories (% (95% CI))	16.7 (15.3, 18.1)	25.3 (23.7, 27.1)	21.3 (19.7, 22.9)	36.7 (34.9, 38.6)	100.0%
Denominators (weighted, unweighted)	213, 265	324, 397	272, 320	470, 506	1279, 1488
<b>Distributions within age categories</b>					
	<b>% (95% CI)</b>	<b>% (95% CI)</b>	<b>% (95% CI)</b>	<b>% (95% CI)</b>	<b>% (95% CI)</b>
<b>Ethnicity</b>					
White	79.0 (72.9, 84.0)	86.0 (81.8, 89.4)	86.4 (81.8, 90.0)	90.8 (87.8, 93.1)	86.7% (84.7, 88.5)
Black or Black African or Black Caribbean or Black British	9.4 (6.1, 14.3)	5.4 (3.4, 8.5)	1.6 (0.6, 4.1)	0.8 (0.3, 2.2)	3.6% (2.7, 4.7)
Asian or Asian British	8.0 (5.0, 12.5)	4.2 (2.5, 7.1)	7.7 (5.1, 11.6)	6.0 (4.2, 8.6)	6.2% (5.0, 7.7)
Mixed or multiple or other ethnic groups	3.6 (1.8, 7.2)	4.4 (2.6, 7.2)	4.2 (2.4, 7.4)	2.4 (1.3, 4.2)	3.5% (2.6, 4.6)
Missing: n=10 (0.7%)					
<b>Self-described sexual identity</b>					
Heterosexual or Straight	92.4 (88.0, 95.3)	96.1 (93.3, 97.7)	97.7 (95.0, 98.9)	98.5 (96.9, 99.3)	96.7% (95.5, 97.5)
Lesbian, Gay, Bisexual or Other	7.6 (4.7, 12.0)	3.9 (2.3, 6.7)	2.3 (1.1, 5.0)	1.5 (0.7, 3.1)	3.3% (2.5, 4.5)
Missing: n=12 (0.8%)					
<b>Social Grade</b>					

Age group	18-24	25-29	30-34	35-44	All ages
A/B - Higher/intermediate managerial, administrative and professional	17.6 (13.0, 23.3)	29.8 (25.1, 35.0)	23.7 (19.0, 29.2)	19.6 (16.2, 23.4)	22.7% (20.5, 25.1)
C1/C2 - Supervisory, clerical and junior managerial, administrative and professional, skilled manual workers	39.3 (32.9, 46.0)	49.1 (43.7, 54.6)	50.3 (44.3, 56.2)	61.4 (56.9, 65.7)	52.3% (49.5, 55.0)
D/E - Semi-skilled and unskilled manual, casual, lowest grade and unemployed	43.1 (36.6, 49.9)	21.1 (16.9, 25.9)	26.0 (21.1, 31.6)	19.0 (15.7, 22.8)	25.0% (22.7, 27.4)
Missing: n=0 (0%)					
<b>Education level</b>					
Degree	28.3 (22.6, 34.7)	58.5 (53.1, 63.8)	59.0 (53.1, 64.7)	54.8 (50.2, 59.2)	52.2% (49.5, 55.0)
Below degree	65.1 (58.5, 71.3)	36.7 (31.6, 42.1)	37.2 (31.6, 43.1)	43.4 (39.0, 47.9)	44.0% (41.3, 46.8)
No qualifications	6.6 (3.9, 10.9)	4.7 (2.9, 7.7)	3.8 (2.1, 6.8)	1.8 (0.9, 3.5)	3.8% (2.8, 4.9)
Missing: n=0 (0%)					
<b>Living together – relationship but not living together – Single</b>					
Married/steady and living together	34.4 (28.3, 41.0)	71.0 (65.8, 75.7)	78.5 (73.2, 83.0)	82.2 (78.4, 85.4)	70.6% (68.0, 73.0)
Married/steady NOT living together	29.2 (23.5, 35.7)	11.7 (8.6, 15.7)	7.7 (5.1, 11.6)	7.3 (5.3, 10.0)	12.1% (10.4, 14.0)
Casual, new, >1, at end or other*	18.5 (13.8, 24.3)	7.7 (5.3, 11.2)	4.2 (2.4, 7.4)	3.8 (2.4, 6.0)	7.3% (6.0, 8.9)
Single	17.9 (13.3, 23.7)	9.5 (6.8, 13.3)	9.6 (6.6, 13.7)	6.7 (4.8, 9.4)	9.9% (8.4, 11.7)
Missing: n=1 (0.1%)					

Age group	18-24	25-29	30-34	35-44	All ages
<b>Been furloughed under the Coronavirus Job Retention Scheme</b>					
No	82.8 (77.0, 87.3)	85.5 (81.2, 88.9)	81.0 (75.9, 85.3)	83.1 (79.4, 86.2)	83.2% (81.0, 85.2)
Yes	17.2 (12.7, 23.0)	14.5 (11.1, 18.8)	19.0 (14.7, 24.1)	16.9 (13.8, 20.6)	16.8% (14.8, 19.0)
Missing: n=12 (0.8%)					
<b>Became unemployed</b>					
No	86.2 (80.8, 90.3)	92.4 (89.0, 94.9)	91.9 (88.0, 94.6)	91.9 (89.0, 94.1)	91.1% (89.4, 92.5)
Yes	13.8 (9.7, 19.2)	7.6 (5.1, 11.0)	8.1 (5.4, 12.0)	8.1 (5.9, 11.0)	8.9% (7.5, 10.6)
Missing: n=12 (0.8%)					
<b>Number of days drinking in past week</b>					
0 days	38.0 (31.7, 44.7)	40.1 (34.9, 45.5)	46.0 (40.2, 52.0)	45.7 (41.2, 50.2)	43.0% (40.3, 45.8)
1-2 days	41.0 (34.6, 47.8)	40.9 (35.6, 46.4)	39.5 (33.9, 45.5)	35.3 (31.1, 39.7)	38.6% (35.9, 41.3)
3-4 days	17.3 (12.8, 23.0)	13.1 (9.8, 17.3)	9.5 (6.5, 13.7)	11.3 (8.7, 14.5)	12.4% (10.7, 14.3)
5-7 days	3.7 (1.8, 7.2)	5.9 (3.8, 9.1)	4.9 (2.9, 8.2)	7.8 (5.7, 10.6)	6.0% (4.8, 7.5)
Missing: n=2 (0.1%)					

Age group	18-24	25-29	30-34	35-44	All ages
<b>Drinking habits compared to pre Covid-19 outbreak</b>					
Less these days	36.9 (30.6, 43.6)	30.9 (26.1, 36.2)	28.8 (23.7, 34.6)	25.8 (22.0, 30.0)	29.6% (27.1, 32.2)
About the same	40.7 (34.2, 47.4)	51.6 (46.1, 57.1)	57.2 (51.1, 63.0)	56.9 (52.4, 61.4)	52.9% (50.1, 55.6)
More these days	22.4 (17.3, 28.6)	17.5 (13.7, 22.0)	14.0 (10.3, 18.7)	17.3 (14.1, 21.0)	17.5% (15.5, 19.7)
Missing: n=19 (1.3%)					
<b>Current smoker</b>					
No	75.9 (69.7, 81.2)	73.4 (68.3, 77.9)	79.8 (74.6, 84.2)	82.3 (78.5, 85.5)	78.5% (76.1, 80.6)
Yes	24.1 (18.8, 30.3)	26.6 (22.1, 31.7)	20.2 (15.8, 25.4)	17.7 (14.5, 21.5)	21.5% (19.4, 23.9)
Missing: n=2 (0.1%)					
<b>Symptoms of depression (PHQ2 score)</b>					
No symptoms of depression (0-2)	47.8 (41.0, 54.6)	63.1 (57.6, 68.2)	67.2 (61.3, 72.5)	74.6 (70.4, 78.3)	65.7% (63.1, 68.3)
Symptoms of depression (3-6)	52.2 (45.4, 59.0)	36.9 (31.8, 42.4)	32.8 (27.5, 38.7)	25.4 (21.7, 29.6)	34.3% (31.7, 36.9)
Missing: n=15 (1.0%)					
<b>Symptoms of anxiety (GAD2 score)</b>					
No symptoms of anxiety (0-2)	46.4 (39.7, 53.3)	58.3 (52.8, 63.5)	63.7 (57.8, 69.2)	69.4 (65.0, 73.4)	61.6% (58.9, 64.2)

Age group	18-24	25-29	30-34	35-44	All ages
Symptoms of anxiety (3-6)	53.6 (46.7, 60.3)	41.7 (36.5, 47.2)	36.3 (30.8, 42.2)	30.6 (26.6, 35.0)	38.4% (35.8, 41.1)

Missing: n=8 (0.5%)

\* In a 'casual' relationship, in a 'new' relationship, in more than one relationship, recently ended a relationship or 'other' relationship status

## Supplementary Table S2: Definitions of outcome variables

Outcome/Variable	Denominator	Definition
Usual contraception used in the year before lockdown	At risk of unplanned pregnancy (n=1,169)	Selected contraceptive method in response to "In the year before the start of the first lockdown (23 March 2020), which of the following did you or a partner use to prevent pregnancy?". Participants selecting more than one method were asked to specify their 'usual' method.
Usual contraception used in the year since lockdown started	At risk of unplanned pregnancy (n=1,169)	Selected contraceptive method in response to "Since the start of the first lockdown, which of the following did you or a partner use to prevent pregnancy?". Participants selecting more than one method were asked to specify their 'usual' method.
Switched contraceptives due to pandemic	At risk of unplanned pregnancy (n=1,169)	Reported "I [temporarily/permanently] changed to a different method to prevent pregnancy" in response to "Since the start of the first lockdown, have any of these things happened because of the pandemic?"



Accessed services successfully	Tried to access a contraceptive service at least once (n=441)	Reported "Services Accessed during lockdown: Contraception services/advice" and did <b>not</b> report "Services tried but failed to access: Contraception services/advice"
Faced difficulty accessing services but was able to access in the end	Tried to access a contraceptive service at least once (n=441)	Reported "Services tried but failed to access: Contraception services/advice" and either "I accessed it eventually" or "I used a different service" to question "what happened in the end?"
Unable to access contraceptive services	Tried to access a contraceptive service at least once (n=441)	Reported "Services tried but failed to access: Contraception services/advice" and did <b>not</b> report "I accessed it eventually" or "I used a different service" to question "what happened in the end?"
Successful use of contraceptive services (odds ratio)	All participants (n=1,488)	Reported "Services Accessed during lockdown: Contraception services/advice"
Barriers accessing contraceptive services (odds ratio)	All participants (n=1,488)	Reported "Services tried but failed to access: Contraception services/advice"
Pregnancy in past year	All participants (n=1,488)	Reported "Yes" to "Are you currently pregnant?" or "In the last year" to "When was most recent pregnancy, even if didn't carry the baby to term?"
London Measure of Unplanned Pregnancy (LMUP) score	Participants with a pregnancy in last 5 years	Questions and scoring detailed at <a href="https://measure.ascody.co.uk/">https://measure.ascody.co.uk/</a>
Unplanned pregnancy in past year	All participants with a pregnancy in past year (n=199)	Reported pregnant in past year and LMUP score <4

## Supplementary Table S3: Service access outcomes – results amongst participants who tried to access contraceptive services (n=441)

### Outcome of attempts to access contraceptive services (% (95% CI))

	Accessed services successfully	Faced difficulty accessing services but was able to access in the end	Unable to access contraceptive services	Denominators (weighted, unweighted)
<b>Total</b>	83.6 (79.5, 87.1)	10.7 (7.9, 14.3)	5.7 (3.7, 8.6)	364, 441
<b>Age</b>				
18-24	80.8 (70.6, 88.0)	14.9 (8.7, 24.6)	4.3 (1.5, 11.7)	82, 104
25-29	83.7 (75.2, 89.7)	11.7 (6.7, 19.5)	4.6 (1.9, 10.9)	104, 136
30-34	81.8 (71.1, 89.1)	9.2 (4.4, 18.5)	9.0 (4.2, 18.2)	74, 89
35-44	87.1 (79.1, 92.3)	7.3 (3.6, 14.2)	5.6 (2.5, 12.2)	104, 112
P-value				p=0.51
<b>Ethnicity</b>				
White	83.7 (79.1, 87.4)	10.3 (7.4, 14.3)	6.0 (3.8, 9.3)	306, 380
Black or Black African or Black Caribbean or Black British	93.4 (66.4, 99.0)	6.6 (1.0, 33.6)		18, 15
Asian or Asian British	73.4 (48.8, 88.9)	21.1 (7.8, 45.8)	5.5 (0.7, 31.4)	20, 20
Mixed or multiple or other ethnic groups	84.5 (57.4, 95.7)	8.4 (1.4, 36.2)	7.1 (1.1, 35.4)	17, 24
P-value				p=0.64
<b>Self-described sexual identity</b>				
Heterosexual or Straight	84.5 (80.3, 88.0)	9.8 (7.1, 13.4)	5.7 (3.7, 8.7)	346, 383

**Outcome of attempts to access contraceptive services (% (95% CI))**

	<b>Accessed services successfully</b>	<b>Faced difficulty accessing services but was able to access in the end</b>	<b>Unable to access contraceptive services</b>	<b>Denominators (weighted, unweighted)</b>
Lesbian, Gay, Bisexual or Other	70.3 (42.8, 88.3)	22.5 (7.6, 50.7)	7.1 (1.0, 37.4)	16, 56
P-value				p=0.047

**Social Grade**

A Upper middle class/ B Middle class	83.8 (74.5, 90.2)	13.3 (7.6, 22.2)	2.9 (0.8, 9.5)	89, 117
C1 Lower middle class/C2 Skilled working class	85.1 (79.1, 89.5)	8.7 (5.4, 13.7)	6.3 (3.5, 10.8)	187, 211
D Working class/ E Lower level of subsistence	80.4 (70.7, 87.5)	12.2 (6.8, 21.0)	7.3 (3.4, 15.1)	88, 113
P-value				p=0.44

**Education level**

Degree	85.5 (79.7, 89.8)	10.7 (7.1, 16.0)	3.8 (1.8, 7.6)	193, 234
Below degree	83.1 (76.4, 88.2)	9.3 (5.7, 15.0)	7.6 (4.4, 13.0)	159, 192
No qualifications	60.8 (29.4, 85.2)	27.6 (8.3, 61.5)	11.6 (1.7, 49.3)	12, 15
P-value				p=0.13

**Living together - relationship but not living together - Single**

Single	87.8 (75.2, 94.4)	6.5 (2.1, 17.9)	5.8 (1.8, 17.1)	50, 61
Casual, new, >1, at end or other	89.1 (70.7, 96.5)	6.9 (1.6, 25.0)	4.0 (0.6, 22.5)	29, 36
Married/steady NOT living together	82.9 (70.0, 90.9)	11.5 (5.2, 23.5)	5.7 (1.8, 16.5)	53, 67
Married/steady and living together	82.2 (76.7, 86.7)	11.9 (8.3, 16.7)	5.9 (3.5, 9.8)	231, 277
P-value				p=0.88

**Outcome of attempts to access contraceptive services (% (95% CI))**

	<b>Accessed services successfully</b>	<b>Faced difficulty accessing services but was able to access in the end</b>	<b>Unable to access contraceptive services</b>	<b>Denominators (weighted, unweighted)</b>
<b>Been furloughed under the Coronavirus Job Retention Scheme</b>				
No	83.6 (79.1, 87.4)	10.5 (7.5, 14.5)	5.9 (3.7, 9.1)	308, 373
Yes	85.2 (72.7, 92.5)	9.9 (4.2, 21.5)	5.0 (1.5, 15.6)	54, 65
P-value				p=0.95
<b>Became unemployed</b>				
No	83.5 (79.0, 87.2)	10.5 (7.6, 14.4)	6.0 (3.9, 9.2)	322, 390
Yes	87.1 (72.2, 94.6)	9.4 (3.4, 23.6)	3.6 (0.7, 17.1)	40, 48
P-value				p=0.79
<b>Number of days drinking</b>				
0 days	82.2 (74.8, 87.7)	11.2 (6.9, 17.7)	6.6 (3.5, 12.2)	140, 166
1-2 days	84.0 (76.8, 89.3)	9.0 (5.2, 15.1)	7.0 (3.8, 12.8)	136, 165
3-4 days	84.2 (72.7, 91.4)	13.0 (6.6, 24.1)	2.8 (0.6, 11.7)	62, 76
5-7 days	88.4 (68.0, 96.5)	10.7 (3.1, 31.1)	0.9 (0.0, 41.4)	25, 34
P-value				p=0.61
<b>Drinking habits compared to pre Covid-19 outbreak</b>				
Less these days	88.8 (81.3, 93.5)	6.4 (3.0, 12.9)	4.8 (2.1, 11.0)	109, 136
About the same	82.3 (75.7, 87.5)	12.5 (8.2, 18.5)	5.2 (2.7, 9.9)	163, 198
More these days	80.5 (70.9, 87.5)	11.7 (6.5, 20.3)	7.7 (3.7, 15.5)	90, 105

**Outcome of attempts to access contraceptive services (% (95% CI))**

	<b>Accessed services successfully</b>	<b>Faced difficulty accessing services but was able to access in the end</b>	<b>Unable to access contraceptive services</b>	<b>Denominators (weighted, unweighted)</b>
P-value				p=0.39
<b>Current smoker</b>				
No	87.5 (83.0, 91.0)	8.3 (5.5, 12.3)	4.2 (2.3, 7.4)	269, 327
Yes	72.4 (62.5, 80.6)	17.4 (11.0, 26.6)	10.1 (5.4, 18.2)	94, 113
P-value				p=0.0026
<b>Symptoms of depression (PHQ2 score)</b>				
No symptoms of depression (0-2)	85.6 (80.2, 89.7)	9.3 (6.1, 14.1)	5.1 (2.8, 9.0)	212, 259
Symptoms of depression (3-6)	80.5 (73.2, 86.2)	12.6 (8.1, 19.2)	6.9 (3.7, 12.4)	145, 174
P-value				p=0.43
<b>Symptoms of anxiety (GAD2 score)</b>				
No symptoms of anxiety (0-2)	87.1 (81.7, 91.1)	8.2 (5.1, 12.9)	4.6 (2.4, 8.6)	202, 235
Symptoms of anxiety (3-6)	78.9 (71.8, 84.6)	13.9 (9.3, 20.3)	7.2 (4.0, 12.4)	159, 203
P-value				p=0.11

## Supplementary Table S4: Stopping or switching contraception because of the pandemic – all participants at risk of unplanned pregnancy who used contraception before Covid

### Stopped or switched contraceptive methods (% (95% CI))

	Stopped using contraceptives	Switched more > less effective	Switched - less > less effective	Switched - more > more effective	Switched less > more effective	Did not switch or stop usual method	Denominators (weighted, unweighted)
<b>Total</b>	3.6 (2.5, 5.1)	2.1 (1.3, 3.3)	4.1 (3.0, 5.8)	6.6 (5.0, 8.5)	0.7 (0.3, 1.6)	82.9 (80.2, 85.4)	811, 957
<b>Age</b>							
18-24	2.6 (0.9, 7.0)	4.1 (1.8, 9.0)	10.7 (6.5, 16.9)	8.5 (4.9, 14.4)	1.4 (0.4, 5.5)	72.7 (64.7, 79.4)	143, 182
25-29	5.1 (2.8, 9.1)	2.7 (1.2, 6.1)	2.1 (0.8, 5.3)	6.4 (3.8, 10.6)	0.4 (0.1, 3.4)	83.2 (77.5, 87.7)	209, 261
30-34	3.5 (1.6, 7.5)	1.8 (0.6, 5.4)	3.4 (1.5, 7.3)	7.9 (4.7, 12.9)	1.0 (0.2, 4.3)	82.4 (76.1, 87.4)	176, 210
35-44	3.0 (1.5, 5.8)	0.7 (0.2, 2.8)	2.8 (1.4, 5.6)	4.9 (2.9, 8.1)	0.5 (0.1, 2.6)	88.2 (83.8, 91.5)	283, 304
P-value							p=0.0020
<b>Ethnicity</b>							
White	3.5 (2.4, 5.2)	1.9 (1.1, 3.3)	3.1 (2.0, 4.7)	6.1 (4.6, 8.1)	0.8 (0.4, 1.9)	84.5 (81.6, 87.0)	707, 857
Black or Black African or Black Caribbean or Black British	10.0 (2.8, 30.1)		16.7 (6.3, 37.4)	13.1 (4.3, 33.5)		60.2 (39.7, 77.6)	26, 20
Asian or Asian British	2.5 (0.4, 15.4)	3.4 (0.6, 16.0)	6.3 (1.9, 19.1)	5.3 (1.4, 18.0)		82.6 (67.9, 91.4)	43, 39

**Stopped or switched contraceptive methods (% (95% CI))**

	<b>Stopped using contraceptives</b>	<b>Switched more &gt; less effective</b>	<b>Switched - less &gt; less effective</b>	<b>Switched - more &gt; more effective</b>	<b>Switched less &gt; more effective</b>	<b>Did not switch or stop usual method</b>	<b>Denominators (weighted, unweighted)</b>
Mixed or multiple or other ethnic groups	1.3 (0.0, 25.2)	5.0 (1.0, 21.9)	15.4 (6.2, 33.4)	8.8 (2.6, 26.0)		69.6 (50.8, 83.5)	31, 39
P-value							p=0.0073

**Self-described sexual identity**

Heterosexual or Straight	3.5 (2.4, 5.1)	2.0 (1.2, 3.3)	4.0 (2.9, 5.7)	6.3 (4.8, 8.2)	0.7 (0.3, 1.7)	83.4 (80.6, 85.9)	771, 844
Lesbian, Gay, Bisexual or Other	5.9 (1.2, 23.9)	4.4 (0.7, 22.7)	4.5 (0.8, 22.8)	15.0 (5.7, 33.8)	0.9 (0.0, 34.0)	69.3 (49.8, 83.7)	29, 104
P-value							p=0.015

**Social Grade**

A/B - Higher/intermediate managerial, administrative and professional	2.8 (1.2, 6.2)	3.3 (1.5, 6.9)	2.7 (1.1, 6.1)	8.0 (5.0, 12.7)	0.4 (0.1, 3.5)	82.8 (76.9, 87.4)	200, 251
C1/C2 - Supervisory, clerical and junior managerial, administrative and professional, skilled manual workers	4.4 (2.8, 6.8)	1.5 (0.7, 3.1)	3.0 (1.8, 5.1)	6.2 (4.3, 8.8)	0.6 (0.2, 1.9)	84.4 (80.7, 87.5)	441, 496
D/E - Semi-skilled and unskilled manual, casual, lowest grade and unemployed	2.5 (0.9, 6.3)	2.1 (0.8, 5.9)	8.8 (5.4, 14.2)	5.8 (3.1, 10.6)	1.6 (0.5, 5.1)	79.2 (72.4, 84.7)	169, 210
P-value							p=0.020

**Education level**

Degree	3.4 (2.1, 5.6)	1.9 (1.0, 3.7)	3.5 (2.1, 5.7)	5.5 (3.7, 8.1)	1.0 (0.4, 2.5)	84.6 (81.0, 87.7)	444, 529
Below degree	3.8 (2.2, 6.4)	2.3 (1.2, 4.6)	5.2 (3.3, 8.1)	7.6 (5.2, 10.9)	0.2 (0.0, 2.0)	80.9 (76.4, 84.7)	349, 408

**Stopped or switched contraceptive methods (% (95% CI))**

	<b>Stopped using contraceptives</b>	<b>Switched more &gt; less effective</b>	<b>Switched - less &gt; less effective</b>	<b>Switched - more &gt; more effective</b>	<b>Switched less &gt; more effective</b>	<b>Did not switch or stop usual method</b>	<b>Denominators (weighted, unweighted)</b>
No qualifications	2.6 (0.1, 39.0)			12.9 (3.2, 40.2)	5.0 (0.5, 35.0)	79.4 (52.4, 93.1)	17, 20
P-value							p=0.21

**Living together - relationship but not living together - Single**

Married/steady and living together	3.8 (2.5, 5.8)	1.2 (0.5, 2.5)	3.7 (2.4, 5.6)	6.1 (4.3, 8.4)	0.8 (0.3, 2.0)	84.5 (81.2, 87.3)	547, 638
Married/steady NOT living together	4.0 (1.6, 9.6)	3.9 (1.5, 9.5)	3.9 (1.5, 9.5)	8.3 (4.4, 15.0)		80.0 (71.6, 86.4)	115, 142
Casual, new, >1, at end or other	0.7 (0.0, 14.5)	4.9 (1.5, 14.8)	10.2 (4.5, 21.4)	4.5 (1.3, 14.4)	1.4 (0.2, 11.8)	78.3 (65.6, 87.3)	58, 68
Single	3.7 (1.3, 10.3)	3.3 (1.1, 9.8)	3.4 (1.1, 10.0)	8.6 (4.3, 16.5)	0.9 (0.1, 7.6)	80.1 (70.6, 87.2)	91, 109
P-value							p=0.20

**Been furloughed under the Coronavirus Job Retention Scheme**

No	3.1 (2.0, 4.7)	2.3 (1.4, 3.7)	4.2 (2.9, 6.1)	7.3 (5.5, 9.5)	0.5 (0.2, 1.5)	82.7 (79.6, 85.4)	656, 775
Yes	5.9 (3.1, 11.1)	1.1 (0.2, 4.9)	4.0 (1.8, 8.7)	3.6 (1.6, 8.2)	1.8 (0.6, 5.9)	83.5 (76.7, 88.7)	150, 177
P-value							p=0.072

**Became unemployed**

No	3.4 (2.3, 4.9)	2.1 (1.3, 3.5)	4.0 (2.8, 5.7)	6.4 (4.8, 8.4)	0.7 (0.3, 1.7)	83.4 (80.5, 86.0)	726, 852
----	----------------	-------------------	-------------------	----------------	-------------------	----------------------	----------



**Stopped or switched contraceptive methods (% (95% CI))**

	<b>Stopped using contraceptives</b>	<b>Switched more &gt; less effective</b>	<b>Switched - less &gt; less effective</b>	<b>Switched - more &gt; more effective</b>	<b>Switched less &gt; more effective</b>	<b>Did not switch or stop usual method</b>	<b>Denominators (weighted, unweighted)</b>
Yes	5.9 (2.4, 13.8)	1.2 (0.2, 8.5)	5.7 (2.3, 13.6)	8.6 (4.1, 17.2)	1.1 (0.1, 8.6)	77.5 (67.0, 85.5)	80, 100
P-value							p=0.61

**Number of days drinking in past week**

0 days	4.2 (2.5, 7.0)	2.4 (1.2, 4.8)	2.1 (1.0, 4.4)	6.6 (4.3, 9.9)	0.3 (0.0, 2.2)	84.5 (80.1, 88.0)	321, 375
1-2 days	3.4 (1.9, 5.9)	1.6 (0.7, 3.7)	3.8 (2.2, 6.4)	7.3 (4.9, 10.6)	0.8 (0.2, 2.6)	83.2 (78.8, 86.9)	334, 399
3-4 days	2.1 (0.5, 7.5)	2.5 (0.7, 8.0)	8.6 (4.5, 15.7)	5.5 (2.4, 11.8)	2.5 (0.7, 8.0)	78.9 (70.1, 85.7)	107, 121
5-7 days	4.6 (1.2, 16.5)	2.1 (0.3, 14.4)	11.4 (4.8, 24.5)	3.7 (0.8, 15.5)		78.2 (63.8, 88.0)	47, 60
P-value							p=0.038

**Drinking habits compared to pre Covid-19 outbreak**

Less these days	2.9 (1.4, 6.0)	3.0 (1.5, 6.2)	3.8 (2.0, 7.1)	7.5 (4.7, 11.6)	1.4 (0.5, 4.0)	81.4 (75.9, 85.9)	239, 294
About the same	4.2 (2.6, 6.6)	1.5 (0.7, 3.2)	3.5 (2.1, 5.8)	5.8 (4.0, 8.5)	0.2 (0.0, 1.7)	84.8 (81.0, 87.9)	421, 488
More these days	3.1 (1.2, 7.7)	2.3 (0.8, 6.6)	6.6 (3.5, 12.0)	7.5 (4.1, 13.1)	1.2 (0.3, 5.2)	79.3 (71.9, 85.2)	144, 168
P-value							p=0.42

**Current smoker**

**Stopped or switched contraceptive methods (% (95% CI))**

	<b>Stopped using contraceptives</b>	<b>Switched more &gt; less effective</b>	<b>Switched - less &gt; less effective</b>	<b>Switched - more &gt; more effective</b>	<b>Switched less &gt; more effective</b>	<b>Did not switch or stop usual method</b>	<b>Denominators (weighted, unweighted)</b>
No	3.5 (2.3, 5.3)	1.8 (1.0, 3.2)	3.2 (2.1, 4.9)	6.4 (4.7, 8.6)	0.8 (0.3, 1.9)	84.2 (81.2, 86.9)	636, 752
Yes	3.8 (1.8, 8.0)	3.0 (1.3, 7.0)	7.6 (4.4, 12.6)	7.2 (4.2, 12.2)	0.5 (0.1, 4.1)	77.9 (71.0, 83.5)	173, 203
P-value							p=0.11

**Symptoms of depression (PHQ2 score)**

No symptoms of depression (0-2)	4.0 (2.6, 6.0)	1.5 (0.8, 3.0)	2.5 (1.5, 4.2)	4.9 (3.4, 7.1)	0.6 (0.2, 1.7)	86.5 (83.3, 89.1)	536, 628
Symptoms of depression (3-6)	2.8 (1.4, 5.7)	3.2 (1.6, 6.1)	6.9 (4.4, 10.7)	9.3 (6.4, 13.5)	0.8 (0.2, 3.0)	76.9 (71.4, 81.6)	266, 319
P-value							p=0.00086

**Symptoms of anxiety (GAD2 score)**

No symptoms of anxiety (0-2)	3.8 (2.4, 5.9)	1.8 (0.9, 3.4)	2.6 (1.5, 4.5)	6.6 (4.7, 9.1)	0.8 (0.3, 2.1)	84.4 (81.0, 87.3)	507, 584
Symptoms of anxiety (3-6)	3.3 (1.8, 6.1)	2.3 (1.1, 4.8)	6.0 (3.8, 9.3)	6.6 (4.3, 10.1)	0.7 (0.2, 2.7)	81.1 (76.2, 85.2)	298, 367
P-value							p=0.26

246 respondents (17.4% of total) answered 'Not applicable' to questions of usual contraception used as they were already pregnant, planning to get pregnant or unable to get pregnant. 142 respondents (9.5%) answered 'no method used' in the year before lockdown. These responses are excluded from the table.