

Primary care providers' knowledge, attitudes and practices of medical abortion: a systematic review

Asvini K Subasinghe , Seema Deb, Danielle Mazza

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjshr-2019-200487>).

Department of General Practice, Faculty of Medicine, Nursing and Health Sciences, Monash University, Notting Hill, Victoria, Australia

Correspondence to

Dr Asvini K Subasinghe, Department of General Practice, Faculty of Medicine, Nursing and Health Sciences, Monash University, Notting Hill, VIC 3168, Australia; asvini.subasinghe@monash.edu

Received 2 October 2019
Revised 13 December 2019
Accepted 14 December 2019



© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Subasinghe AK, Deb S, Mazza D. *BMJ Sex Reprod Health* Published Online First: [please include Day Month Year]. doi:10.1136/bmjshr-2019-200487

ABSTRACT

Background Despite the availability of medical abortifacients, and their potential use in primary care, only a small proportion of primary healthcare professionals provide medical abortion services. Understanding the perspectives of primary care providers on delivering medical abortion is pertinent to identifying barriers to medical abortion service provision and increasing access for women globally.

Objective To understand the knowledge, attitudes and practices of primary healthcare providers regarding medical abortion services.

Design Four databases (Medline, EMBASE, Web of Science (WOS) and Scopus) were searched using search terms related to medical abortion and primary care. The Joanna Briggs Institute Critical Appraisal tools were used to appraise the methodological quality of studies included.

Results Some 22 studies were identified, conducted across 15 countries, comprising 6072 participants. Study participants comprised doctors and residents (n=8), nurses and nursing students (n=5), and pharmacists (n=3) and six studies were conducted with mixed samples of providers. Medical abortion was deemed acceptable by some doctors, but fear of criminal prosecution, in countries where abortion is still restrictive, left doctors and nurses circumspect about providing medical abortion. Pharmacists referred women to other providers with only a small proportion dispensing medical abortifacients. General practitioners, nurses and trainees had mixed knowledge of medical abortion and emphasised the need for training on delivery of medical abortion and dissemination of guidelines. Conversely, pharmacists reported poor knowledge regarding medical abortion regimens and complications.

Conclusions Increased dissemination of training and resources is pertinent to supporting primary care providers delivering medical abortion services and to increasing access for women on a global scale.

Key messages

- Within the primary care setting, doctors are more supportive of medical abortion service provision than are nurses and pharmacists.
- Fear of criminal prosecution and conservative attitudes towards abortion impacted decisions on whether health professionals provided the service.
- Most primary care providers have poor knowledge of medical abortion service provision and would benefit from increased training and education.

INTRODUCTION

Provision of medical abortion services through the primary care sector has the potential to increase access for women globally. General practitioners (GPs)/family physicians and nurses are ideally positioned to deliver medical abortions because they are the first point of contact for women with contraception and pregnancy-related issues.

Approximately one in four women will choose to terminate a pregnancy in their lifetime.¹ Globally, the estimated abortion rate between 2010 and 2014 was 35 per 1000 women aged 15–44 years,² which was 5 per 1000 women less than estimates in 1990–1994. However, no change in the abortion rate was seen in the developing world. The constant rate of abortions in low- and middle-income countries is largely attributable to poor access to effective contraception.³

One of the most important developments in terms of increasing access to abortion is the practice of medical abortions.⁴ Medical abortion is a safe and non-invasive alternative to surgical abortion used in the first trimester of pregnancy. The World Health Organization (WHO)

recommends a medical abortion regimen comprising mifepristone and misoprostol.⁵ When this regimen is administered ≤ 63 days' gestation, the success rate of termination is approximately 98%.⁶ Additionally, this regimen is approved by the US Food and Drug Administration for use up to 70 days' gestation with a growing body of evidence indicating continued effectiveness in later gestations.⁷

While GPs, nurses and pharmacists can provide services and medication for women choosing medical abortion, it is still unclear whether medical abortion provision has been successfully integrated into the primary care setting. Understanding the perspectives of primary care providers on medical abortion service delivery is pertinent to identifying barriers and enablers to the successful integration of medical abortion into general practice and is critical to increasing access to these services for women globally.

Limited data show that GPs and pharmacists perceive medical abortion to be out of their scope or practice and that those delivering the service feel stigmatised, under-resourced and unsure about side effects associated with the procedure.^{8–10} However, to date there has been no systematic review of the literature relating to the knowledge, perceptions and behaviours of primary care providers on medical abortion service delivery.

The aim of this systematic literature review was to identify, assess and synthesise research addressing primary care providers' knowledge, attitudes and practices (KAP) towards providing medical abortions.

METHODS

Search strategy

We searched four databases (Medline, EMBASE, Web of Science (WOS) and Scopus) and manually searched reference lists of included papers. Our search was conducted between March and May 2019 and included all peer-reviewed articles with no restrictions on year of publication. The search strategy used to retrieve articles from Scopus is replicated here: TITLE-ABS-KEY (“induced abortion” OR “medical abort*” OR “mifepristone” OR “misoprostol” OR “methotrexate” OR “unintended pregnanc*”) AND TITLE (“knowledge” OR “perception*” OR “attitude*”) AND TITLE (“primary care*” OR (“general pract*” OR (“doctor*”) OR (“physician*”) OR (“pharmac*”) OR (“nurs*”)) AND (LIMIT-TO (LANGUAGE, “English”)).

The systematic search covered three topics: (1) induced abortion, (2) healthcare provider and (3) knowledge, attitudes and practices. For each of the topics, relevant medical subject headings (MeSH), alternate spellings and truncated versions of words were compiled. All search terms were then combined using ‘OR’ to provide a large range of studies for each theme. The three lists of search terms for each of the topics were then combined with ‘AND’ to generate

high-sensitivity and low-specificity citations that were relevant to all three elements of the research question. The reference lists of the retrieved articles were screened to identify further relevant papers. Each search was limited to studies in English.

Inclusion criteria

Studies were included in this review if they met the following criteria: all primary qualitative and quantitative research studies in which surveys, in-depth interviews, focus group discussions or self-administered questionnaires were used to investigate the KAP of medical abortion services among primary care providers such as GPs, practice nurses and pharmacists as well as students in these professions. Studies were excluded from analysis if KAP were assessed among medical and pharmacy students, non-primary care health providers, or patients, in relation to medical abortion services, or if outcome measures related to emergency contraception use.

Selection of studies

The first author (AKS) selected and collected the articles for the review. All records identified from the electronic searches were then imported into bibliographic software Covidence for reference management. All duplicates were deleted, and a single copy of each record was retained. Titles and abstracts were reviewed to determine whether studies met the inclusion criteria. Full-text articles were then obtained and analysed according to the inclusion criteria. Each article was independently reviewed by two of the authors (AKS and SD) for inclusion. Differences of opinion were resolved by discussion.

Assessment of the quality of the study and data extraction

To assess the methodological quality of the included studies, we used the Joanna Briggs Institute Critical Appraisal tools¹¹ commonly used for assessing qualitative and quantitative studies for inclusion in reviews. Studies that did not meet the pre-set methodological quality criteria were excluded (online supplementary appendix 1).

Following the appraisal of the included studies, we extracted the following information: author, year, region and country, aim, study population, and method of data collection.

Synthesis

The results from the included studies were analysed using content analysis that has previously been used for synthesising results in systematic literature reviews of qualitative and quantitative studies.¹² The 22 studies included in this review were first assessed to categorise key descriptive themes. The key descriptive themes were then systematised in a matrix, and similarities, differences and contradictions were examined.




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of search strategy and selection process.

Analytical themes were developed to answer our question about the KAP of primary care providers towards medical abortions.

RESULTS

Search results and study characteristics

From 342 studies identified, 192 duplicates were removed (figure 1). Titles and abstracts for the remaining 150 studies were reviewed, and then 61 study abstracts were considered for further review. After conducting a risk of bias assessment for articles that met the inclusion criteria, one article was removed due to receiving a high risk of bias score¹³ (online supplementary appendix 1). Subsequently 22 studies were included in this review. Overall, these 22 studies provided a total of 6072 participants (table 1).

The 22 studies included in the review were published between 1981 and 2019 and described healthcare professionals' perceptions of and attitudes towards medical abortions (table 1). The studies were conducted in 15 countries: Italy and France,¹ Turkey,² Northern Ireland,¹ Scotland,¹ USA,⁷ Honduras, Mexico, Nicaragua and Puerto Rico,¹ Australia,⁴ Bangladesh,¹ Senegal,¹ Kenya² and Vietnam.¹

In eight of the studies^{9 14-20} researchers investigated KAP of medical abortion among doctors and medical interns. Five studies²¹⁻²⁵ were conducted on samples of nurses and nursing students, three research groups^{8 26 27} investigated KAP in pharmacists and six studies²⁸⁻³³ included data from mixed groups of health professionals.

In 12 of the studies^{14 15 18-25 32 33} researchers used surveys as their method of data collection. Interviews were used in five studies,^{17 27 28 30 31} and in the remaining five studies^{8 9 16 24 26 29} multiple methods of data collection were used such as focus group discussions, in-depth interviews, surveys and mystery client visits to pharmacies.

Primary care provider KAP data were further categorised into analytical themes based on content.

Poor knowledge of medical abortion provision

Primary care providers generally were unaware of the efficacy and possible adverse effects related to medical abortion.^{8 9 14 26 27} Some GPs believed that uterine perforation, cervical laceration and pelvic infection were complications related to medical abortion.¹⁴ Internists from California had poorer knowledge about the safety and efficacy of mifepristone compared with family practitioners (59% vs 90%, $p < 0.004$).²⁰ Similarly, nursing students from Italy and Sweden²⁴ and nurses in a Kenyan study demonstrated deficient knowledge in all aspects of induced abortion.²¹ The majority of pharmacists in Vietnam knew medical abortifacients by trade names rather than generic names⁸ whereas those practising in Kenya and Senegal had poor knowledge of mifepristone and misoprostol treatment regimens.^{26 27}

Dual influence of legislative restrictions on abortion

Legislative restrictions were perceived to be both a frustration and a necessity, depending on the location of the health professional. The majority of primary healthcare providers believed that medical abortion should be available on request.^{15 18 21-23 29 33} Some providers considered access to medical abortion as a right for women who requested it regardless of the law.^{16 22 30 32} In one study, GPs in Queensland, Australia, where abortion was restrictive at the time, were frustrated with legislative restrictions on abortion. They commented that these restrictions not only limited their confidence in providing abortions but also created gaps in their medical training.¹⁷ However, family physicians in rural Idaho, USA were strongly opposed towards providing RU-486 to their patients due to conservative political and religious views which were shared by the community in which they practised.¹⁹ This opposition was also shared by nurse-midwives ($n = 510$) in the USA, 66% of whom said they would leave their place of work if they were allowed to perform abortions.²²

Table 1 Characteristics of studies included in the review

Author(s)	Year	Countries	Abortion legislation*	Aims/purpose	Study population	Methodology
Agostino <i>et al</i> ²³	1991	Italy Sweden	A, W12 A, W18	To describe attitudes and knowledge concerning abortion and FP among nursing students in Italy and Sweden	229 nursing students 105 Swedish 124 Italian	Self-administered questionnaire in classrooms (20–25 students at a time) + open-ended discussion between researcher and classroom
Akin <i>et al</i> ¹⁴	2012	Turkey	A, W10	To better understand the knowledge, attitudes and perspectives on medical abortion held by physicians	268 GPs 187 obgyns	Self-administered questionnaire
Black <i>et al</i> ¹⁵	2001	Northern Ireland	B	To elicit attitudes of doctors in Northern Ireland, practising obstetrics and gynaecology or FP	16 FP doctors 48 junior obgyn 38 consultant obgyn	Questionnaire
Dawson <i>et al</i> ¹⁶	2017	Australia	C	To explore the provision and referral of MTOPs by GPs in NSW, Australia	32 GPs 24 non-MTOP GP providers 8 MTOP GP providers	28 interviews 1 focus group (4 GPs)
De Moel-Mandel ³²	2019	Australia	C	To identify enablers and barriers to the provision of medication abortion in the primary healthcare setting of regional and rural areas of Victoria, Australia.	39 GPs 30 primary healthcare nurses	Questionnaire
Douglas <i>et al</i> ¹⁷	2013	Australia	C	To explore doctors' understanding of the law related to the provision of abortion in these two states and their views about the effect of the law on their practice	15 QLD doctors 7 NSW doctors	Interview – 10 scenarios
Espinoza <i>et al</i> ⁹	2004	Mexico Puerto Rico Honduras Nicaragua	D A, W12 E E	To examine physicians' knowledge and attitudes with regard to medication abortion	43 Mexican: 93% physicians 13 Puerto Rico: 78% physicians 21 Honduras: 93% physicians 23 Nicaragua: 63% Physicians	8 focus group discussions with 70 physicians Focus group: ▶ survey ▶ presentation on MTOP for common knowledge ▶ discussion
Grindlay <i>et al</i> ²⁰	2012	Iowa, USA	C	To evaluate patients' and providers' experiences with telemedicine provision of medical abortion aimed at learning more about the acceptability of telemedicine, abortion services and the impact that it has on patients, staff and clinic operations	20 telemedicine patients 5 inpatient MTOP patients 15 clinic staff	In-depth interview
Hwang <i>et al</i> ³¹	2005	California, USA	C	To assess the potential for advanced practice clinicians' provision of medical abortion	762 nurse practitioners	Questionnaire
Kidula <i>et al</i> ²⁰	1992	Kisii district, Kenya	B	To determine nurses' knowledge of various aspects of induced abortion, attitude towards induced abortion and abortion patients, and their involvement in abortion	218 nurses	Self-administered questionnaire: partially coded and open-ended questions under supervision of trained research assistant
McKee <i>et al</i> ²¹	1994	USA	C	To determine the attitudes of nurse midwives towards performing abortions	1208 nurses	Self-administered questionnaire

Continued

Table 1 Continued

Author(s)	Year	Countries	Abortion legislation*	Aims/purpose	Study population	Methodology
Miller <i>et al</i> ¹⁸	1998	USA	C	To document practices and attitudes of physicians regarding adolescent abortion and contraception, as well as physician willingness to prescribe medical abortion if approved by the FDA	541 paediatricians 46 family medicine 38 internal medicine 22 obgyn 15 medicine – paediatrics 4 psychiatry 2 other	Self-administered questionnaire (21 items)
Newton <i>et al</i> ²⁹	2016	Victoria, Australia	C	To explore factors that contribute to Australian women's decision to have a surgical or medical abortion, through the perspective of abortion service providers	4 GPs 3 obgyn 2 primary healthcare nurses 1 psychologist 1 sexual health physician 1 medical practitioner 3 service managers	Semi-structured interview
Ngo <i>et al</i> ⁸	2012	Ho Chi Minh City, Vietnam	F	To investigate pharmacy workers' knowledge and provision of abortifacients in Ho Chi Minh City, Vietnam	100 pharmacy workers Mystery client: 30 pharmacies visited	Survey + mystery client visit
Obiadullah <i>et al</i> ²⁸	1981	Bangladesh, all districts except Chittagon Hill Tracts (inaccessible)	D	To investigate attitudes towards abortion of physicians working in rural health complexes and subdivision and district hospitals	376 physicians	Interviews – eight scenarios, field visits
Purcell <i>et al</i> ²⁷	2017	Scotland (urban)	A, W24	To make visible the contemporary body work of abortion and the challenges which accompany it in the context of these shifts, and to use this context to explore the conceptual boundaries of body work	17 nurses 8 doctors 7 clinical support workers 5 sonographers	Semi-structured interview
Reiss <i>et al</i> ²⁵	2016	Nairobi, Mombasa and Kisumu, Kenya	B	To assess pharmacy workers' knowledge and provision of abortion information and methods in Kenya	235 pharmacy workers	Interview Mystery client visits to pharmacies
Reiss <i>et al</i> ²⁶	2017	Dakar, Senegal	E	To understand knowledge and provision of misoprostol among pharmacy workers	110 pharmacy workers	Interview
Rosenblatt <i>et al</i> ¹⁹	1995	Rural Idaho, USA	C	To determine why rural physicians are unwilling to provide abortions	114 family physicians 9 obgyn 15 general surgeons	Survey
Schwartz <i>et al</i> ²⁰	2005	California, USA	C	To assess what internal medicine residents know about mifepristone and whether they intend to incorporate medication abortion into their clinical practices	158 internists 30 family physicians 23 gynaecologists	Self-administered questionnaire
Swenson <i>et al</i> ²²	1994	5 states in the USA (not specified)	C	To learn more about the abortion-related experiences and value orientation of nurses	844 community health nurses	Self-administered questionnaire
Yanikkerem <i>et al</i> ²⁴	2018	Turkey	A, W10	To evaluate Turkish nursing students' attitudes towards voluntary induced abortion	1089 students	Three-part questionnaire

Abortion legislation categories. A: On request: gestational limits vary. B: To preserve a woman's physical, mental and social well-being. C: Federal system in which abortion law restrictions differ at state level. D: To preserve a woman's life. E: Prohibited. F: No restriction.

*Source: Center for Reproductive Rights (2019) World Abortion Laws 2019 <https://reproductiverights.org/sites/default/files/documents/World-Abortion-Map.pdf>.

FDA, Food and Drug Administration; FP, family planning; GP, general practitioner; MTOP, medical termination of pregnancy; NSW, New South Wales; obgyn, obstetrician gynaecologist; QLD, Queensland; USA, United States of America; W10, gestational limit of 10 weeks; W12, gestational limit of 12 weeks; W18, gestational limit of 18 weeks; W24, gestational limit of 24 weeks.

Fear and concern

Fear of being labelled an abortion doctor and risk of criminal prosecution were reported to be reasons for

not providing medical abortion services by GPs in Australia.^{16 17} Similarly, doctors in a study conducted

in South America, reported to offer mifepristone and methotrexate clandestinely because only surgical abortion was legal.⁹ These doctors also asserted that they had a fear relating to tele-abortion: the risk of potential misuse and self-medication by women, if mifepristone were to become approved in their countries.⁹ In another study conducted in Scotland on a small sample of nurses and doctors there was also a shared concern that women may be unable to cope with seeing an expelled embryo if abortion drugs were self-administered.²⁸ Interestingly, from the lens of the provider, internists and family practitioners from California reported concerns about receiving adequate 'back up' access to vacuum aspiration services, lack of access to ultrasound and management of bleeding

Responsibility, scope of practice and demand

There were mixed opinions on who should be responsible for providing medical abortion services. Family planning doctors in Northern Ireland believed gynaecologists should be responsible,¹⁵ while some GPs in Australia asserted that dedicated abortion clinics would provide a better service to women than those working in primary care.¹⁶ Conversely, in one study conducted in the USA in 1998, with 46 family planning doctors, more than 50% believed that if medical abortion was routinely available that it should be available from primary care physicians.¹⁸ Some nurses were comfortable providing medical abortion services if given the opportunity,^{23 32} but most referred women to other providers, similar to practices among pharmacists in Africa.^{22 26 27}

Despite demand for medical abortion being reported as low in two studies conducted with GPs in Australia,^{16 17} these GPs still contended that they had concerns about being inundated with requests for medical abortion which would dominate their scope of practice.¹⁶

Non-conventional medical abortion models of care

Some alternative medical abortion practices were noted in studies relating to telemedicine and traditional culturally specific methods. In one study conducted in Iowa, USA telemedicine was seen to be acceptable to providers in terms of having a greater reach of physicians and more efficient use of resources compared with standard medical abortion service provision.³¹ The only challenges cited were documenting the Rhesus status of patients and organising anti-D injections, and amalgamating patient documentation. Conversely, doctors in Honduras, Mexico, Nicaragua and Puerto Rico did not support telemedicine in their countries because they suspected that women would misuse the drugs at home.⁹ Interestingly these same doctors reported prescribing herbs and teas to induce an abortion or facilitate a complete termination.⁹

Need for resources

All primary care providers stated that there was a need for more resources and training materials related to medical abortion service provision.^{9 16 17 33} Lack of training opportunities was cited as the largest barrier to performing medical abortion or assisting another provider in performing medical abortion by nurse practitioners in California.³² Only a small proportion of nurse practitioners and GPs stated they were not interested in receiving medical abortion training because of the concern for needing surgical back-up.³³ GP providers in Australia discussed gaps in peer support to ensure continuity of care and professional development and that appropriate policies and procedures need to be developed, particularly on patient follow-up, within practices.¹⁶ GPs in South America also reported a lack of information and commented on the poor quality of resources available for them on medical abortion.⁹ Suggestions for strategic dissemination of information included public information campaigns, creating an informal website for health professionals about medical abortion, and distributing printed materials among doctors.⁹

DISCUSSION

From our review of 22 studies including 6072 participants, we found that GPs were largely supportive of medical abortion services being available to women, while nurses and pharmacists reported mixed attitudes. Most primary care providers demonstrated relatively poor knowledge of processes and noted that more training and educational materials on medical abortion service provision would be welcomed. Some primary care providers practising in countries where abortion is still restrictive had concerns about providing medical abortion services, due to fear of prosecution and stigma. Despite most primary care providers claiming to support medical abortion, this rarely was reflected in their practice as provision was still poor. Finally, we also demonstrate a gap in the literature for information on medical abortion service provision within the primary care setting, as demonstrated by studies on nurses and pharmacists comprising fewer than 30% of studies included in this review.

Knowledge appeared to be relatively poor across all primary care providers. Some GPs were aware of procedures and follow-up care because they provided the service. Poor knowledge was understandably more common among primary care providers practising in countries where abortion is still restrictive such as Turkey, Kenya, Senegal and some countries in South America. Therefore, national colleges of practitioners around the world should advocate on behalf of the evidence to change legislation so that the best-available evidence around medical abortion provision can be implemented in the form of national guidelines and instruction manuals.

Most doctors supported medical abortion provision, but strong opposition to medical abortion, based on conservative or political views, was still prevalent among some doctors and nurses.^{19–22} This opposition is problematic when considering the WHO's recommendation for task-sharing medical abortion models of care whereby doctors and nurses are encouraged to share responsibilities and delegations during provision of medical abortion services to women.³⁴ Therefore, findings from this review support more technical training and values clarification for nurses and midwives to increase provision of safe and high-quality medical abortion care.³⁵

Only a small proportion of primary care health professionals reported providing medical abortion services and dispensing medical abortifacients. Legislative restrictions on abortion, personal objections, and a lack of training and availability of resources were common factors that negatively impacted on service provision. More attention should be given to training and education on personal challenges relating to medical abortion service provision in medical, nursing and pharmacist training. University and college curriculums should comprise clear and comprehensive guidelines on how to manage the ethical dilemmas associated with medical abortion along with training modules on comprehensive medical abortion care, coupled with counselling in family planning and abortion.³⁵

Strengths

To the best of our knowledge, this is the first systematic review to evaluate primary care providers KAP related to medical abortion provision. Studies included in the review spanned several continents, thus providing data on medical abortion service provision in primary care setting within countries with highly disparate legislation on abortion. Additionally, given the strong movement, led by the WHO,⁵ in support of providing medical abortions at the primary care level, this work is pertinent to understanding how we can improve medical abortion service provision within the primary healthcare sector to fundamentally increase access for women.

Limitations

Studies included in this review comprised various methods of data collection and posed different research questions across studies. Most studies included assessments of knowledge and attitudes relating to medical abortion services without direct assessment of practices. Additionally, adjustment for confounding factors such as age, sex and length of employment were rarely considered.

CONCLUSIONS

From this systematic literature review of 22 studies we show that doctors working within the primary care setting are more supportive of medical abortion

Additional Educational Resources

- ▶ Dawson A, Bateson D, Estoesta J, *et al.* Towards comprehensive early abortion service delivery in high income countries: insights for improving universal access to abortion in Australia. *BMC Health Serv Res* 2016;**16**:612.
- ▶ Barnard S, Kim C, Park MH, *et al.* Doctors or mid-level providers for abortion. *Cochrane Database Syst Rev* 2015;**7**:CD011242. DOI: 10.1002/14651858.CD011242.pub2.
- ▶ World Health Organization. *Clinical Practice Handbook for Safe Abortion* [Electronic Handbook]. 2014. https://apps.who.int/iris/bitstream/handle/10665/97415/9789241548717_eng.pdf;jsessionid=F68376B09CA2FA832C14B3C8342DC63C?sequence=1.

service provision compared with nurses and pharmacists. Fear of criminal prosecution and conservative attitudes towards abortion determined whether or not health professionals provided the service and or referrals. We also highlight that the majority of primary care providers have poor knowledge of medical abortion service provision and would benefit from quality training and education. Changes in legislation and accessibility to resources are required globally to ensure that primary care providers have the confidence and training to perform abortions safely and become exemplars of abortion advocacy in their respective countries.

Contributors AKS and DM were responsible for the conception and design of the study. AKS conducted the literature search and extracted the data. AKS and SD independently reviewed articles for inclusion. AKS analysed and interpreted the data and wrote the manuscript. All authors critically reviewed the manuscript and provided important intellectual content.

Funding This study was funded by the National Health and Medical Research Council (NHMRC), Australia (1153592) Centre of Research Excellence in Sexual and Reproductive Health for Women in Primary Care (SPHERE).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Asvini K Subasinghe <http://orcid.org/0000-0001-6045-2728>

REFERENCES

- 1 Cameron S, Glasier A, Lohr PA, *et al.* Induced abortion. *Hum Reprod* 2017;**32**:1160–9.
- 2 Sedgh G, Bearak J, Singh S, *et al.* Abortion incidence between 1990 and 2014: global, regional, and subregional levels and trends. *Lancet* 2016;**388**:258–67.
- 3 Singh SDJ. *Adding it up: costs and benefits of contraceptive services*. New York, NY, USA: Guttmacher Institute and UNFPA, 2012.

- 4 Guttmacher Institute. Abortion worldwide 2017: uneven progress and unequal access; 2018. <https://www.guttmacher.org/report/abortion-worldwide-2017>
- 5 World Health Organization (WHO). *Medical management of abortion (Licence: CC BY-NC-SA 3.0 IGO)*. Geneva, Switzerland: WHO, 2018.
- 6 Kulier R, Kapp N, Gülmezoglu AM, *et al*. Medical methods for first trimester abortion. *Cochrane Database Syst Rev* 2011;123.
- 7 Food and Drug Administration. *Mifeprex (mifepristone) information*, 2016.
- 8 Ngo TD, Park MH, Nguyen TH. Pharmacy workers' knowledge and provision of abortifacients in Ho Chi Minh City, Vietnam. *Int J Gynecol Obstet* 2012;117:187–8.
- 9 Espinoza H, Abuabara K, Ellertson C. Physicians' knowledge and opinions about medication abortion in four Latin American and Caribbean region countries. *Contraception* 2004;70:127–33.
- 10 Dressler J, Maughn N, Soon JA, *et al*. The perspective of rural physicians providing abortion in Canada: qualitative findings of the BC Abortion Providers Survey (BCAPS). *PLoS One* 2013;8:e67070.
- 11 Joanna Briggs Institute. Critical appraisal tools for use in JBI systematic reviews, 2017. Available: https://joannabriggs.org/sites/default/files/2019-05/JBI_RCTs_Appraisal_tool2017_0.pdf [Accessed 19 Feb 2019].
- 12 Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol* 2008;8:45.
- 13 Serena B, Sullivan P, Stack J, *et al*. Attitudes toward Michigan's abortion law and experience of Michigan physicians with requests for abortion. *Mich Med* 1971;70:309–16.
- 14 Akin A, Dogan BG, Ozvaris SB, *et al*. Introducing medical abortion in turkey: perceptions of physicians. *Int J Gynaecol Obstet* 2012;118:S57–61.
- 15 Black G, Hunter A, Heasley N. A survey of attitudes to abortion law in Northern Ireland amongst obstetricians, gynaecologists and family planning doctors. *J Fam Plann Reprod Health Care* 2001;27:221–2.
- 16 Dawson AJ, Nicolls R, Bateson D, *et al*. Medical termination of pregnancy in general practice in Australia: a descriptive-interpretive qualitative study. *Reprod Health* 2017;14:39.
- 17 Douglas H, Black K, de Costa C. Manufacturing mental illness (and lawful abortion): doctors' attitudes to abortion law and practice in New South Wales and Queensland. *J Law Med* 2013;20:560–76.
- 18 Miller NH, Miller DJ, Pinkston Koenigs LM. Attitudes of the physician membership of the Society for Adolescent Medicine toward medical abortions for adolescents. *Pediatrics* 1998;101:E4.
- 19 Rosenblatt RA, Mattis R, Hart LG. Abortions in rural Idaho: physicians' attitudes and practices. *Am J Public Health* 1995;85:1423–5.
- 20 Schwarz EB, Luetkemeyer A, Foster DG, *et al*. Willing and able? Provision of medication for abortion by future internists. *Women's Health Issues* 2005;15:39–44.
- 21 Kidula NA, Kamau RK, Ojwang SB, *et al*. A survey of the knowledge, attitude and practice of induced abortion among nurses in Kisii district, Kenya. *J Obstet Gynaecol East Cent Africa* 1992;10:10–12.
- 22 McKee K, Adams E. Nurse midwives' attitudes toward abortion performance and related procedures. *J Nurse Midwifery* 1994;39:300–11.
- 23 Swenson I, Swanson J, Oakley D. Community health nurses' perceptions, knowledge, and involvement in abortion services. *Womens Health Issues* 1994;4:219–26.
- 24 Agostino MB, Wahlberg V. Nursing students' attitudes towards abortion and family planning in Italy and Sweden. *Scand J Caring Sci* 1991;5:87–92.
- 25 Yanikkerem E, Üstgörtl S, Karakus A, *et al*. Turkish nursing students' attitudes towards voluntary induced abortion. *J Pak Med Assoc* 2018;68:410–6.
- 26 Reiss K, Footman K, Akora V, *et al*. Pharmacy workers' knowledge and provision of medication for termination of pregnancy in Kenya. *J Fam Plann Reprod Health Care* 2016;42:208–12.
- 27 Reiss K, Footman K, Burke E, *et al*. Knowledge and provision of misoprostol among pharmacy workers in Senegal: a cross sectional study. *BMC Pregnancy Childbirth* 2017;17:211.
- 28 Purcell C, Cameron S, Lawton J, *et al*. The changing body work of abortion: a qualitative study of the experiences of health professionals. *Sociol Health Illn* 2017;39:78–94.
- 29 Obiadullah M, Khan AR, Measham AR, *et al*. Induced abortion in rural Bangladesh: mortality levels and physicians' attitudes. *Rural Demogr* 1981;8:91–120.
- 30 Newton D, Bayly C, McNamee K, *et al*. How do women seeking abortion choose between surgical and medical abortion? Perspectives from abortion service providers. *Aust N Z J Obstet Gynaecol* 2016;56:523–9.
- 31 Grindlay K, Lane K, Grossman D. Women's and providers' experiences with medical abortion provided through telemedicine: a qualitative study. *Womens Health Issues* 2013;23:e117–22.
- 32 Hwang AC, Koyama A, Taylor D, *et al*. Advanced practice clinicians' interest in providing medical abortion: results of a California survey. *Perspect Sex Reprod Health* 2005;37:92–7.
- 33 Moel-Mandel C, Graham M, Taket A. Snapshot of medication abortion provision in the primary health care setting of regional and rural Victoria. *Aust J Rural Health* 2019;22:237–44.
- 34 World Health Organization (WHO). *Who recommendations: optimizing health worker roles to improve access to key maternal and newborn health interventions through task shifting*. Geneva, Switzerland: WHO, 2012.
- 35 Rehnström Loi U, Gemzell-Danielsson K, Faxelid E, *et al*. Health care providers' perceptions of and attitudes towards induced abortions in sub-Saharan Africa and Southeast Asia: a systematic literature review of qualitative and quantitative data. *BMC Public Health* 2015;15:139.