

Sexual behaviour, pregnancy intention and sexually transmitted infection risk varies extensively among transgender and non-binary patients in the UK

There is a dearth of evidence regarding the sexual and reproductive health (SRH) needs and experiences of transgender and non-binary (TNB) people (those whose current gender identity differs from the sex they were assigned at birth).¹ However, the TNB community remain underrepresented in medical research.² This is important, as a key part of gender affirmation treatments include cross-sex hormones and surgery that may limit TNB people's future reproductive capabilities.

We sought to explore pregnancy intention, HIV and sexually transmitted infection (STI) risk, and contraception use of TNB patients, with the intention that the results may guide healthcare provision.

TNB individuals attending a National Health Service (NHS) gender identity clinic in Edinburgh over a 4-month period completed an anonymous self-administered questionnaire. The questionnaire asked about demographic characteristics, pregnancy intention, sexual behaviour, STI/HIV risk and pre-exposure prophylaxis (PrEP) use. Data were recorded and analysed using Microsoft Excel (2007).

Some 65 questionnaires were disseminated between September 2019 and January 2020. A total of 43 questionnaires were returned partially or fully completed (a 66.2% return rate). Thirty-six respondents completed the questionnaire sufficiently to be included in the analysis, giving a completion rate of 55.6%.

Twelve (33.3%) respondents identified as trans men and 18 (50%) identified as trans women. Six (16.6%) identified as non-binary assigned female at birth (NBAFAB). Further data are framed in the context of gender identity. **Table 1** displays information regarding the demographics and sexual behaviours of respondents in relation to gender identity.

Ten (27.8%) respondents wanted a baby now or in the future. Eight (22.2%)

Table 1 Respondents' demographics and sexual behaviour

Division	Subdivision	Respondents (n)	Percentage of total respondents (n=36)
Gender	Transgender man	12	33.3
	Transgender woman	18	50.0
	Non-binary (AFAB)	6	16.6
Ethnicity	White-British	35	97.2
	Mixed White-British/Arabic	1	2.8
Relationship status	Single	23	63.9
	Partnered/married	13	36.1
Accommodation	Temporary rented	19	52.8
	Homeowner	17	47.2
Occupation	Employed	17	47.2
	Unemployed	4	11.1
	Student	13	36.1
	Not fit for work	2	5.6
Education	Standard grade level or higher	36	100
Smoking status	Smoker	9	25.0
	Non-smoker	27	75.0
Alcohol intake	≥2 times per week	8	22.2
	<2 times per week	28	77.8
Fertility	Fertile	28	77.8
	Infertile	8	22.2
			Percentage of each variable
Sexual intercourse among trans men	In the last year	9	75.0
	Not in the last year	3	25.0
Type of sexual intercourse among trans men	Oral	5	55.6
	Penis-vaginal	2	22.2
	Mixed (penis-vaginal, penis-anal)	2	22.2
Sexual intercourse among trans women	In the last year	8	44.4
	Not in the last year	10	55.6
Type of sexual intercourse among trans women	Oral	1	12.5
	Penis-vaginal	4	50.0
	Penis-anal	1	12.5
	Mixed (penis-vaginal, penis-anal)	2	25.0
Sexual intercourse among NBAFAB individuals	In the last year	3	50.0
	Not in the last year	3	50.0
Type of sexual intercourse among NBAFAB individuals	Penis-vaginal	2	66.7
	Oral	1	33.3

NBAFAB, non-binary assigned female at birth.

were identified as being in pregnancy-possible couplings (PPCs), based on fertility status, sexual behaviours and the sex assigned at birth of each partner. However, most of these (n=7, 87.5%) were not using any form of contraception. Of the seven respondents in PPCs

and also not using contraception, five (71.4%) expressed that they were trying to avoid pregnancy.

Sixteen (44.4%) respondents were at risk of STIs/HIV based on self-reported sexual behaviours, with most having no sexual health check-up (n=10, 62.5%)

or HIV test (n=13, 81.3%) in the last year.

Twenty-one (58.3%) respondents had heard of PrEP for the prevention of HIV. Only 5/21 (13.9%) respondents would consider taking it.

The reproductive desires of TNB individuals vary widely and are comparable to those of cisgender individuals.² The majority of our respondents in PPCs were not using contraception, reflecting an incorrect belief among many TNB individuals that gender-affirming hormones act as contraception.³ Those who were identified as being at risk of STIs/HIV were unlikely to have been tested for HIV/STIs in the last year. Disengagement from sexual health services, potentially due to stigma, discrimination and past negative healthcare encounters, may limit the potential to identify and address STI/HIV risk among TNB individuals. As the TNB community are disproportionately affected by STIs/HIV, it is of paramount importance to design sexual health services with this population in mind. PrEP awareness is low among the TNB population, despite a higher prevalence of HIV among this group. Globally, studies have shown that willingness to use PrEP is high among TNB patients.⁴ It is crucial that information regarding PrEP is disseminated effectively throughout the TNB community by sexual healthcare providers.

Even in this small sample, we identified several TNB patients who had unmet SRH needs. Recent UK guidelines provide a framework for the redesign of sexual health services with the TNB population in mind.⁵ Further research is needed to identify how best to engage and support the TNB community.

Alexandra Marion Hague ,¹

John Joseph Reynolds-Wright ,^{2,3}

¹Division of Clinical and Surgical Sciences, Medical School, University of Edinburgh, Edinburgh, UK

²Queen's Medical Research Institute, University of Edinburgh MRC Centre for Reproductive Health, Edinburgh, UK

³NHS Lothian, Chalmers Centre, Edinburgh, UK

Correspondence to Alexandra Marion Hague, Division of Clinical and Surgical Sciences, Medical School, University of Edinburgh, Edinburgh EH8 9YL, UK; alexhague@hotmail.co.uk

Twitter John Joseph Reynolds-Wright @doctorjrw

Acknowledgements The authors want to thank the staff and patients of the Chalmers Centre Gender Identity Clinic for participating in the study. They also want to thank Professors Sharon T Cameron and Richard A Anderson for their advice on the design of the questionnaire.

Contributors AMH and JJRW contributed equally to the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not required.

Ethics approval Ethical approval was obtained from the West of Scotland Research Ethics Committee, Glasgow, UK (Ref. 19/WS/0114).

Provenance and peer review Not commissioned; externally peer reviewed.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.



BMJ Sex Reprod Health 2020;**0**:1–2.
doi:10.1136/bmjshr-2020-200655

ORCID iDs

Alexandra Marion Hague <http://orcid.org/0000-0003-1147-9814>

John Joseph Reynolds-Wright <http://orcid.org/0000-0001-6597-1666>

REFERENCES

- Olson-Kennedy J, Cohen-Kettenis PT, Kreukels BPC, *et al.* Research priorities for gender nonconforming/transgender youth: gender identity development and biopsychosocial outcomes. *Curr Opin Endocrinol Diabetes Obes* 2016;**23**:172–9.
- Hoffkling A, Obedin-Maliver J, Sevelius J. From erasure to opportunity: a qualitative study of the experiences of transgender men around pregnancy and recommendations for providers. *BMC Pregnancy Childbirth* 2017;**17**:1–14.
- Faculty of Sexual & Reproductive Healthcare (FSRH). Clinical effectiveness unit (CEU) statement: contraceptive choices and sexual health for transgender and non-binary people, 2017. Available: <https://www.fsrh.org/documents/fsrh-ceu-statement-contraceptive-choices-and-sexual-health-for/> [Accessed 18 May 2020].
- Pacífico de Carvalho N, Mendicino CCP, Cândido RCF, *et al.* HIV pre-exposure prophylaxis (PrEP) awareness and acceptability among trans women: a review. *AIDS Care* 2019;**31**:1234–40.
- British Association for Sexual Health and HIV (BASHH). BASHH recommendations for integrated sexual health services for trans, including non-binary people, 2020. Available: <http://www.gpone.wales.nhs.uk/sitesplus/documents/1000/bashh-recommendations-for-integrated-sexual-health-services-for-trans-including-non-binary-people-2019pdf.pdf> [Accessed 18 May 2020].