

Table S1. Intention-to-treat analysis of abortion-related variables for medical abortion at 17-20 weeks gestation.

	Immediate Group (n=55)	Delayed Group (n=57)	
Abortion intervals	Median (IQR)	Median (IQR)	P-value ^a
Time to fetal expulsion (Hours)	6.3 (4.8-9.6)	8.3 (6.0-11.0)	0.15
Time from fetal expulsion to abortion completion ^b (Hours)	5.2 (2.9-6.4)	4.3 (1.5-6.5)	0.34
	n (%)	n (%)	
Spontaneous expulsion of fetus	55 (100)	57 (100)	-
Additional procedures following fetal expulsion			0.34
None (spontaneous complete expulsion of placenta)	29 (53)	23 (40)	
Manual removal of retained placenta (no vacuum aspiration performed)	6 (11)	6 (11)	
Vacuum aspiration for retained placenta, products of conception or bleeding	20 (36)	28 (49)	
IUD inserted after vacuum aspiration ^c	15 (33)	-	
Abortion complications			
Given intravenous antibiotics for infection	2 (4)	0 (0)	0.24
Hemorrhage >500 ml	8 (15)	5 (9)	0.34

^a Wilcoxon rank-sum tests for medians. Chi-squared or Fisher's exact test for proportions

^b Defined as expulsion or evacuation of the placenta

^c Includes only those getting the IUD immediately (per protocol)