





OPEN ACCESS

Adherence to treatment and prevalence of side effects when medical abortion is delivered via telemedicine: a prospective observational cohort study during COVID-19

John Joseph Reynolds-Wright ^{1,2}, Anne Johnstone,^{1,2} Karen McCabe,^{1,2} Emily Evans,³ Sharon Cameron ^{1,2}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjsex-2021-201263>).

¹MRC Centre for Reproductive Health, The University of Edinburgh, Edinburgh, UK
²Chalmers Centre, NHS Lothian, Edinburgh, UK

³Edinburgh Clinical Research Facility, The University of Edinburgh, Edinburgh, UK

Correspondence to

Dr John Joseph Reynolds-Wright, The University of Edinburgh MRC Centre for Reproductive Health, Edinburgh, Edinburgh, UK; jjrw@doctors.org.uk

Received 5 July 2021
Accepted 9 October 2021



Check for updates

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Reynolds-Wright JJ, Johnstone A, McCabe K, *et al.* *BMJ Sex Reprod Health* Published Online First: [please include Day Month Year]. doi:10.1136/bmjsex-2021-201263

ABSTRACT

Background The Scottish government introduced legislation during the COVID-19 outbreak to permit medical abortion at home with telemedicine. All women received an initial telephone consultation. For those choosing medical abortion, we provided self-administered medications to eligible women with pregnancies under 12 weeks' gestation.

Aims To assess adherence to the recommended abortion drug regimen, with particular focus on the number of misoprostol doses used and the interval between mifepristone and misoprostol administration and the induction–expulsion interval. Additionally, to evaluate use of analgesia, antiemetics and antibiotics, and the side effects, pain and bleeding profile of medical abortion at home.

Methods We conducted a prospective cohort study of 663 women choosing medical abortion at home via telemedicine at an NHS abortion service in Edinburgh, Scotland between 1 April and 9 July 2020. Interviewer-administered questionnaires were completed at telephone follow-up 4 and 14 days following treatment. Outcome measures were self-reported and included use of mifepristone and misoprostol, induction–expulsion interval (time from misoprostol administration until expulsion of pregnancy), antiemetics, antibiotics, analgesia use, pain scores, rates of side effects, bleeding and preparedness for treatment.

Results Among the respondents, 652/663 women (98%) answered at least one questionnaire, and 594/663 (89.6%) used both abortion medications as directed (24–72 hours between medications). The mean (SD) induction–expulsion interval was 4.3 (4.3) hours. Antiemetics were used by 611/663

Key messages

- The majority of patients self-administer abortion medications correctly following telephone counselling.
- Many experience side effects including pain, which do not appear to be adequately controlled with current analgesia regimens.
- Telemedicine is effective to prepare women for medical abortion at home.

(92%), 383/599 (64%) completed the course of prophylactic antibiotics, and 616/663 (93%) used analgesia, with mean (SD) worst-pain scores of 6.7 (2.2) out of 10. Regarding side effects, 510/663 (77%) experienced either nausea, vomiting, diarrhoea or headache, 101/663 (15%) experienced headache and 510/663 (77%) experienced bleeding that was heavier than a period; 554/663, (84%) felt prepared for their treatment by teleconsultation.

Conclusion Patients are able to correctly self-administer abortion medications following a telemedicine consultation. Further research is required to optimise pain management and gastrointestinal side effects during medical abortion.

INTRODUCTION

Until March 2020, medical abortion care in Britain routinely involved an in-person consultation and ultrasound scan to assess gestation.^{1 2} Administration of mifepristone needed to occur in a registered clinic or hospital, but women with pregnancies

under 10 weeks' gestation could self-administer misoprostol at home.

In response to the COVID-19 pandemic, legislation across the UK changed to allow home use of mifepristone. In England and Wales, gestation was limited to 10 weeks, but in Scotland clinical guidance supported medical abortion at home up to 12 weeks' gestation.³ NHS Lothian, the sole provider of abortion care in Edinburgh and the surrounding region, treats just over 2600 women each year⁴ and is based at Chalmers Centre for Sexual and Reproductive Health. All care and medications are at no cost to the patient, as is the norm in the National Health Service (NHS). As a result of the new legislation and clinical guidance,^{3,5} the service moved wholly to provision of abortion care by telemedicine and without routine ultrasound on 1 April 2020.⁶ Surgical abortion and in-patient medical abortion were locally available for women with pregnancies up to 12 weeks' and 20 weeks' gestation, respectively; however, access was restricted due to COVID-19.

All women received a remote consultation and those who indicated they would prefer medical abortion were provided with verbal information on how to use the medications and directed to the service's online audiovisual resources on abortion care.⁷ Consultations were by telephone and lasted between 30 and 60 min. Women could choose to collect a medication pack from a designated collection point at the clinic (accessible without entering the building) or to receive this via courier at their home. The contents of the medication pack are shown in **Box 1**. The pack contained a detailed information sheet written in simple English and with pictures. Antibiotics were provided to all women as pre-abortion STI testing was not available. Antibiotics were not provided if there was a clinical contraindication or the woman declined.

Women were provided with a direct telephone line to speak to a specialist nurse in the clinic during daytime hours or to the gynaecology ward at the regional hospital overnight. They could call if they had questions or concerns about an aspect of the procedure.

We have previously reported on the high levels of safety, effectiveness and acceptability of medical abortion at home delivered by our telemedicine model,⁶ which are comparable to in-person models of care. Other studies in Britain and the USA have reported similar findings.^{8–10} Yet, there are no studies reporting on adherence to medical abortion drug regimens or the interval between misoprostol administration and expulsion of pregnancy (induction–expulsion interval) when self-administered at home or in the context of a telemedicine-delivered service. Likewise, there are no studies reporting on the rates of side effects, pain score and analgesia use in telemedicine/at-home settings.

The aim of this analysis was therefore to assess adherence to the recommended abortion drug regimen, with particular focus on the number of misoprostol doses used and the interval between mifepristone and misoprostol administration and induction–expulsion

Box 1 Medication pack contents

The medication pack and drug regimens:

- ▶ A single dose of mifepristone 200 mg* to be taken orally and followed 24–72 hours later by
- ▶ An initial dose of misoprostol 800 µg (and up to three supplemental doses of 400 µg) to be administered sublingually, vaginally or buccally as preferred by the patient at 3–4 hourly intervals until expulsion of pregnancy.
- ▶ Five doses of cyclizine 50 mg – women were directed to use one tablet an hour before mifepristone and another an hour before misoprostol.
- ▶ A 7-day course of prophylactic doxycycline 100 mg twice daily, as routine testing for sexually transmitted infections was not available during the early COVID-19 pandemic.†
- ▶ Five doses of dihydrocodeine 30 mg – all women were encouraged to purchase a supply of paracetamol and ibuprofen.
- ▶ Short-acting hormonal contraception or condoms were included if requested.
- ▶ A low-sensitivity urine pregnancy test (positive at 1000 IU/mL human chorionic gonadotrophin) with instructions on how to use this at 2 weeks after misoprostol administration.
- ▶ Detailed written information on the use of the medications.

*The abortion medications were labelled in the order that they should be administered (ie, mifepristone = 1, misoprostol = 2) and with plain language labels.

†Antibiotics were provided to all women unless contraindicated or declined.

interval. We also aimed to evaluate use of analgesia, antiemetics and antibiotics, and the side effects, pain and bleeding profile of medical abortion at home.

METHODS

We conducted a prospective observational study of women having medical abortion at home (<12 weeks' gestation) via telemedicine.⁶ All women receiving telemedicine medical abortion from 1 April 2020 until 9 July 2020 were contacted by a researcher and asked to complete a questionnaire by telephone on days 4 and 14 after misoprostol administration. We contacted women at day 4 as this was when the abortion was likely to be completed and the experience fresh in the woman's mind, without disturbing them during their treatment. We contacted women at day 14 to obtain the result of their post-abortion low-sensitivity pregnancy test. We asked for the result of this to evaluate effectiveness of treatment (reported previously⁶) and used this opportunity to ask further questions. The questionnaires relied on recall only and women were not asked to note timings prospectively or keep a diary.

The day 4 questionnaire asked about intake of the abortion medications, including administration of misoprostol (route, number of doses and timing in relation to mifepristone), perceived timing of pregnancy expulsion, pain score during abortion (worst pain experienced on an 11-point Likert scale, where 0 is no pain and 10 is worst pain imaginable, as well as comparison of pain with expectations), analgesia use, antiemetic use, bleeding (using a 5-point Likert scale in comparison with a usual period, ranging from 'much less than a normal period' to 'much more than a normal period') and side effects (nausea, diarrhoea, vomiting and headache).

The day 14 questionnaire asked about prophylactic antibiotic use, preparedness for treatment (assessed on a 5-point Likert scale from 'very unprepared' to 'very prepared') and reflections on duration of the telephone consultation. See online supplemental file 1 for the full questionnaire and [box 1](#) for the medication regimen used.

As medical abortion at home for women with pregnancies between 10 weeks and 12 weeks' gestation was a novel practice, we have reported these data separately within the results.

The main outcomes were: 1) proportion of women administering mifepristone and misoprostol within the correct time frame (misoprostol administered within 24–72 hours of mifepristone), 2) total number of misoprostol doses used, 3) induction–expulsion interval (time from misoprostol administration until pregnancy tissue passed, when patient certain pregnancy expelled) in hours. Secondary outcomes were worst pain score, analgesia regimens used, rates of anti-emetic use, rates of compliance with prophylactic antibiotic, bleeding duration and side effect profiles (nausea, diarrhoea, vomiting and headache).

Statistics

An independent statistical consultant performed all statistical analyses using SAS Enterprise Guide v 7.15 (SAS Institute Inc., Cary, North Carolina, USA) and Microsoft Excel 2016. Descriptive statistics with proportions are presented.

Approvals

The project received approval from the NHS Lothian Sexual and Reproductive Health Service quality improvement team and was not deemed to require ethical approval following review by the local NHS research ethics committee scientific officer.

Patient and public involvement

Patients and members of the public were not directly involved in the design of this study.

RESULTS

In the study period, 826 women had a teleconsultation. Sixty-eight women did not proceed to have an abortion following consultation. Of the remaining

758 women who proceeded to abortion, 663 (87%) had a medical abortion at home and were included in the study cohort.⁶ Almost all (n=652, 98.3%) the women provided responses to at least one follow-up questionnaire. Complete questionnaires at both day 4 and day 14 post-abortion were available for 605 (91.3%) women. Forty-five (6.8%) women responded to the day 4 contact only and 2 (0.3%) responded only to the day 14 contact. The full demographics and characteristics of the cohort are reported in full elsewhere.⁶

Use of abortion medications

Mifepristone was administered by 649/663 women (98%). Two women (0.3%) did not use mifepristone and 12 women (1.8%) did not respond to the questionnaire. Misoprostol was used by 594/649 patients (92%) at the recommended interval following mifepristone administration; 542/663 (82%) used a single dose of misoprostol 800 µg and 89/663 (13%) used one further 400 µg dose. A total of 460/663 (69%) administered misoprostol sublingually. The mean (4.3 hours) and median (3 hours) induction–expulsion intervals were similar and calculated for women who were certain they had passed the pregnancy (those who were not, did not provide timings). [Table 1](#) presents this in more detail and the subset of patients with pregnancies between 10 weeks and 11 weeks 6 days' gestation. The proportion of women using medications correctly, mode of administration and induction–expulsion interval were similar in the higher and lower gestation groups.

Use of antibiotic and antiemetic medications

Antibiotic prophylaxis (doxycycline) was provided to 599/663 (90%) of patients. Of these, 383/599 (64%) took the full course as directed and 53/599 (9%) used some of the antibiotics. Antiemetic use was reported by 611/663 (92%) patients and of those who gave the total number of doses used, the mean was 2.1 doses. Further detail is provided in [table 2](#).

Experience of pain and use of analgesia

At day 4, 624/663 (94%) patients reported that they had experienced pain and 616/663 (93%) used at least one form of analgesic medication. The mean (SD) score for worst pain experienced (by day 4) was 6.7 (2.2). Compared with expectations, 375/663 (57%) rated their pain as better or the same as they were expecting. For analgesia, 610/663 (92%) patients used paracetamol (median total dose during abortion process was 2000 mg, 403/663 (61%) used ibuprofen (median total dose 800 mg), and 427/663 (64%) used dihydrocodeine (median total dose 60 mg); 34/663 patients (5%) used at least one additional form of analgesia that they had obtained outside of the clinic or already had in their possession, including cannabis or derivatives (n=15), co-codamol preparations (n=12), tramadol (n=5), other non-steroidal drugs (n=2) and buprenorphine (n=1) (NB. Some patients used more

Table 1 Use of abortion medications

Route of misoprostol administration	Entire cohort (n=663)	Gestation under 10 weeks (n=642)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
Sublingual	460 (69.4%)	444 (69.2%)	16 (76.2%)
Vaginal	170 (25.6%)	166 (25.9%)	4 (19.0%)
Buccal	1 (0.2%)	1 (0.2%)	0
Unknown	32 (4.8%)	31 (4.8%)	1 (4.8%)
Doses of misoprostol	Entire cohort (n=663)	Gestation under 10 weeks (n=642)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
800 µg dose only	544 (82.1%)	529 (82.4%)	15 (71.4%)
800 µg dose and 1×400 µg further dose	89 (13.4%)	84 (13.1%)	5 (23.8%)
800 µg dose and 2×400 µg further doses	14 (2.1%)	13 (2.0%)	1 (4.8%)
Unknown	16 (2.4%)	16 (2.5%)	0
Used medications correctly?*	Entire cohort administering mifepristone (n=649)	Gestation under 10 weeks administering mifepristone (n=628)	Gestation 10 weeks – 11 weeks and 6 days administering mifepristone (n=21)
Yes	594 (91.5%)	576 (91.7%)	18 (85.7%)
No	52 (8.0%)	49 (7.8%)	3 (14.3%)
Unknown	3 (0.5%)	3 (0.5%)	0
Certain of pregnancy expulsion?†	Entire cohort (n=663)	Gestation under 10 weeks (n=642)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
Yes	479 (72.2%)	462 (72.0%)	17 (81.0%)
No or Not Sure	161 (24.4%)	157 (24.5%)	4 (19.0%)
Unknown	23 (3.5%)	23 (3.6%)	0
Induction–expulsion interval (hours)‡	Entire cohort certain pregnancy expelled (n=479)	Gestation under 10 weeks certain pregnancy expelled (n=462)	Gestation 10 weeks – 11 weeks and 6 days certain pregnancy expelled (n=17)
Mean	4.3	4.3	4.4
Median	3	3	2.5
SD	4.3	4.3	4.2
Induction–expulsion interval (hours)§	Cohort used medications correctly and certain pregnancy expelled (n=442)	Cohort used medications incorrectly and certain pregnancy expelled (n=37)	
Mean	4.2	5	
Median	3	3.25	
SD	4.2	5.3	

*Correct use of mifepristone and misoprostol is defined as interval between mifepristone and misoprostol administration between 24 and 72 hours.

†Participants reporting they were certain that they had passed pregnancy tissue following misoprostol administration. 'No' means that they did not pass pregnancy, 'Not sure' means they do not know if they passed the pregnancy or not, 'Unknown' indicates the number of people not responding to the question.

‡ Misoprostol administration (induction) until expulsion of pregnancy intervals in hours, when participant was certain pregnancy expelled.

§ Induction–expulsion interval when certain pregnancy expelled, grouped by correct or incorrect use of abortion medications.

than one of these treatments). The mean (6.8) and median (7) pain scores in this group were similar to the cohort overall. Further detail is provided in [table 3](#).

Side effects and bleeding profile

Regarding side effects, 510/663 (77%) women experienced either nausea, vomiting, diarrhoea or headache; 21/663 (3%) reported all four side effects during treatment. Regarding bleeding profile, 510/663 (77%) reported bleeding that was more than their usual menstrual period. Of those patients reporting the

same or less bleeding than a period, only four required further abortion medication to complete the abortion.

Further detail is provided in [table 4](#).

Reflecting on timing and preparedness for treatment

Patients were asked on day 14 about the duration of the consultation and how prepared they felt for treatment; 562/663 (85%) felt the consultation duration was 'just right', 15/663 (2%) thought it was too long and 2/663 (0.3%) thought it was too short. Likewise, 554/663 (84%) felt adequately prepared for abortion

Table 2 Antibiotic and antiemetic provision and reported use

Prophylactic antibiotics given*	Entire cohort (n=663)	Gestation under 10 weeks (n=642)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
Yes	599 (90.3%)	581 (90.5%)	18 (85.7%)
No	44 (6.6%)	42 (6.5%)	2 (9.5%)
Unknown	20 (3.0%)	19 (3.0%)	1 (4.8%)
Prophylactic antibiotics used*	Given antibiotics (n=599)	Gestation under 10 weeks given antibiotics (n=581)	Gestation 10 weeks – 11 weeks and 6 days given antibiotics (n=18)
Took all	383 (63.9%)	369 (63.5%)	14 (77.8%)
Took some	53 (8.8%)	52 (9.0%)	1 (5.6%)
Took none	97 (16.2%)	95 (16.4%)	2 (11.1%)
Unknown	66 (11.0%)	65 (11.2%)	1 (5.6%)
Antiemetic use†	Entire cohort (n=663)	Gestation under 10 weeks (n=642)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
Yes	611 (92.2%)	591 (92.1%)	20 (95.2%)
No	32 (4.8%)	31 (4.8%)	1 (4.8%)
Unknown	20 (3.0%)	20 (3.1%)	0
Number of doses of antiemetic used, of those answering 'Yes' to antiemetic use†	Cohort reporting number of doses (n=608, 3 did not report number of doses used)	Gestation under 10 weeks reporting number of doses (n=588)	Gestation 10 weeks – 11 weeks and 6 days reporting number of doses (n=20)
Median	2	2	2
Mean	2.1	2.1	2.7
SD	0.9	0.9	1.1

*Antibiotics given were a 7-day course of oral doxycycline 100 mg twice daily. 'Took all' means completed the whole course, 'Took some' means they used at least one dose but did not complete the full course, 'Took none' means they did not use any doses'.

†All patients were provided with five doses of cyclizine 50 mg and directed to use one tablet before mifepristone, one before misoprostol, and the remainder as required.

treatment, 12/663 (2%) felt neither prepared nor unprepared and 11/663 (2%) felt unprepared.

DISCUSSION

Main findings

This study contributes novel data on the patient use of medications when receiving medical abortion at home following a telemedicine consultation.

The majority of patients in our cohort were able to self-administer their medications correctly, required two doses of misoprostol or fewer, and had an induction–expulsion interval comparable to that reported in the literature—that is, the majority reporting expulsion between 3 and 6 hours after misoprostol administration.^{11 12} Of those with pregnancies less than 10

weeks' gestation, 8/10 patients required only a single 800 µg dose of misoprostol.

Provision of antibiotics was high and of those receiving antibiotics, approximately two-thirds completed the full course. Antiemetic use was similarly high. Clinical advice at the time was to use a dose prophylactically before mifepristone and again before misoprostol. This may be why the mean number of doses used was 2. Despite this, nearly one-third of women experienced vomiting and many experienced nausea and diarrhoea. Gastrointestinal side effects and their optimal management require further investigation.

Although use of analgesia was widespread, including weak opioids, pain scores were still relatively high, suggesting an unmet need for optimal pain management. Headache was experienced by one in six patients and could be related to analgesia or abortion medications.

By day 4 the majority reported experiencing more bleeding than a period, which is consistent with literature, although we were not able to quantify this exactly in this study.¹³

Preparedness for the procedure was high when assessed at 14 days and the proportion rating themselves as prepared is similar to the proportion when asked at 4 days post-treatment.⁶ The duration of the teleconsultations was acceptable to the majority of patients. These findings are reflected in qualitative data from this cohort, indicating that women found the process straightforward and were well prepared by the telephone consultation.¹⁴

This cohort included a small number of patients receiving abortion care at home between 10 weeks and 11 weeks 6 days' gestation; however, their experiences were comparable to the cohort overall for use of abortion medications and other outcomes of interest.

What this study contributes

The Scottish government recently conducted a public consultation on the continued use of telemedicine. Some respondents were concerned that evidence was lacking on women's adherence to medical abortion treatment in the context of telemedicine and self-administration of medications. This study robustly examines those concerns and demonstrates high rates of adherence to medical abortion regimens.¹⁵

The overall package of information—audiovisual and written information online, verbal information delivered by telephone consultation and written information leaflets included in treatment packs—was sufficient to adequately prepare women for medical abortion at home and empower them to correctly self-administer abortion medications.

We have identified that pain and other side effects are imperfectly managed with the current analgesia and antiemetic regimens used. Pain scores during medical abortion in other studies are inconsistently reported, but those that do report them are at similar levels to

Table 3 Pain experience and analgesia use reported at day 4 following treatment

Experienced pain	Entire cohort (n=663)	Gestation under 10 weeks (n=642)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
Yes	624 (94.1%)	603 (93.9%)	21 (100%)
No	22 (3.3%)	22 (3.4%)	0
Unknown	17 (2.6%)	17 (2.6%)	0
Pain score of those answering 'Yes' to pain experienced*	Cohort reporting pain score (n=610, 14 did not report pain score)	Gestation under 10 weeks (n=589)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
Median	7	7	7
Mean	6.7	6.7	7.0
SD	2.2	2.2	2.0
Pain experienced compared with expectation, of those providing a pain score	Cohort responding (n=587, 23 did not respond)	Gestation under 10 weeks (n=568, 21 did not respond)	Gestation 10 weeks – 11 weeks and 6 days (n=19, 2 did not respond)
Better than expected	227 (38.7%)	217 (38.2%)	10 (52.6%)
Same as expected	148 (25.2%)	144 (25.4%)	4 (21.1%)
Worse than expected	212 (36.1%)	207 (36.4%)	5 (26.3%)
Used analgesia	Entire cohort (n=663)	Gestation under 10 weeks (n=642)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
Yes†	616 (92.9%)	596 (92.8%)	20 (95.2%)
No	30 (4.5%)	29 (4.5%)	1 (4.8%)
Unknown	17 (2.6%)	17 (2.6%)	0
Paracetamol			
Yes	610 (92.0%)	593 (92.4%)	17 (81.0%)
No	36 (5.4%)	32 (5.0%)	4 (19.0%)
Ibuprofen			
Yes	403 (60.9%)	390 (60.7%)	13 (61.9%)
No	243 (36.7%)	235 (36.6%)	8 (38.1%)
Dihydrocodeine‡			
Yes	427 (64.4%)	413 (64.3%)	14 (66.7%)
No	219 (33.0%)	212 (33.0%)	7 (33.3%)

*Pain score on an 11-point Likert scale from 0 to 10, and comparison of pain experienced with expectations of pain.

†Yes includes use of paracetamol, ibuprofen, dihydrocodeine, or any other pain relief treatment.

‡Patients were given five doses of dihydrocodeine 30 mg in their treatment packs and directed to purchase their own paracetamol and ibuprofen.

those in our study, between 5 to 7 out of 10 on visual analogue or numerical Likert scales.¹⁶

Although there have been various trials of analgesic medications^{17–23} during medical abortion, the optimal dosing and frequency remain unclear. Further research is required to identify an effective and acceptable regimen that is easy for patients to self-administer at home and with minimal side effects. We were only able to offer a single antiemetic in the form of oral cyclizine during this study. There may be other antiemetics that are better suited to managing nausea during medical abortion, such as metoclopramide,²⁴ but further study is required.

Strengths and limitations

More than 98% of women in the study cohort completed at least one of the questionnaires. This group represents almost all of the women receiving medical abortion at home during the study period. This may have been influenced by the COVID-19

restrictions in place at the time of the study, meaning that patients were easily contactable by telephone.

Pain score, bleeding and side effect data were collected within a short time frame to minimise errors of recall, but they relied on women's recall rather than on diaries or prospective recording.

This is a single-centre study in a high-resource setting with a well-developed community medical abortion service. The findings of this study may be reproducible more broadly—any centre that can offer clearly written patient information combined with verbal and online advice and telephone support before, during and after an abortion could adopt this approach.

This was an observational study which has inherent limitations in comparison with randomised controlled trials and comparison is only available in relation to previously published data prior to COVID-19 and requirements for some or all medications being administered in a clinical facility. Nevertheless, this study

Table 4 Side effects and bleeding profile reported at day 4 following treatment

Side effects		Entire cohort (n=663)	Gestation under 10 weeks (n=642)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
Nausea	Yes	415 (62.6%)	402 (62.6%)	13 (61.9%)
	No	231 (34.8%)	223 (34.7%)	8 (38.1%)
	Unknown	17 (2.6%)	17 (2.6%)	0
Vomiting	Yes	223 (33.6%)	213 (33.2%)	10 (47.6%)
	No	423 (63.8%)	412 (64.2%)	11 (52.4%)
	Unknown	17 (2.6%)	17 (2.6%)	0
Diarrhoea	Yes	199 (30.0%)	192 (29.9%)	7 (33.3%)
	No	446 (67.3%)	432 (67.3%)	14 (66.7%)
	Unknown	18 (2.7%)	18 (2.8%)	0
Headache	Yes	101 (15.2%)	97 (15.1%)	4 (19.0%)
	No	544 (83.6%)	527 (82.1%)	17 (81.0%)
	Unknown	18 (2.7%)	18 (2.8%)	0
Bleeding profile		Entire cohort reporting bleeding (n=637, no bleeding=6, unknown=20)	Gestation under 10 weeks (n=616)	Gestation 10 weeks – 11 weeks and 6 days (n=21)
Much more than a period		305 (47.9%)	292 (47.4%)	13 (61.9%)
A bit more than a period		205 (32.2%)	197 (32.0%)	8 (38.1%)
Same as a period		81 (12.7%)	81 (13.1%)	0
A bit less than a period		23 (3.6%)	23 (3.7%)	0
Much less than a period		23 (3.6%)	23 (3.7%)	0

provides the first detailed descriptive information on women's use of medication for medical abortion at home when delivered via telemedicine.

CONCLUSION

Telemedicine can effectively deliver information to patients to prepare them to safely and correctly use abortion medications as part of medical abortion at home with remote support. Further research is required to optimise the management of pain and other side effects during medical abortion.

Twitter John Joseph Reynolds-Wright @doctorjjrw

Contributors JJR-W and SC designed the study, analysed the data and drafted the manuscript. AJ and KM contributed to questionnaire design and collected the data. EE undertook statistical analysis. All authors reviewed the final manuscript prior to submission. JJR-W is the guarantor of this paper.

Funding The study was conducted by staff at the MRC Centre for Reproductive Health, which is supported by grant MR/N022556/1. The Edinburgh Family Planning Trust provided funding to employ clinical research nurses.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. The original data are not available in a public repository. Please contact the corresponding author for any data requests to be considered.

Open access This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

ORCID iDs

John Joseph Reynolds-Wright <http://orcid.org/0000-0001-6597-1666>

Sharon Cameron <http://orcid.org/0000-0002-1168-2276>

REFERENCES

- Royal College of obstetricians and gynaecologists. The care of women requesting induced abortion: evidence-based clinical guideline number 7. London, 2011. Available: https://www.rcog.org.uk/globalassets/documents/guidelines/abortion-guideline_web_1.pdf [Accessed 4 Jun 2020].
- Low ST, Chen ZE, Cameron S. Women's experiences of self-referral to an abortion service: qualitative study. *BMJ Sex Reprod Health* 2021;47:37–42.
- Chief Medical Officer for Scotland. Abortion – Covid-19 – approval for mifepristone to be taken at home and other contingency measures, 2020. Available: <https://www.sehd.scot.nhs.uk/cmo/CMO%282020%2909.pdf> [Accessed 20 Oct 2021].
- Public Health Scotland. Termination of pregnancy statistics year ending 2019, 2020. Available: <https://beta.isdscotland.org/>

- find-publications-and-data/population-health/sexual-health/termination-of-pregnancy-statistics/ [Accessed 20 Oct 2021].
- 5 Royal College of Obstetricians and Gynaecologists. Coronavirus (COVID-19) infection and abortion care: information for healthcare professionals. London, 2020. Available: <https://bsacp.org.uk/wp-content/uploads/2020/05/2020-04-09-coronavirus-covid-19-infection-and-abortion-care.pdf> [Accessed 4 Jun 2020].
 - 6 Reynolds-Wright JJ, Johnstone A, McCabe K, *et al*. Telemedicine medical abortion at home under 12 weeks' gestation: a prospective observational cohort study during the COVID-19 pandemic. *BMJ Sex Reprod Health* 2021;47:-246–51. doi:10.1136/bmjshr-2020-200976
 - 7 Reynolds-Wright JJ, Bellevue F, Daberius A, *et al*. Information on early medical abortion for women using an audiovisual animation vs face-to-face consultation: a consortium randomized and quasi-randomized trial. *Acta Obstet Gynecol Scand* 2020;99:1611–7.
 - 8 Aiken A, Lohr PA, Lord J, *et al*. Effectiveness, safety and acceptability of no-test medical abortion (termination of pregnancy) provided via telemedicine: a national cohort study. *BJOG* 2021;128:1464–74.
 - 9 Kerestes C, Murayama S, Tyson J, *et al*. Provision of medication abortion in Hawai'i during COVID-19: practical experience with multiple care delivery models. *Contraception* 2021;104:49–53.
 - 10 Chong E, Shochet T, Raymond E, *et al*. Expansion of a direct-to-patient telemedicine abortion service in the United States and experience during the COVID-19 pandemic. *Contraception* 2021;104:43–8.
 - 11 Kapp N, Eckersberger E, Lavelanet A, *et al*. Medical abortion in the late first trimester: a systematic review. *Contraception* 2019;99:77–86.
 - 12 Kulier R, Kapp N, Gülmezoglu AM, *et al*. Medical methods for first trimester abortion. *Cochrane Database Syst Rev* 2011;CD002855.
 - 13 Baird DT, Rodger M, Cameron IT, *et al*. Prostaglandins and antigestagens for the interruption of early pregnancy. *J Reprod Fertil Suppl* 1988;36:173–9.
 - 14 Boydell N, Reynolds-Wright JJ, Cameron ST, *et al*. Women's experiences of a telemedicine abortion service (up to 12 weeks) implemented during the coronavirus (COVID-19) pandemic: a qualitative evaluation. *BJOG* 2021;128:1752–61.
 - 15 Scottish Government - Minister for Public Health. Women's Health and Sport. Future arrangements for early medical abortion at home consultation analysis, 2021. Available: <https://www.gov.scot/publications/future-arrangements-early-medical-abortion-home-consultation-analysis/> [Accessed 20 Oct 2021].
 - 16 Fiala C, Cameron S, Bombas T, *et al*. Pain during medical abortion, the impact of the regimen: a neglected issue? A review. *Eur J Contracept Reprod Health Care* 2014;19:404–19.
 - 17 Avraham S, Gat I, Duvdevani N-R, *et al*. Pre-emptive effect of ibuprofen versus placebo on pain relief and success rates of medical abortion: a double-blind, randomized, controlled study. *Fertil Steril* 2012;97:612–5.
 - 18 Livshits A, Machtinger R, David LB, *et al*. Ibuprofen and paracetamol for pain relief during medical abortion: a double-blind randomized controlled study. *Fertil Steril* 2009;91:1877–80.
 - 19 Raymond EG, Weaver MA, Louie KS, *et al*. Prophylactic compared with therapeutic ibuprofen analgesia in first-trimester medical abortion: a randomized controlled trial. *Obstet Gynecol* 2013;122:558–64.
 - 20 Friedlander EB, Soon R, Salcedo J. Prophylactic pregabalin to decrease pain during medication abortion: a randomized controlled trial. *Obstet Gynecol* 2018;132:612–8.
 - 21 Ojha K, Gillott DJ, Wood P, *et al*. Clinical outcomes from a prospective study evaluating the role of ambulation during medical termination of pregnancy. *Contraception* 2012;85:398–401.
 - 22 Reynolds-Wright JJ, Woldetsadik MA, Morroni C. Pain management for medical abortion before 14 weeks' gestation. *Cochrane Database Syst Rev* 2020:1465–858. doi:10.1002/14651858.CD013525
 - 23 Colwill AC, Bayer LL, Bednarek P, *et al*. Opioid analgesia for medical abortion: a randomized controlled trial. *Obstet Gynecol* 2019;134:1163–70. doi:10.1097/AOG.00000000000003576
 - 24 Dragoman MV, Grossman D, Nguyen MH, *et al*. Two prophylactic pain management regimens for medical abortion ≤63 days' gestation with mifepristone and misoprostol: a multicenter, randomized, placebo-controlled trial. *Contraception* 2021;103:163–70.

Supplementary File 1: Questionnaire

DAY FOUR QUESTIONS

Question 1:

Did you take the first tablet (mifepristone, 1 tablet that you swallow)?

Circle: Yes / No

If yes, do you remember when:

Date: _____ Time: _____

Or circle: Don't remember

Question 2:

Do you remember the date and time you took the first dose of your misoprostol tablets (4 tablets under the tongue or inside the vagina)?

Circle: Yes / No

If yes:

Date: _____ Time: _____

Did you take the tablets (circle): under the tongue / inside the vagina /
between your cheek and gum ?

Question 3:

Did you use any additional doses of misoprostol? (2 more tablets under the tongue or inside the vagina)

Circle: Yes / No

If yes:

How many further doses (i.e. pairs of tablets) did you take? _____

Question 4:

Do you remember the date and time that you passed the pregnancy?

Circle: Yes / No / Unsure

If yes:

Date: _____ Time: _____

Supplementary File 1: Questionnaire

DAY FOUR QUESTIONS

Question 1:

Did you take the first tablet (mifepristone, 1 tablet that you swallow)?

Circle: Yes / No

If yes, do you remember when:

Date: _____ Time: _____

Or circle: Don't remember

Question 2:

Do you remember the date and time you took the first dose of your misoprostol tablets (4 tablets under the tongue or inside the vagina)?

Circle: Yes / No

If yes:

Date: _____ Time: _____

Did you take the tablets (circle): under the tongue / inside the vagina /
between your cheek and gum ?

Question 3:

Did you use any additional doses of misoprostol? (2 more tablets under the tongue or inside the vagina)

Circle: Yes / No

If yes:

How many further doses (i.e. pairs of tablets) did you take? _____

Question 4:

Do you remember the date and time that you passed the pregnancy?

Circle: Yes / No / Unsure

If yes:

Date: _____ Time: _____

Supplementary File 1: Questionnaire

DAY FOUR QUESTIONS

Question 1:

Did you take the first tablet (mifepristone, 1 tablet that you swallow)?

Circle: Yes / No

If yes, do you remember when:

Date: _____ Time: _____

Or circle: Don't remember

Question 2:

Do you remember the date and time you took the first dose of your misoprostol tablets (4 tablets under the tongue or inside the vagina)?

Circle: Yes / No

If yes:

Date: _____ Time: _____

Did you take the tablets (circle): under the tongue / inside the vagina /
between your cheek and gum ?

Question 3:

Did you use any additional doses of misoprostol? (2 more tablets under the tongue or inside the vagina)

Circle: Yes / No

If yes:

How many further doses (i.e. pairs of tablets) did you take? _____

Question 4:

Do you remember the date and time that you passed the pregnancy?

Circle: Yes / No / Unsure

If yes:

Date: _____ Time: _____

Question 5:

Did you experience pain during the procedure?

Circle: Yes / No

If yes:

- What was the worst pain you experienced on a scale of 0-10 (0 being no pain, 10 being worst pain imaginable): _____
- Was this pain *worse than expected / as bad as expected / better than expected?* (please circle)

Question 6:

Did you use any pain killers during the procedure?

Circle: Yes / No

If yes:

What did you use:

Drug name	Used? (Circle)	How many doses (write strength and number e.g. 400mg x 3)
Paracetamol	Yes / No	
Ibuprofen	Yes / No	
Dihydrocodeine	Yes / No	
Other (ask about own supply of other meds and also recreational drugs e.g. cannabis):	Yes / No	

Question 7:

Did you experience any of the following (circle):

- Nausea: Yes / No
- Vomiting: Yes / No
- Diarrhoea: Yes / No
- Headache: Yes / No

Question 8:

Have you experienced any bleeding?

Circle: Yes / No

If yes:

How did this compare to a typical period for you:

- Much more bleeding than a period
- A bit more bleeding than a period
- The same amount of bleeding as a period
- A bit less bleeding than a period
- Much less bleeding than a period

Question 9:

Did you use any of the anti-sickness pills provided (cyclizine)?

Circle: Yes / No

If yes:

How many tablets did you use in total? _____

DAY FOURTEEN QUESTIONS

Question 10: Were you given antibiotics? Yes / No
If yes, did you use them?: Took full course / Took some / Took none

Question 11:

Looking back, what did you think of duration of the consultation?

- Much too long
- A bit longer than I wanted
- Just right
- A bit shorter than I wanted
- Much too short

Question 12:

Looking back, how well prepared were you for your abortion?

- Very prepared
- Somewhat prepared
- Neutral
- Somewhat unprepared
- Very unprepared