




OPEN ACCESS

Rhesus isoimmunisation in unsensitised RhD-negative individuals seeking abortion at less than 12 weeks' gestation: a systematic review

Michelle C Chan ,¹ Roopan Kaur Gill,² Caron Rahn Kim³

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmj-srh-2021-201225>).

¹Department of Obstetrics and Gynecology, The University of British Columbia Faculty of Medicine, Vancouver, British Columbia, Canada

²Department of Obstetrics and Gynecology, University of Toronto, Toronto, Ontario, Canada

³Department of Reproductive Health and Research, World Health Organization, Geneva, Switzerland

Correspondence to

Dr Michelle C Chan, Obstetrics and Gynecology, The University of British Columbia Faculty of Medicine, Vancouver, Canada; mcchan@ualberta.ca

Received 29 May 2021

Accepted 27 October 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Chan MC, Gill RK, Kim CR. *BMJ Sex Reprod Health* Published Online First: [please include Day Month Year]. doi:10.1136/bmj-srh-2021-201225

ABSTRACT

Aim The aim of this review was to systematically review the outcome of routine anti-D administration among unsensitised rhesus (RhD)-negative individuals who have an abortion. This review is registered with Prospero.

Methods A search for all published and ongoing studies, without restrictions on language or publication status, was performed using the following databases from their inception: EBM Reviews Ovid - Cochrane Central Register of Controlled Trials, MEDLINE Ovid (Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily), Embase.com, Popline and Google Scholar. Study types included: randomised controlled trials, controlled trials, cohort and case-control studies from 1971 onwards. The population included women who undergo an abortion (induced, incomplete, spontaneous or septic abortion), medical or surgical <12 weeks, and isoimmunisation in a subsequent pregnancy. The primary outcomes were: (1) development of a positive Kleihauer-Betke test and (2) development of Rh alloimmunisation in a subsequent pregnancy.

Results A total of 2652 studies were screened with 105 accessed for full-text review. Two studies have been included with high bias appreciated. Both studies found few women to be sensitised in forming antibodies after an abortion. The limited studies available and heterogeneity prevent the conduction of a meta-analysis.

Conclusions Rh immunoglobulin has well-documented safety. However, it is not without risks and costs, is a possible barrier to delivering efficient services, and may have limited availability in some countries. The evidence base and quality of studies are currently limited. There is unclear benefit from the recommendation for

Key messages

- Rhesus (Rh) immunoglobulin has well-documented safety but is not without risks, costs or acquisition complexity and may cause unnecessary barriers and delay for patients accessing abortion care.
- There is limited evidence on Rh sensitisation for first-trimester abortions at <12 weeks' gestation (including threatened, spontaneous, surgical or medical abortion).
- Clinical practice guidelines currently based on expert opinions vary according to geographical region.

Rh testing and immunoglobulin administration in early pregnancy. More research is needed as clinical practice guidelines are varied, based on expert opinions and moving away from testing and administration at time of abortion.

Implications There is limited evidence surrounding medical benefit of Rh testing and immunoglobulin administration in early pregnancy. Further research is needed to define alloimmunisation and immunoglobulin benefit to update standards of care. Additionally, other factors should be considered in forming clinical policies and guidelines such as costs, feasibility and impact on access to care for patients.

BACKGROUND

The administration of rhesus (Rh) immunoglobulin was first introduced in 1968 and significantly reduced immunisation to D-antigen.¹ Evidence is sparse for such intervention after abortion in early pregnancy.¹ Abortion for the purposes of this

review included induced (medical or surgical), incomplete, missed, spontaneous or septic abortion at less than 12 weeks' gestation.

There is a movement to forgo Rh testing and anti-D immunoglobulin administration in people presenting for early abortion, miscarriage or ectopic pregnancy.^{2,3} For example, the 2019 National Institute for Health and Care Excellence (NICE) guidelines on abortion recommend not offering anti-D prophylaxis to people undergoing medical abortion up to and including 10 weeks' gestation.⁴ Rationales to forgo Rh testing and anti-D immunoglobulin administration include lack of evidence that anti-D immunoglobulin prevents Rh alloimmunisation in lower gestational ages and immunoglobulin administration is not without risk as it is a blood product with added costs and care complexity.^{2,3} For example, added costs and complexity include testing patient blood type and cold storage required for anti-D immunoglobulin.

A systematic review by Karanth *et al* (2013) examined anti-D administration after spontaneous miscarriage for preventing Rh alloimmunisation with the Cochrane Pregnancy and Childbirth Group.⁵ Karanth *et al* concluded there is insufficient evidence available to evaluate the current practices of anti-D administration in unsensitised RhD-negative mothers after spontaneous miscarriage.

To date, there has been one systematic review examining subsequent isoimmunisation after abortion at less than 13+6 weeks' gestation in 2019.⁴ In preparation for updating the World Health Organization (WHO) 2012 Safe Abortion guidelines, the objective of the review was to systematically review the effect of routine anti-D administration among unsensitised RhD-negative individuals who have an abortion.

METHODS

This review follows Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidance.⁶ A protocol for this systematic review is included as online supplemental appendix 1–2. This review is registered with Prospero (CRD42020149073). The population included women who undergo an abortion: induced (medical or surgical), incomplete, missed, spontaneous or septic abortion at less than 12 weeks' gestation. The intervention examined was routine anti-D administration compared with no anti-D administration.

A search was performed for all published and ongoing studies, without language restrictions, using the following databases from their inception to 13 May 2021: EBM Reviews Ovid - Cochrane Central Register of Controlled Trials, MEDLINE Ovid (Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily), Embase.com, Popline and Google Scholar (online supplement 1–2). [NB. Popline and Google Scholar were only searched to 23 July 2019.]

We included randomised controlled trials, controlled trials, cohort studies and case-control studies. The following study types were excluded: case studies, review articles, editorials, letters, advisories, non-comparative studies, unpublished manuscripts, conference abstracts, diagnostic studies, animal studies, cost-benefit analyses, and studies with basic science outcomes. Non-English publications and articles published prior to 1971 were excluded given WHO only started to recommend that Rh testing and treatment with immunoglobulin be made part of the standard protocol of medical care for pregnant women from 1971 onwards.

Primary outcomes included: (a) development of a positive Kleihauer–Betke test (a test that detects fetal cells in the maternal blood) and (b) development of RhD alloimmunisation in a subsequent pregnancy. Secondary outcomes included: (a) detection of atypical blood group antibodies by positive indirect Coombs test after 6 months of exposure (non-prespecified outcome), (b) need for increased surveillance for suspected fetal blood sampling and fetal transfusions in subsequent pregnancies, (c) neonatal morbidity such as neonatal anaemia, jaundice, bilirubin encephalopathy, erythroblastosis, prematurity, hypoglycaemia (low blood sugar) in subsequent pregnancies and (d) maternal adverse events of anti-D administration including anaphylactic reaction. Currently, flow cytometry has been found to be a more accurate estimate of sensitisation.^{1,7} However, this review utilised the Kleihauer–Betke test as an inclusion criterion because it is more available and commonly used in clinical settings.

Three authors (MCC, CRK and RKG) independently screened all the titles, abstracts and full texts identified from the initial search to determine eligibility for inclusion. The intervention examined was routine anti-D administration compared with no anti-D administration. Studies that did not explicitly include routine anti-D administration as the intervention were excluded. Wrong comparators would include the use of different testing for the primary outcome of developing RhD alloimmunisation confirmed with a positive Kleihauer–Betke test. Conflicts were resolved through discussion and consensus first between MCC and RKG and, if further discussion was needed, with the third reviewer CRK. We used a standard template for data abstraction (table 1).

Data synthesis

A narrative synthesis was performed; the findings are summarised based on study design, population, intervention, comparison, results, strengths and weaknesses based on information reported by the study authors and subjective assessment of the review authors (table 1). The limited number and heterogeneity of the included studies prevent the conduction of a meta-analysis. Risk of bias was assessed by MCC and CRK with quality of evidence assessment frameworks. Cochrane was used

Table 1 Rhesus (Rh) isoimmunisation in un sensitised Rh-negative individuals seeking abortion at <12 weeks

Author, year, funding	Study design, location, year(s) of data collection, follow-up	Population	Intervention	Comparison	Results	Strengths	Weaknesses	Risk of bias
Gavin (1972) Funding: Ortho Research Foundation and the Kaiser Foundation Hospital	Design: prospective double-blind Location: Kaiser Foundation Hospital in Los Angeles, California - Bellflower and Panorama City Duration of data collection: 1 November 1969–15 August 1970 Follow-up: 4 months with indirect Coombs and paternal genotyping (this was only done in 50% of couples)	491 women underwent therapeutic abortions 180 treated for spontaneous abortion 57 were found to be Rh-negative and indirect Coombs at the time of abortion n=57 ▶ Three refused participation ▶ Nine lost to follow-up ▶ One had a husband found to be Rh-negative Remaining 44 - Combined with 13 patients identified the same way at Bellflower and Panorama City * <i>Ec topic was included as a 'method of abortion' ... same with hysterotomy</i>	Within 72 hours of abortion given RhoGAM - 21 patients 8–16 weeks and a 24-week preterm delivery received RhoGAM	Within 72 hours given placebo - 36 patients received placebo 4–20 weeks	2/36 sensitisations from placebo group Patient 1 – G1 with TA at 81 days under GA, 1:16 positive indirect Coombs 7 and 9 months after, no history of transfusions or previous pregnancies Patient 2 – G1 TA at 76 days under GA, 1:2 positive indirect Coombs test 3 months and 19 days after abortion, no history of termination or previous pregnancies No patients receiving RhoGAM sensitised	1. Described as “double-blinded RCT” (poor) 2. Objective measurement of outcome, titres reported 3. No defined sample size 4. No power calculations 5. There is no ‘Table 1’ on the demographics of the patients 6. No statistics performed (eg, intention to treat analysis) * Giving someone no RhoGAM at 4 weeks is not helpful, as we know the yolk sac hasn’t even been developed yet	High	
Goldman and Eckerling (1977) Funding: source not noted	Design: prospective? Location: not noted, but Tel Aviv Duration of data collection: not noted Follow-up: 9 months or more ▶ Antibody titres at 6 weeks, 3 months and 6 months after abortion ▶ Patients with antibodies at 3 months → ask to continue serological follow-up for 24 months after abortion	170 postabortion patients >6 months in Rh-negative mothers (no Rh antibodies in her blood and a negative Coombs test) - When possible, husband’s ABO blood group determined to ensure compatibility with wife’s	48 patients in treated group ▶ Prior to Rh serological test to confirm ABO group and Rh type, indirect Coombs and antibody titres were sensitised	5/122 patients in control group sensitised	Incidence of Rh immunisation in a period of 6 months or more postabortion in Rh-negative mothers	1. Table documenting “admin” – who got anti-D and who didn’t 2. Follow-up of Rh-negative patients	1. Patients were not randomised; placed in the control group when anti-D was not available 2. No allocation concealment 3. No follow-up rate reported 4. No sample size calculation 5. No power calculation	High

G1, primiparous; GA, general anaesthesia; TA, therapeutic abortion.




Figure 1 PRISMA flow diagram.

for randomised trials and ROBINS-I tool for observational studies.^{8 9}

RESULTS

The search strategy identified 2652 publications and three duplicate records were removed, leaving 2649 publications for abstract screening (figure 1). From abstract screening, 105 publications underwent full-text review. The two included studies were cohort studies taking place in the United States and Israel.^{10 11} Both studies included women with induced abortion in addition to other types of abortion.

A 1972 study funded by Ortho Research Foundation and the Kaiser Foundation Hospital performed a prospective double-blind study collecting data on 491 women undergoing therapeutic abortions between 1 November 1969 and 15 August 1970 in California.¹¹ The primary aim of this study was to investigate Rh immunisation incidence in a postabortion period of 6 months or more in RhD-negative people. From the 491 therapeutic abortion patients, and 180 spontaneous abortion patients, 57 were RhD-negative based on indirect Coombs at the time of abortion management. The resulting sample size was 44 because three refused participation, nine were lost to follow-up and one husband was found to also be RhD-negative (paternal genotyping was only performed in 50% of couples). Within 72 hours of abortion the patient received RhoGAM or a placebo. The study found two of 36 sensitisations in the placebo group with indirect Coombs positive after abortion. All patients receiving RhoGAM were not found to be sensitised. Despite a 'double-blind' approach there was no description of allocation concealment, therefore the authors of this review considered this an impact on study quality. However, the objective measurement of outcome (titres) is a strength of the study. Other

factors contributing to the limitations of the study and its quality include: no description of allocation, no follow-up rate reported for patients, no defined sample size, no power calculations for sample size, no intention to treat analysis and no report of a 'Table 1' for the demographics of the population studied.

The second included study was published in 1972 and followed 170 RhD-negative postabortion patients for 6 months or more with antibody titre testing.¹⁰ The primary aim of this study was to identify Rh sensitisation after an abortion in those who did and did not receive RhoGAM. There was no clear description of data collection duration. For example, patients were followed for different times depending on their antibody status. All postabortion participants were observed for 6 months with antibody titres performed at 6 weeks, 3 months and 6 months. In sensitised participants with antibodies detected at 3 months, repeat follow-up serology was performed 24 months after abortion. The study followed patients identified as RhD-negative mothers without antibodies in their blood and a negative Coombs test. When possible, a husband's ABO blood group was also determined. There were 48 in the intervention group, receiving an injection of RhoGAM, and 122 in the control group. The main outcome was incidence of rhesus sensitisation in a period of 6 months or more postabortion in RhD-negative mothers. The study identified five mothers who became sensitised in the control group. There were no sensitisations found in the intervention group. The authors of this review assessed the limitations of the study and its quality which included: patients not randomised (patients were placed in the control group if RhoGAM was not available), lack of allocation concealment, lack of reported follow-up rate, no sample size or power calculation. The study did not identify sensitised patients within the intervention group.

DISCUSSION

Our review identifies limited evidence on the effectiveness of anti-D administration in women who are RhD-negative and undergo an early abortion. This is consistent with past literature reviews which also provided limited results.^{1 12 13} Our findings are consistent that there are few studies showing maternal sensitisation or fetal haemolytic disease due to foeto-maternal haemorrhage from first-trimester abortion.^{1 12 13} Furthermore, evidence surrounding sensitisation secondary to ectopic pregnancy is very limited.¹³

There is no evidence of clear benefit in early pregnancy or the gestational age at which sensitisation can occur among aspiration or medication abortion.^{2 3} Current clinical practice guidelines are generally based on expert opinions.^{4 14} The prevalence of RhD-negative varies among populations.¹⁵ A review comparing Canadian and Netherlands policies surrounding Rh immunoglobulin administration in first-trimester pregnancy

found no differences in sensitisation.² The populations between Canada and the Netherlands were found to be similar, however there is a difference in practice: Canada recommending Rh immunoglobulin administration and the Netherlands, at the time of this study, offering Rh immunoglobulin to RhD-negative women having spontaneous abortions over 10 weeks 0 days' gestation and induced abortions over 7 weeks 0 days.² Furthermore, in the Netherlands, where no anti-D immunoglobulin is provided routinely in early abortion, showed lower sensitisation rates than Canada.² There does not appear to be a clear benefit in Rh immunoglobulin administration in the first trimester.

A recent prospective study examined 42 pregnant people at 5–12 weeks' gestational age for the minimum fetal red blood cell concentration required to cause maternal Rh sensitisation.¹⁶ The study found fetal blood cell exposure in the first trimester to be well below a calculated threshold for Rh sensitisation.¹⁶ Of note, the study used flow cytometry which is a more accurate marker of feto-maternal haemorrhage.¹⁶ However, the authors identified that larger studies are needed to confirm their study findings.¹⁶ These findings also support the suggestion there is no advantage in Rh immunoglobulin administration in the first trimester.

This review is robust from the systematic search of the literature. However, the studies included for this review were of poor quality.^{10 11} Included studies are quite dated whereby flow cytometry is now considered the most accurate method of testing.¹ Given this, it is important to consider changes in clinical practice and standards contextually. For example, previous surgical techniques likely included more sharp curettage relative to today's standards. Additionally, the included studies had small sample sizes, with approximately 50 women in each study.

Given minimal quality evidence and no clear benefit of RhD immunoglobulin administration after early abortion, and some disadvantages of doing so, further research is needed. Some disadvantages to the routine administration of RhD immunoglobulin include: cost, cold storage, and barriers to access in the context of increasing self-managed abortions. There is also a dissonance surrounding clinically used versus laboratory research measurements of isoimmunisation. For example, the Kleihauer–Betke test is widely available and clinically used in contrast to new evidence that supports flow cytometry as the most accurate method of assessment. A more robust study should utilise flow cytometry to define sensitisation. Included studies in this review utilised the Kleihauer–Betke test which is more accessible and practical in clinical settings compared with flow cytometry. Up-to-date studies with accessible routine laboratory methods and standard definitions for sensitisation or follow-up times to appreciate sensitisation are needed rather than referring to dated studies with limited quality, such as the

ones included in this review. In the process of reinforcing standards of practice, practical guideline application is needed for potential global clinical practice change. This would include considerations not limited to: (a) anti-D administration logistics and practicalities, (b) access to instruments or tools to appreciate sensitisation and (c) access, including people self-managing abortion, as medical abortion is becoming more prevalent and thus do anti-D immunoglobulin recommendations help or hinder access to safe abortion care.

CONCLUSIONS

Rh immunoglobulin has well-documented safety. However, it is not without associated risks and costs, and it introduces barriers to accessing care that may impact especially on populations with restricted access (ie, legal, financial or resource-wise). The evidence base and quality of studies are currently limited, but there is some reassurance and experience from several national guidelines to justify no longer requiring its use in the first trimester. There is unclear benefit from the recommendation for Rh testing and immunoglobulin administration in early pregnancy. Clinical practice guidelines are based on expert opinions which are varied and moving away from testing and administration at the time of abortion. Therefore, more robust research would help give the reassurance needed to achieve substantial changes to guide clinical practice.

Twitter Michelle C Chan @chanmichellec

Acknowledgements The authors appreciate the help provided by Robyn Paynter, Cochrane Fertility Regulation Group, in the development of the search strategy. They also appreciate librarian assistance from the College of Physicians and Surgeons of British Columbia.

Contributors Three authors (MCC, CRK and RKG) independently screened all the titles, abstracts and full texts identified from the initial search to determine eligibility for inclusion. Conflicts were resolved through discussion and consensus first between MCC and RKG and, if further discussion was needed, with the third reviewer CRK. Risk of bias was assessed by MCC and CRK.

Funding UNDP/UNFPA/UNICEF/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP)

Disclaimer The authors alone are responsible for the views expressed in this review article and they do not necessarily represent the views, decisions or policies of the institutions with which they are affiliated.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different

terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Michelle C Chan <http://orcid.org/0000-0002-4671-1487>

REFERENCES

- Fiala C, Fux M, Gemzell Danielsson K. Rh-prophylaxis in early abortion. *Acta Obstet Gynecol Scand* 2003;82:892–903.
- Wiebe ER, Campbell M, Aiken ARA, *et al*. Can we safely stop testing for Rh status and immunizing Rh-negative women having early abortions? A comparison of Rh alloimmunization in Canada and the Netherlands. *Contraception* 2019;1:100001.
- Mark A, Foster AM, Grossman D, *et al*. Foregoing Rh testing and anti-D immunoglobulin for women presenting for early abortion: a recommendation from the National Abortion Federation's Clinical Policies Committee. *Contraception* 2019;99:265–6.
- National Institute for Health and Care Excellence. Abortion care - Review [C] anti-D prophylaxis for women up to 13+6 weeks' gestation (NICE Guideline NG140), 2019NICE. Available: <https://www.nice.org.uk/guidance/ng140/evidence/c-antid-prophylaxis-for-women-up-to-136-weeks-gestation-pdf-6905052975>
- Karanth L, Jaafar SH, Kanagasabai S, *et al*. Anti-D administration after spontaneous miscarriage for preventing rhesus alloimmunisation. *Cochrane Database Syst Rev* 2013;2013:CD009617.
- Moher D, Liberati A, Tetzlaff JAD. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement, 2019. Available: <https://www.equator-network.org/reporting-guidelines/prisma/> [Accessed 01 Aug 2019].
- Horvath S, Luning Prak ET, Schreiber CA. A highly sensitive flow cytometry protocol shows fetal red blood cell counts in first-trimester maternal circulation well below the threshold for Rh sensitization. *Contraception* 2018;98:332.
- Higgins JPT, Thomas J, Chandler J, eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.1*. Cochrane, 2020. www.training.cochrane.org/handbook
- Sterne JA, Hernán MA, Reeves BC, *et al*. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;355:i4919–10.
- Goldman JA, Eckerling B. Prevention of Rh immunization after abortion with anti-Rh (D)-immunoglobulin. *Obstet Gynecol* 1972;40:366–70.
- Gavin PS. Rhesus sensitization in abortion. *Obstet Gynecol* 1972;39:37–40.
- Hannafin B, Lovecchio F, Blackburn P. Do Rh-negative women with first trimester spontaneous abortions need Rh immune globulin? *Am J Emerg Med* 2006;24:487–9
- Jabara S, Barnhart KT. Is Rh immune globulin needed in early first-trimester abortion? A review. *Am J Obstet Gynecol* 2003;188:623–7.
- World Health Organization (WHO). *Safe abortion: technical and policy guidance for health systems*. 2nd edn. Geneva, Switzerland: WHO, 2012. <https://www.tandfonline.com/doi/full/10.1016/S0968-8080%2812%2939623-7>
- Moise K, Lockwood C, Kleinman SBarss V, ed. *Rhd alloimmunization in pregnancy: overview*, 2021. [https://www.uptodate.com/contents/rhd-alloimmunization-in-pregnancy-overview?search=rh negative&topicRef=6773&source=see_link](https://www.uptodate.com/contents/rhd-alloimmunization-in-pregnancy-overview?search=rh%20negative&topicRef=6773&source=see_link)
- Horvath S, Tsao P, Huang Z-Y, *et al*. The concentration of fetal red blood cells in first-trimester pregnant women undergoing uterine aspiration is below the calculated threshold for Rh sensitization. *Contraception* 2020;102:1–6.

Supplements:

Supplement 1 – Search Strategy

The following databases from their inception:

- EBM Reviews Ovid - Cochrane Central Register of Controlled Trials
- MEDLINE Ovid (Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily)
- Embase.com
- Popline <https://www.popline.org/advancedsearch>
- Google Scholar

The following trials registries:

- The World Health Organization International Clinical Trials Registry Platform www.who.int/trialsearch
- ClinicalTrials.gov www.clinicaltrials.gov.

Ovid MEDLINE(R) and Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily 1946 to July 22, 2019

Date searched: July 23, 2019

#	Searches	Results
1	Abortion, Induced/ or Abortion, Eugenic/ or Abortion, Legal/ or Abortion, Missed/ or Abortion, Therapeutic/ or Abortion, Criminal/ or Abortion Applicants/	40273
2	(abortion* or MTOP or VTOP or preabortion* or postabortion* or fetid* or foetid* or ((interrup* or terminat*) adj4 pregnan*) or (labo?r adj2 terminat*) or (menstrua* adj3 (induc* or regulat*))) .ti,ab,kf.	71537
3	Abortifacient Agents/ or Abortifacient Agents, Nonsteroidal/ or Abortifacient Agents, Steroidal/ or Menstruation-Inducing Agents/ or ((Carboprost/ or Dinoprost/ or Mifepristone/ or Misoprostol/ or Methotrexate/) and (Abortion, Induced/ or Abortion, Eugenic/ or Abortion, Legal/ or Abortion, Missed/ or Abortion, Therapeutic/ or Abortion, Criminal/ or Abortion Applicants/))	5190
4	(abortifacient* or ((mifepristone or misoprostol or methotrexate or dinoprost* or carboprost or sulprostone or gemeprost or meteneprost or lilopristone or onapristone or epostane or	5396

	oxytocin or prostaglandin*) and (abortion* or "interruption of pregnancy" or "termination of pregnancy" or pregnancy-termination*))) .ti,ab,kf.	
5	"Dilatation and Curettage"/ or Vacuum Curettage/	3000
6	("D&C" or "D & C" or "D/C" or D&E or "D & E" or D&X or "D & X" or MVA or ((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*))) .ti,ab,kf.	303323
7	or/1-6	373037
8	"Rh Isoimmunization"/ or "Rho(D) Immune Globulin"/ or "RHO(D) Antibody"/ or Isoantibodies/ or Isoantigens/ or Rh-Hr Blood-Group System/ or Fetomaternal Transfusion/	32528
9	(alloimmun* or allo-immun* or anti-D or anti-Rh or anti-RhD or ((Fetomaternal or fetomaternal or foetomaternal or foeto-maternal or transplacental or trans-placental) adj3 (haemorrhag* or hemorrhag* or transfusion)) or hydrops or ((Rh or RhD or "Rho(D)" or "Rho (D)" or "Rh o (D)" or RhIG or Rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or immunoglobulin* or immune-globulin or immunoprophylaxis or immuno-prophylaxis or isoantibod* or isoimmun* or iso-immun* or negative or prophyla* or sensiti*))) .ti,ab,kf.	24035
10	(WinRho or Rhogam).ti,ab,kf.	72
11	or/8-10	48774
12	and/7,11	2335
13	12 not ((exp Animals/ not Humans/) or (cat or cats or dog or dogs or macaque* or murine or moose or rabbit* or rat or rats or sheep).ti,ab,kf.)	1987

Embase.com

Date searched: July 23, 2019

#11	#10 AND [embase]/lim NOT ([embase]/lim AND [medline]/lim)	584
#10	#4 AND #8 AND [humans]/lim	1,964
#9	#4 AND #8	2,356
#8	#5 OR #6 OR #7	223,873

#7	wirho:ti,ab,kw OR rhogam:ti,ab,kw	113
#6	alloimmun*:ti,ab,kw OR 'allo immun*':ti,ab,kw OR 'anti d':ti,ab,kw OR 'anti rh':ti,ab,kw OR 'anti rhd':ti,ab,kw OR immunoglobulin*:ti,ab,kw OR 'immune globulin':ti,ab,kw OR immunoprophylaxis:ti,ab,kw OR isoantibod*:ti,ab,kw OR isoimmun*:ti,ab,kw OR 'iso immun*':ti,ab,kw OR (((rh OR rhd OR rhod OR 'rho d' OR 'rh o d' OR rhlg OR rhesus) NEAR/5 (hemoly* OR haemoly* OR incompat* OR immun* OR negativeOR prophyla* OR sensiti*)):ti,ab,kw)	216,255
#5	'rhesus isoimmunization'/exp OR 'rhesus antibody'/exp OR 'rhesus d antibody'/exp OR 'blood group rhesus system'/exp	14,837
#4	#1 OR #2 OR #3	145,876
#3	((dilation OR dilatation OR electric OR gestation OR manual OR pregnan* OR sharp OR suction OR surgical OR uterine OR uterus OR vacuum) NEAR/3 (aspiration* OR curettage OR evacuation* OR extraction OR terminat*)):ti,ab,kw	32,094
#2	abortion*:ti,ab,kw OR mtop:ti,ab,kw OR vtop:ti,ab,kw OR 'pre abortion*':ti,ab,kw OR preabortion*:ti,ab,kw OR 'post abortion*':ti,ab,kw OR postabortion*:ti,ab,kw OR feticid*:ti,ab,kw OR foeticid*:ti,ab,kw OR (((interrupt* OR terminat*) NEAR/4 pregnan*):ti,ab,kw)	84,086
#1	'induced abortion'/exp OR 'abortion'/exp	105,065

EBM Reviews - Cochrane Central Register of Controlled Trials June 2019

Date searched: July 23, 2019

#	Searches	Results
1	(abortion* or MTOP or VTOP or preabortion* or postabortion* or feticid* or foeticid* or ((interrup* or terminat*) adj4 pregnan*) or (menstrua* adj3 (induc* or regulat*))).ti,ab.	4280
2	((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*)).ti,ab.	2966
3	or/1-2	5627
4	(alloimmun* or allo-immun* or anti-D or anti-Rh or anti-RhD or ((Fetomaternal or feto-maternal or foetomaternal or foeto-maternal or transplacental or trans-placental) adj3 (haemorrhag* or hemorrhag* or transfusion)) or ((Rh or RhD or "Rho(D)" or "Rho (D)" or "Rh o (D)" or RhIG or Rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or immunoglobulin* or immune-	697

	globulin or immunoprophylaxis or immuno-prophylaxis or isoantibod* or isoimmun* or iso-immun* or negative or prophyla* or sensiti*))) .ti,ab.	
5	(WinRho or Rhogam).ti,ab.	10
6	or/4-5	702
7	and/3,6	26

POPLINE

Date searched: July 23, 2019

ALL FIELDS: abortion* OR MTOP OR VTOP OR preabortion* OR postabortion* OR feticid* OR foeticid* OR ((interrup* OR terminat*) AND pregnan*) OR (menstrua* AND (induc* OR regulat*)) OR ((dilation OR dilatation OR electric OR gestation OR manual OR pregnan* OR sharp OR suction OR surgical OR uterine OR uterus OR vacuum) AND (aspiration* OR curettage OR evacuation* OR extraction OR terminat*))
AND
ALL FIELDS: alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR anti-RhD OR ((Fetomaternal OR fetomaternal OR foetomaternal OR foeto-maternal OR transplacental OR trans-placental) AND (haemorrhag* OR hemorrhag* OR transfusion)) OR ((Rh OR RhD OR "Rho(D)" OR "Rho (D)" OR "Rh o (D)" OR RhIG OR Rhesus) AND (hemoly* OR haemoly* OR incompat* OR immun* OR immunoglobulin* OR immune-globulin OR immunoprophylaxis OR immuno-prophylaxis OR isoantibod* OR isoimmun* OR iso-immun* OR negative OR prophyla* OR sensiti*)) OR WinRho OR Rhogam
80 results

ClinicalTrials.gov

Date searched: July 23, 2019

abortion OR interruption OR termination OR aspiration OR extraction OR curettage OR evacuation (alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR Fetomaternal OR fetomaternal OR foetomaternal OR foeto-maternal OR RhD OR Rho(D) OR RhIG OR Rhesus OR immun* OR isoantibod* OR isoimmun* OR iso-immun* OR WinRho OR Rhogam)= 0 studies
--

WHO ICTRP

Date searched: July 23, 2019

TITLE: anti
AND
alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR anti-RhD OR Fetomaternal OR feto-maternal OR foetomaternal OR foeto-maternal OR Rh OR RhD OR Rho OR RhIG OR Rhesus
= 15 studies

Google Scholar

Date searched: July 23, 2019

(abortion* OR ((interrup* OR terminat*) AND pregnan*) OR ((dilation OR dilatation OR electric OR gestation OR manual OR pregnan* OR suction OR uterine OR uterus OR vacuum) AND (aspiration* OR curettage OR evacuation* OR extraction OR terminat*))) AND (alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR anti-RhD OR Fetomaternal OR feto-maternal OR foetomaternal OR foeto-maternal OR Rh OR RhD OR Rho OR RhIG OR Rhesus)

A search update was performed 13 May 2021, with no further studies meeting inclusion criteria.

Medline

Ovid MEDLINE(R) and Epub Ahead of Print, In-Process, In-Data-Review & Other Non-Indexed Citations and Daily 1946 to May 13, 2021

#	Searches	Results
1	Abortion, Induced/ or Abortion, Eugenic/ or Abortion, Legal/ or Abortion, Missed/ or Abortion, Therapeutic/ or Abortion, Criminal/ or Abortion Applicants/	41594
2	(abortion* or MTOP or VTOP or preabortion* or postabortion* or fetucid* or foeticid* or ((interrup* or terminat*) adj4 pregnan*) or (labo?r adj2 terminat*) or (menstrua* adj3 (induc* or regulat*))).ti,ab,kf.	76673
3	Abortifacient Agents/ or Abortifacient Agents, Nonsteroidal/ or Abortifacient Agents, Steroidal/ or Menstruation-Inducing Agents/ or ((Carboprost/ or Dinoprost/ or Mifepristone/ or Misoprostol/ or Methotrexate/) and (Abortion, Induced/ or Abortion, Eugenic/ or Abortion, Legal/ or Abortion, Missed/ or Abortion, Therapeutic/ or Abortion, Criminal/ or Abortion Applicants/))	5423
4	(abortifacient* or ((mifepristone or misoprostol or methotrexate or dinoprost* or carboprost or sulprostone or gemeprost or meteneprost or lilopristone or onapristone or epostane or oxytocin or prostaglandin*) and (abortion* or "interruption of pregnancy" or "termination of pregnancy" or pregnancy-termination*))).ti,ab,kf.	5695
5	"Dilatation and Curettage"/ or Vacuum Curettage/	3102
6	("D&C" or "D & C" or "D/C" or D&E or "D & E" or D&X or "D & X" or MVA or ((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*))).ti,ab,kf.	360810

7	1 or 2 or 3 or 4 or 5 or 6	434642
8	"Rh Isoimmunization"/ or "Rho(D) Immune Globulin"/ or "RHO(D) Antibody"/ or Isoantibodies/ or Isoantigens/ or Rh-Hr Blood-Group System/ or Fetomaternal Transfusion/	33384
9	(alloimmun* or allo-immun* or anti-D or anti-Rh or anti-RhD or ((Fetomaternal or feto maternal or foetomaternal or foeto-maternal or transplacental or trans-placental) adj3 (haemorrhag* or hemorrhag* or transfusion)) or hydrops or ((Rh or RhD or "Rho(D)" or "Rho (D)" or "Rh o (D)" or RhIG or Rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or immunoglobulin* or immunoglobulin or immunoprophylaxis or immuno-prophylaxis or isoantibod* or isoimmun* or iso-immun* or negative or prophyla* or sensiti*))) .ti,ab,kf.	25777
10	(WinRho or Rhogam).ti,ab,kf.	75
11	8 or 9 or 10	51075
12	7 and 11	2545
13	12 not ((exp Animals/ not Humans/) or (cat or cats or dog or dogs or macaque* or murine or moose or rabbit* or rat or rats or sheep).ti,ab,kf.)	2182
14	limit 13 to dt=20190723-20210514	141

In line 14, I used the Create Date – the day the citation record is entered into Medline. This will capture everything published since July 23, 2019, but also anything published slightly before July 23 that was added at a later time.

Embase

I only have access to Embase through the Ovid platform. The search results followed a fairly similar pattern to what I could see in your previous Embase.com results. While I did not remove Medline articles in the search, I manually removed them afterward. In line 13, I removed one result that overlapped with the Medline results.

Embase 1974 to 2021 May 13 (Ovid Version)

#	Searches	Results
1	exp induced abortion/ or exp abortion/	104290
2	(abortion* or mtop or vtop or "pre abortion*" or preabortion* or "post abortion*" or postabortion or fetucid* or foeticid* or ((interrupt* or terminat*) adj4 pregnan*).ti,ab,kw.	84873
3	((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*).ti,ab,kw.	35104

4	1 or 2 or 3	148743
5	exp rhesus isoimmunization/ or exp rhesus antibody/ or exp rhesus d antibody/ or exp blood group rhesus system/	10242
6	(alloimmun* or "allo immun*" or "anti d" or "anti rh" or "anti rhd" or immunoglobulin* or "immune globulin" or immunoprophylaxis or isoantibod* or isoimmun* or "iso immun*" or ((rh or rhd or rhod or "rho d" or "rh o d" or rhlg or rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or negative or prophyla* or sensiti*))) .ti,ab,kw.	234457
7	(winrho or rhogam).ti,ab,kw.	114
8	5 or 6 or 7	238756
9	4 and 8	2354
10	limit 9 to human	1954
11	limit 10 to dd=20190723-20210514	78
12	from 11 keep 5	1
13	11 not 12	77

In line 10, I limited the search using the Date Delivered field – which acts similarly to the Create Date in Ovid Medline.

EBM Reviews

EBM Reviews - Cochrane Central Register of Controlled Trials April 2021

#	Searches	Results
1	(abortion* or MTOP or VTOP or preabortion* or postabortion* or fetid* or foetid* or ((interrup* or terminat*) adj4 pregnan*) or (menstrua* adj3 (induc* or regulat*))) .ti,ab.	5027
2	((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*)) .ti,ab.	3551
3	1 or 2	6719
4	(alloimmun* or allo-immun* or anti-D or anti-Rh or anti-RhD or ((Fetomaternal or feto-maternal or foetomaternal or foeto-maternal or transplacental or trans-placental) adj3 (haemorrhag* or hemorrhag* or transfusion)) or ((Rh or RhD or "Rho(D)" or "Rho (D)" or "Rh o (D)" or RhIG or Rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or immunoglobulin* or immune-globulin or	774

	immunoprophylaxis or immuno-prophylaxis or isoantibod* or isoimmun* or iso immun* or negative or prophyla* or sensiti*)))ti,ab.	
5	(WinRho or Rhogam).ti,ab.	10
6	4 or 5	778
7	3 and 6	26

No new results

POPLINE

Unfortunately, this database was retired in September 2019, so I was unable to search it.

ClinicalTrials.gov

Broken down into 2 searches (to fit the character limit):

(abortion OR interruption OR termination OR aspiration OR extraction OR curettage OR evacuation) AND (alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR Fetomaternal OR fetomaternal OR foetomaternal OR foeto-maternal OR RhD OR Rho(D) OR RhIG)

(abortion OR interruption OR termination OR aspiration OR extraction OR curettage OR evacuation) AND (Rhesus OR immun* OR isoantibod* OR isoimmun* OR iso-immun* OR WinRho OR Rhogam)

Found 1 result:

https://clinicaltrials.gov/ct2/show/NCT04701034?cond=%28abortion+OR+interruption+OR+termination+OR+aspiration+OR+extraction+OR+curettage+OR+evacuation%29+AND+%28Rhesus+OR+immun*+OR+isoantibod*+OR+isoimmun*+OR+iso-immun*+OR+WinRho+OR+Rhogam%29&draw=2&rank=1

WHO ICTRP

Searching in the Title field:

anti AND alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR anti-RhD OR Fetomaternal OR fetomaternal OR foetomaternal OR foeto-maternal OR Rh OR RhD OR Rho OR RhIG OR Rhesus

Results:

Main ID	Public Title	Date of Registration
NCT04734379	Rho Kinase (ROCK) Inhibitor in Tauopathies - 1	2021-01-18
NCT04682912	Blood Types in Children With COVID-19	2020-12-22
NCT04600674	Delayed Cord Clamping in Rhesus Disease of the Newborn	2020-10-19
JPRN-jRCT2052200064	Effectiveness and Safety of RH-01 in Patients with Cardiac Disease and Indication of Cardiac Rehabilitation: A Multicenter Parallel-group Randomized Control Trial	2020-10-13
CTRI/2020/09/027588	Association of ABO Blood Group And RH Factor With Facial Characteristics Represented by Mandibular Position And Mandibular Angular Relation - An Observational Study.	2020-09-03
NCT04484896	Message Framing on Recruiting Rh-D Negative Blood Donors	2020-07-21
ChiCTR2000033786	Effect and mechanism of micro negative pressure therapy on burn wound: a randomized controlled trial.	2020-06-12
NCT04364308	The Relationship Between Umbilical Cord ph and Feto-maternal Doppler Studies in Scheduled Nonlaboring Term Singleton Caesarean Deliveries	2020-04-19
NCT04556188	The Clinical Influence of Developing a Sustainable	2020-03-17

	Cardiac Surgery Service to Reduce the Burden of Rheumatic Heart Disease in Sub-Saharan Africa	
NCT04285398	Prospective Natural History Study of Retinitis Pigmentosa	2020-02-24
NCT04234932	Short-term Effect of Rho-kinase Inhibitor on Retinal Circulation	2020-01-15
NCT04156893	RH Genotype Matched RBC Transfusions	2019-11-06
NCT04156906	RHD Genotype Matching for Anti-D	2019-11-06
NCT04120883	Oral Hydroxychloroquine (HCQ) for Retinitis Pigmentosa Caused by P23H-Rhodopsin (RHO)	2019-10-08
NCT04123626	A Study to Evaluate the Safety and Tolerability of QR-1123 in Subjects With Autosomal Dominant Retinitis Pigmentosa Due to the P23H Mutation in the RHO Gene	2019-10-01

Google Scholar was not repeated in the updated search, given it is not a database.

To enable PROSPERO to focus on COVID-19 submissions, this registration record has undergone basic automated checks for eligibility and is published exactly as submitted. PROSPERO has never provided peer review, and usual checking by the PROSPERO team does not endorse content. Therefore, automatically published records should be treated as any other PROSPERO registration. Further detail is provided [here](#).

Citation

Michelle Chan, Caron Kim, Roopan Gill. A systematic review: Rh isoimmunization in unsensitized Rh negative individuals seeking abortion <12 weeks. PROSPERO 2020 CRD42020149073 Available from: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020149073

Review question

Among women who undergo an abortion, medical or surgical, what is the rate of isoimmunization in subsequent pregnancy.

Searches

Run on 23 July 2019 by Fertility Regulation Group Information Specialist, Robin Paynter (MLIS), conducted a search for all published, unpublished, and ongoing studies, without restrictions on language or publication status.

The following databases from their inception:

- EBM Reviews Ovid - Cochrane Central Register of Controlled Trials (CENTRAL)
- MEDLINE Ovid (Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily)
- Embase.com
- Popline <https://www.popline.org/advancedsearch>
- Google Scholar

The following trials registries:

- The World Health Organization International Clinical Trials Registry Platform www.who.int/trialsearch
- ClinicalTrials.gov www.ClinicalTrials.gov.

Types of study to be included

Types of studies:

Included: Primary studies including RCTs, controlled trials, cohort studies, case-control studies

Excluded: Case series, review articles, editorials, letters, advisories, non-comparative studies, unpublished manuscripts, conference abstracts, diagnostic studies, animal studies, cost-benefit analysis, study with basic science outcomes

- Date cut off: nothing before 1971 *WHO recommended in this year to its nations that Rh testing and treatment with immune globulin be made part of the standard protocol of medical care for pregnant women

- Non-English publications

Condition or domain being studied

In most countries anti-D Immunoglobulin G is provided for rhesus (Rh)-negative patients. The administration

of Rh-immunoglobulin was first introduced in 1968 and significantly reduced immunization to D-antigen (Fiala 2003). There is sparse evidence for such intervention after abortion in early pregnancy (Fiala 2003). Early abortion includes threatened, spontaneous, surgical or medical abortion.

Currently, the WHO recommends Rh testing and administration of anti-D in populations with higher prevalence of Rh-negative individuals. Recently, the National Abortion Federation of North America published a statement that women presenting for early abortion can forego Rh testing and anti-D immunoglobulin administration (Mark 2019).

Participants/population

Reproductive aged women who have recently undergone surgical or medical management of pregnancy <12 weeks (includes induced, incomplete, spontaneous, or septic abortion)

Intervention(s), exposure(s)

Intervention:

Routine administration of Anti-D

Comparator(s)/control

Comparisons:

No administration of Anti-D

Context

Global setting

Main outcome(s)

Primary outcomes

1. Development of a positive Kleihauer Betke test (a test that detects fetal cells in the maternal blood).
2. Development of RhD alloimmunisation in a subsequent pregnancy.

Measures of effect

As defined by studies

Additional outcome(s)

Secondary outcomes

1. Detection of atypical blood group antibodies by positive indirect Coombs test after six months of exposure (non?prespecified outcome).
2. Need for increased surveillance for suspected fetal blood sampling and fetal transfusions in subsequent pregnancies.
3. Neonatal morbidity such as neonatal anaemia, jaundice, bilirubin encephalopathy, erythroblastosis, prematurity, hypoglycaemia (low blood sugar) in subsequent pregnancies.
4. Maternal adverse events of anti?D administration including anaphylactic reaction

Measures of effect

As defined by studies

Data extraction (selection and coding)

With less than 20 articles, all data extraction will be performed by Dr. Michelle Chan. Data extraction will be

obtained directly from the original published article. Recording of information for the data abstraction will occur in a table format, with the following headings. (1) Author, Year, Funding (2) Study Design, location, year(s) of data collection, follow-up (3) Population (4) Intervention (5) Comparison (6) Results (7) Strengths (8) Weaknesses (9) Risk of Bias

Risk of bias (quality) assessment

Risk of bias quality assessment will be addressed with criteria depending on study design.

RCT - List of factors for RCT quality framework: Adequate randomization (computer generated); Allocation concealment (opaque sealed envelopes); Blinding of participants and personnel; Blinding of outcome assessment;

Objective measurement of outcome; Incomplete data/lost to follow-up; Small sample size (need to define the size); Power calculations by appropriate outcome measure.

Observational studies - List of factors for observational quality framework: Population, Participation and follow up rates as above; Defined exposure (esp with comparator); Exposure defining, duration, timing; Measurement of isoimmunization; Confounding issue

Strategy for data synthesis

Data synthesis will primarily be descriptive analysis based on the outcomes. Pooled data will be reviewed with the GRADE system.

Given the anticipated small number of articles, this review will be primarily narrative. A descriptive analysis will provide summary of the studies including: study location, years of data collection, follow-up, population, intervention, comparison, results, strengths and weaknesses. Risk of bias will also be included.

A broader summary and tables of primary outcomes:

1. Development of a positive Kleihauer Betke test (a test that detects fetal cells in the maternal blood).
2. Development of RhD alloimmunisation in a subsequent pregnancy.

and secondary outcomes:

1. Detection of atypical blood group antibodies by positive indirect Coombs test after six months of exposure (non?prespecified outcome).
2. Need for increased surveillance for suspected fetal blood sampling and fetal transfusions in subsequent pregnancies.
3. Neonatal morbidity such as neonatal anaemia, jaundice, bilirubin encephalopathy, erythroblastosis, prematurity, hypoglycaemia (low blood sugar) in subsequent pregnancies.
4. Maternal adverse events of anti?D administration including anaphylactic reaction

and secondary outcomes (will also be included.

Analysis of subgroups or subsets

Not applicable

Contact details for further information

Michelle C Chan
michelle.c.chan@gmail.com

Organisational affiliation of the review

University of British Columbia/ World Health Organization, Department of Reproductive Health & Research

Review team members and their organisational affiliations

Dr Michelle Chan. University of British Columbia/ World Health Organization, Department of Reproductive Health & Research

Dr Caron Kim. World Health Organization, Department of Reproductive Health & Research

Dr Roopan Gill. World Health Organization, Department of Reproductive Health & Research

Collaborators

Ms Robyn Paynter. Cochrane Fertility Regulation Group Information Specialist, MLS

Type and method of review

Systematic review

Anticipated or actual start date

23 July 2019

Anticipated completion date

29 February 2020

Funding sources/sponsors

Not applicable, no funders

Conflicts of interest

Language

English

Country

Canada, Switzerland, United States of America

Stage of review

Review Ongoing

Subject index terms status

Subject indexing assigned by CRD

Subject index terms

Abortion, Induced; Abortion, Spontaneous; Female; Humans; Pregnancy; Rh Isoimmunization

Date of registration in PROSPERO

28 April 2020

Date of first submission

30 August 2019

Stage of review at time of this submission

Stage	Started	Completed
Preliminary searches	Yes	No
Piloting of the study selection process	Yes	No
Formal screening of search results against eligibility criteria	Yes	No
Data extraction	Yes	No
Risk of bias (quality) assessment	No	No
Data analysis	No	No

The record owner confirms that the information they have supplied for this submission is accurate and complete and they understand that deliberate provision of inaccurate information or omission of data may be construed as scientific misconduct.

The record owner confirms that they will update the status of the review when it is completed and will add publication details in due course.

Versions

28 April 2020

Supplements:

Supplement 1 – Search Strategy

The following databases from their inception:

- EBM Reviews Ovid - Cochrane Central Register of Controlled Trials
- MEDLINE Ovid (Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily)
- Embase.com
- Popline <https://www.popline.org/advancedsearch>
- Google Scholar

The following trials registries:

- The World Health Organization International Clinical Trials Registry Platform www.who.int/trialsearch
- ClinicalTrials.gov www.clinicaltrials.gov.

Ovid MEDLINE(R) and Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily 1946 to July 22, 2019

Date searched: July 23, 2019

#	Searches	Results
1	Abortion, Induced/ or Abortion, Eugenic/ or Abortion, Legal/ or Abortion, Missed/ or Abortion, Therapeutic/ or Abortion, Criminal/ or Abortion Applicants/	40273
2	(abortion* or MTOP or VTOP or preabortion* or postabortion* or fetid* or foetid* or ((interrup* or terminat*) adj4 pregnan*) or (labo?r adj2 terminat*) or (menstrua* adj3 (induc* or regulat*))) .ti,ab,kf.	71537
3	Abortifacient Agents/ or Abortifacient Agents, Nonsteroidal/ or Abortifacient Agents, Steroidal/ or Menstruation-Inducing Agents/ or ((Carboprost/ or Dinoprost/ or Mifepristone/ or Misoprostol/ or Methotrexate/) and (Abortion, Induced/ or Abortion, Eugenic/ or Abortion, Legal/ or Abortion, Missed/ or Abortion, Therapeutic/ or Abortion, Criminal/ or Abortion Applicants/))	5190
4	(abortifacient* or ((mifepristone or misoprostol or methotrexate or dinoprost* or carboprost or sulprostone or gemeprost or meteneprost or lilopristone or onapristone or epostane or	5396

	oxytocin or prostaglandin*) and (abortion* or "interruption of pregnancy" or "termination of pregnancy" or pregnancy-termination*))) .ti,ab,kf.	
5	"Dilatation and Curettage"/ or Vacuum Curettage/	3000
6	("D&C" or "D & C" or "D/C" or D&E or "D & E" or D&X or "D & X" or MVA or ((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*))) .ti,ab,kf.	303323
7	or/1-6	373037
8	"Rh Isoimmunization"/ or "Rho(D) Immune Globulin"/ or "RHO(D) Antibody"/ or Isoantibodies/ or Isoantigens/ or Rh-Hr Blood-Group System/ or Fetomaternal Transfusion/	32528
9	(alloimmun* or allo-immun* or anti-D or anti-Rh or anti-RhD or ((Fetomaternal or fetomaternal or foetomaternal or foeto-maternal or transplacental or trans-placental) adj3 (haemorrhag* or hemorrhag* or transfusion)) or hydrops or ((Rh or RhD or "Rho(D)" or "Rho(D)" or "Rh o (D)" or RhIG or Rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or immunoglobulin* or immune-globulin or immunoprophylaxis or immuno-prophylaxis or isoantibod* or isoimmun* or iso-immun* or negative or prophyla* or sensiti*))) .ti,ab,kf.	24035
10	(WinRho or Rhogam).ti,ab,kf.	72
11	or/8-10	48774
12	and/7,11	2335
13	12 not ((exp Animals/ not Humans/) or (cat or cats or dog or dogs or macaque* or murine or moose or rabbit* or rat or rats or sheep).ti,ab,kf.)	1987

Embase.com

Date searched: July 23, 2019

#11	#10 AND [embase]/lim NOT ([embase]/lim AND [medline]/lim)	584
#10	#4 AND #8 AND [humans]/lim	1,964
#9	#4 AND #8	2,356
#8	#5 OR #6 OR #7	223,873

#7	winrho:ti,ab,kw OR rhogam:ti,ab,kw	113
#6	alloimmun*:ti,ab,kw OR 'allo immun*':ti,ab,kw OR 'anti d':ti,ab,kw OR 'anti rh':ti,ab,kw OR 'anti rhd':ti,ab,kw OR immunoglobulin*:ti,ab,kw OR 'immune globulin':ti,ab,kw OR immunoprophylaxis:ti,ab,kw OR isoantibod*:ti,ab,kw OR isoimmun*:ti,ab,kw OR 'iso immun*':ti,ab,kw OR (((rh OR rhd OR rhod OR 'rho d' OR 'rh o d' OR rhlg OR rhesus) NEAR/5 (hemoly* OR haemoly* OR incompat* OR immun* OR negativeOR prophyla* OR sensiti*)):ti,ab,kw)	216,255
#5	'rhesus isoimmunization'/exp OR 'rhesus antibody'/exp OR 'rhesus d antibody'/exp OR 'blood group rhesus system'/exp	14,837
#4	#1 OR #2 OR #3	145,876
#3	((dilation OR dilatation OR electric OR gestation OR manual OR pregnan* OR sharp OR suction OR surgical OR uterine OR uterus OR vacuum) NEAR/3 (aspiration* OR curettage OR evacuation* OR extraction OR terminat*)):ti,ab,kw	32,094
#2	abortion*:ti,ab,kw OR mtop:ti,ab,kw OR vtop:ti,ab,kw OR 'pre abortion*':ti,ab,kw OR preabortion*:ti,ab,kw OR 'post abortion*':ti,ab,kw OR postabortion*:ti,ab,kw OR feticid*:ti,ab,kw OR foeticid*:ti,ab,kw OR (((interrupt* OR terminat*) NEAR/4 pregnan*):ti,ab,kw)	84,086
#1	'induced abortion'/exp OR 'abortion'/exp	105,065

EBM Reviews - Cochrane Central Register of Controlled Trials June 2019

Date searched: July 23, 2019

#	Searches	Results
1	(abortion* or MTOP or VTOP or preabortion* or postabortion* or feticid* or foeticid* or ((interrup* or terminat*) adj4 pregnan*) or (menstrua* adj3 (induc* or regulat*))).ti,ab.	4280
2	((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*)).ti,ab.	2966
3	or/1-2	5627
4	(alloimmun* or allo-immun* or anti-D or anti-Rh or anti-RhD or ((Fetomaternal or feto-maternal or foetomaternal or foeto-maternal or transplacental or trans-placental) adj3 (haemorrhag* or hemorrhag* or transfusion)) or ((Rh or RhD or "Rho(D)" or "Rho (D)" or "Rh o (D)" or RhIG or Rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or immunoglobulin* or immune-	697

	globulin or immunoprophylaxis or immuno-prophylaxis or isoantibod* or isoimmun* or iso-immun* or negative or prophyla* or sensiti*))) .ti,ab.	
5	(WinRho or Rhogam).ti,ab.	10
6	or/4-5	702
7	and/3,6	26

POPLINE

Date searched: July 23, 2019

ALL FIELDS: abortion* OR MTOP OR VTOP OR preabortion* OR postabortion* OR fetid* OR foetid* OR ((interrup* OR terminat*) AND pregnan*) OR (menstrua* AND (induc* OR regulat*)) OR ((dilation OR dilatation OR electric OR gestation OR manual OR pregnan* OR sharp OR suction OR surgical OR uterine OR uterus OR vacuum) AND (aspiration* OR curettage OR evacuation* OR extraction OR terminat*))
AND
ALL FIELDS: alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR anti-RhD OR ((Fetomaternal OR fetomaternal OR foetomaternal OR foeto-maternal OR transplacental OR trans-placental) AND (haemorrhag* OR hemorrhag* OR transfusion)) OR ((Rh OR RhD OR "Rho(D)" OR "Rho (D)" OR "Rh o (D)" OR RhIG OR Rhesus) AND (hemoly* OR haemoly* OR incompat* OR immun* OR immunoglobulin* OR immune-globulin OR immunoprophylaxis OR immuno-prophylaxis OR isoantibod* OR isoimmun* OR iso-immun* OR negative OR prophyla* OR sensiti*)) OR WinRho OR Rhogam
80 results

ClinicalTrials.gov

Date searched: July 23, 2019

abortion OR interruption OR termination OR aspiration OR extraction OR curettage OR evacuation (alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR Fetomaternal OR fetomaternal OR foetomaternal OR foeto-maternal OR RhD OR Rho(D) OR RhIG OR Rhesus OR immun* OR isoantibod* OR isoimmun* OR iso-immun* OR WinRho OR Rhogam)= 0 studies
--

WHO ICTRP

Date searched: July 23, 2019

TITLE: anti
AND
alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR anti-RhD OR Fetomaternal OR feto-maternal OR foetomaternal OR foeto-maternal OR Rh OR RhD OR Rho OR RhIG OR Rhesus
= 15 studies

Google Scholar

Date searched: July 23, 2019

(abortion* OR ((interrup* OR terminat*) AND pregnan*) OR ((dilation OR dilatation OR electric OR gestation OR manual OR pregnan* OR suction OR uterine OR uterus OR vacuum) AND (aspiration* OR curettage OR evacuation* OR extraction OR terminat*))) AND (alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR anti-RhD OR Fetomaternal OR feto-maternal OR foetomaternal OR foeto-maternal OR Rh OR RhD OR Rho OR RhIG OR Rhesus)

A search update was performed 13 May 2021, with no further studies meeting inclusion criteria.

Medline

Ovid MEDLINE(R) and Epub Ahead of Print, In-Process, In-Data-Review & Other Non-Indexed Citations and Daily 1946 to May 13, 2021

#	Searches	Results
1	Abortion, Induced/ or Abortion, Eugenic/ or Abortion, Legal/ or Abortion, Missed/ or Abortion, Therapeutic/ or Abortion, Criminal/ or Abortion Applicants/	41594
2	(abortion* or MTOP or VTOP or preabortion* or postabortion* or fetucid* or foeticid* or ((interrup* or terminat*) adj4 pregnan*) or (labo?r adj2 terminat*) or (menstrua* adj3 (induc* or regulat*))).ti,ab,kf.	76673
3	Abortifacient Agents/ or Abortifacient Agents, Nonsteroidal/ or Abortifacient Agents, Steroidal/ or Menstruation-Inducing Agents/ or ((Carboprost/ or Dinoprost/ or Mifepristone/ or Misoprostol/ or Methotrexate/) and (Abortion, Induced/ or Abortion, Eugenic/ or Abortion, Legal/ or Abortion, Missed/ or Abortion, Therapeutic/ or Abortion, Criminal/ or Abortion Applicants/))	5423
4	(abortifacient* or ((mifepristone or misoprostol or methotrexate or dinoprost* or carboprost or sulprostone or gemeprost or meteneprost or lilopristone or onapristone or epostane or oxytocin or prostaglandin*) and (abortion* or "interruption of pregnancy" or "termination of pregnancy" or pregnancy-termination*))).ti,ab,kf.	5695
5	"Dilatation and Curettage"/ or Vacuum Curettage/	3102
6	("D&C" or "D & C" or "D/C" or D&E or "D & E" or D&X or "D & X" or MVA or ((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*))).ti,ab,kf.	360810

7	1 or 2 or 3 or 4 or 5 or 6	434642
8	"Rh Isoimmunization"/ or "Rho(D) Immune Globulin"/ or "RHO(D) Antibody"/ or Isoantibodies/ or Isoantigens/ or Rh-Hr Blood-Group System/ or Fetomaternal Transfusion/	33384
9	(alloimmun* or allo-immun* or anti-D or anti-Rh or anti-RhD or ((Fetomaternal or feto maternal or foetomaternal or foeto-maternal or transplacental or trans-placental) adj3 (haemorrhag* or hemorrhag* or transfusion)) or hydrops or ((Rh or RhD or "Rho(D)" or "Rho (D)" or "Rh o (D)" or RhIG or Rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or immunoglobulin* or immunoglobulin or immunoprophylaxis or immuno-prophylaxis or isoantibod* or isoimmun* or iso-immun* or negative or prophyla* or sensiti*))) .ti,ab,kf.	25777
10	(WinRho or Rhogam).ti,ab,kf.	75
11	8 or 9 or 10	51075
12	7 and 11	2545
13	12 not ((exp Animals/ not Humans/) or (cat or cats or dog or dogs or macaque* or murine or moose or rabbit* or rat or rats or sheep).ti,ab,kf.)	2182
14	limit 13 to dt=20190723-20210514	141

In line 14, I used the Create Date – the day the citation record is entered into Medline. This will capture everything published since July 23, 2019, but also anything published slightly before July 23 that was added at a later time.

Embase

I only have access to Embase through the Ovid platform. The search results followed a fairly similar pattern to what I could see in your previous Embase.com results. While I did not remove Medline articles in the search, I manually removed them afterward. In line 13, I removed one result that overlapped with the Medline results.

Embase 1974 to 2021 May 13 (Ovid Version)

#	Searches	Results
1	exp induced abortion/ or exp abortion/	104290
2	(abortion* or mtop or vtop or "pre abortion*" or preabortion* or "post abortion*" or postabortion or fetucid* or foeticid* or ((interrupt* or terminat*) adj4 pregnan*).ti,ab,kw.	84873
3	((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*).ti,ab,kw.	35104

4	1 or 2 or 3	148743
5	exp rhesus isoimmunization/ or exp rhesus antibody/ or exp rhesus d antibody/ or exp blood group rhesus system/	10242
6	(alloimmun* or "allo immun*" or "anti d" or "anti rh" or "anti rhd" or immunoglobulin* or "immune globulin" or immunoprophylaxis or isoantibod* or isoimmun* or "iso immun*" or ((rh or rhd or rhod or "rho d" or "rh o d" or rhlg or rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or negative or prophyla* or sensiti*))) .ti,ab,kw.	234457
7	(winrho or rhogam).ti,ab,kw.	114
8	5 or 6 or 7	238756
9	4 and 8	2354
10	limit 9 to human	1954
11	limit 10 to dd=20190723-20210514	78
12	from 11 keep 5	1
13	11 not 12	77

In line 10, I limited the search using the Date Delivered field – which acts similarly to the Create Date in Ovid Medline.

EBM Reviews

EBM Reviews - Cochrane Central Register of Controlled Trials April 2021

#	Searches	Results
1	(abortion* or MTOP or VTOP or preabortion* or postabortion* or fetid* or foetid* or ((interrup* or terminat*) adj4 pregnan*) or (menstrua* adj3 (induc* or regulat*))) .ti,ab.	5027
2	((dilation or dilatation or electric or gestation or manual or pregnan* or sharp or suction or surgical or uterine or uterus or vacuum) adj3 (aspiration* or curettage or evacuation* or extraction or terminat*)) .ti,ab.	3551
3	1 or 2	6719
4	(alloimmun* or allo-immun* or anti-D or anti-Rh or anti-RhD or ((Fetomaternal or feto-maternal or foetomaternal or foeto-maternal or transplacental or trans-placental) adj3 (haemorrhag* or hemorrhag* or transfusion)) or ((Rh or RhD or "Rho(D)" or "Rho (D)" or "Rh o (D)" or RhIG or Rhesus) adj5 (hemoly* or haemoly* or incompat* or immun* or immunoglobulin* or immune-globulin or	774

	immunoprophylaxis or immuno-prophylaxis or isoantibod* or isoimmun* or iso immun* or negative or prophyla* or sensiti*)))ti,ab.	
5	(WinRho or Rhogam).ti,ab.	10
6	4 or 5	778
7	3 and 6	26

No new results

POPLINE

Unfortunately, this database was retired in September 2019, so I was unable to search it.

ClinicalTrials.gov

Broken down into 2 searches (to fit the character limit):

(abortion OR interruption OR termination OR aspiration OR extraction OR curettage OR evacuation) AND (alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR Fetomaternal OR fetomaternal OR foetomaternal OR foeto-maternal OR RhD OR Rho(D) OR RhIG)

(abortion OR interruption OR termination OR aspiration OR extraction OR curettage OR evacuation) AND (Rhesus OR immun* OR isoantibod* OR isoimmun* OR iso-immun* OR WinRho OR Rhogam)

Found 1 result:

https://clinicaltrials.gov/ct2/show/NCT04701034?cond=%28abortion+OR+interruption+OR+termination+OR+aspiration+OR+extraction+OR+curettage+OR+evacuation%29+AND+%28Rhesus+OR+immun*+OR+isoantibod*+OR+isoimmun*+OR+iso-immun*+OR+WinRho+OR+Rhogam%29&draw=2&rank=1

WHO ICTRP

Searching in the Title field:

anti AND alloimmun* OR allo-immun* OR anti-D OR anti-Rh OR anti-RhD OR Fetomaternal OR fetomaternal OR foetomaternal OR foeto-maternal OR Rh OR RhD OR Rho OR RhIG OR Rhesus

Results:

Main ID	Public Title	Date of Registration
NCT04734379	Rho Kinase (ROCK) Inhibitor in Tauopathies - 1	2021-01-18
NCT04682912	Blood Types in Children With COVID-19	2020-12-22
NCT04600674	Delayed Cord Clamping in Rhesus Disease of the Newborn	2020-10-19
JPRN-jRCT2052200064	Effectiveness and Safety of RH-01 in Patients with Cardiac Disease and Indication of Cardiac Rehabilitation: A Multicenter Parallel-group Randomized Control Trial	2020-10-13
CTRI/2020/09/027588	Association of ABO Blood Group And RH Factor With Facial Characteristics Represented by Mandibular Position And Mandibular Angular Relation - An Observational Study.	2020-09-03
NCT04484896	Message Framing on Recruiting Rh-D Negative Blood Donors	2020-07-21
ChiCTR2000033786	Effect and mechanism of micro negative pressure therapy on burn wound: a randomized controlled trial.	2020-06-12
NCT04364308	The Relationship Between Umbilical Cord ph and Feto-maternal Doppler Studies in Scheduled Nonlaboring Term Singleton Caesarean Deliveries	2020-04-19
NCT04556188	The Clinical Influence of Developing a Sustainable	2020-03-17

	Cardiac Surgery Service to Reduce the Burden of Rheumatic Heart Disease in Sub-Saharan Africa	
NCT04285398	Prospective Natural History Study of Retinitis Pigmentosa	2020-02-24
NCT04234932	Short-term Effect of Rho-kinase Inhibitor on Retinal Circulation	2020-01-15
NCT04156893	RH Genotype Matched RBC Transfusions	2019-11-06
NCT04156906	RHD Genotype Matching for Anti-D	2019-11-06
NCT04120883	Oral Hydroxychloroquine (HCQ) for Retinitis Pigmentosa Caused by P23H-Rhodopsin (RHO)	2019-10-08
NCT04123626	A Study to Evaluate the Safety and Tolerability of QR-1123 in Subjects With Autosomal Dominant Retinitis Pigmentosa Due to the P23H Mutation in the RHO Gene	2019-10-01

Google Scholar was not repeated in the updated search, given it is not a database.

To enable PROSPERO to focus on COVID-19 submissions, this registration record has undergone basic automated checks for eligibility and is published exactly as submitted. PROSPERO has never provided peer review, and usual checking by the PROSPERO team does not endorse content. Therefore, automatically published records should be treated as any other PROSPERO registration. Further detail is provided [here](#).

Citation

Michelle Chan, Caron Kim, Roopan Gill. A systematic review: Rh isoimmunization in unsensitized Rh negative individuals seeking abortion <12 weeks. PROSPERO 2020 CRD42020149073 Available from: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020149073

Review question

Among women who undergo an abortion, medical or surgical, what is the rate of isoimmunization in subsequent pregnancy.

Searches

Run on 23 July 2019 by Fertility Regulation Group Information Specialist, Robin Paynter (MLIS), conducted a search for all published, unpublished, and ongoing studies, without restrictions on language or publication status.

The following databases from their inception:

- EBM Reviews Ovid - Cochrane Central Register of Controlled Trials (CENTRAL)
- MEDLINE Ovid (Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily)
- Embase.com
- Popline <https://www.popline.org/advancedsearch>
- Google Scholar

The following trials registries:

- The World Health Organization International Clinical Trials Registry Platform www.who.int/trialsearch
- ClinicalTrials.gov www.ClinicalTrials.gov.

Types of study to be included

Types of studies:

Included: Primary studies including RCTs, controlled trials, cohort studies, case-control studies

Excluded: Case series, review articles, editorials, letters, advisories, non-comparative studies, unpublished manuscripts, conference abstracts, diagnostic studies, animal studies, cost-benefit analysis, study with basic science outcomes

- Date cut off: nothing before 1971 *WHO recommended in this year to its nations that Rh testing and treatment with immune globulin be made part of the standard protocol of medical care for pregnant women

- Non-English publications

Condition or domain being studied

In most countries anti-D Immunoglobulin G is provided for rhesus (Rh)-negative patients. The administration

of Rh-immunoglobulin was first introduced in 1968 and significantly reduced immunization to D-antigen (Fiala 2003). There is sparse evidence for such intervention after abortion in early pregnancy (Fiala 2003). Early abortion includes threatened, spontaneous, surgical or medical abortion.

Currently, the WHO recommends Rh testing and administration of anti-D in populations with higher prevalence of Rh-negative individuals. Recently, the National Abortion Federation of North America published a statement that women presenting for early abortion can forego Rh testing and anti-D immunoglobulin administration (Mark 2019).

Participants/population

Reproductive aged women who have recently undergone surgical or medical management of pregnancy <12 weeks (includes induced, incomplete, spontaneous, or septic abortion)

Intervention(s), exposure(s)

Intervention:

Routine administration of Anti-D

Comparator(s)/control

Comparisons:

No administration of Anti-D

Context

Global setting

Main outcome(s)

Primary outcomes

1. Development of a positive Kleihauer Betke test (a test that detects fetal cells in the maternal blood).
2. Development of RhD alloimmunisation in a subsequent pregnancy.

Measures of effect

As defined by studies

Additional outcome(s)

Secondary outcomes

1. Detection of atypical blood group antibodies by positive indirect Coombs test after six months of exposure (non?prespecified outcome).
2. Need for increased surveillance for suspected fetal blood sampling and fetal transfusions in subsequent pregnancies.
3. Neonatal morbidity such as neonatal anaemia, jaundice, bilirubin encephalopathy, erythroblastosis, prematurity, hypoglycaemia (low blood sugar) in subsequent pregnancies.
4. Maternal adverse events of anti?D administration including anaphylactic reaction

Measures of effect

As defined by studies

Data extraction (selection and coding)

With less than 20 articles, all data extraction will be performed by Dr. Michelle Chan. Data extraction will be

obtained directly from the original published article. Recording of information for the data abstraction will occur in a table format, with the following headings. (1) Author, Year, Funding (2) Study Design, location, year(s) of data collection, follow-up (3) Population (4) Intervention (5) Comparison (6) Results (7) Strengths (8) Weaknesses (9) Risk of Bias

Risk of bias (quality) assessment

Risk of bias quality assessment will be addressed with criteria depending on study design.

RCT - List of factors for RCT quality framework: Adequate randomization (computer generated); Allocation concealment (opaque sealed envelopes); Blinding of participants and personnel; Blinding of outcome assessment;

Objective measurement of outcome; Incomplete data/lost to follow-up; Small sample size (need to define the size); Power calculations by appropriate outcome measure.

Observational studies - List of factors for observational quality framework: Population, Participation and follow up rates as above; Defined exposure (esp with comparator); Exposure defining, duration, timing; Measurement of isoimmunization; Confounding issue

Strategy for data synthesis

Data synthesis will primarily be descriptive analysis based on the outcomes. Pooled data will be reviewed with the GRADE system.

Given the anticipated small number of articles, this review will be primarily narrative. A descriptive analysis will provide summary of the studies including: study location, years of data collection, follow-up, population, intervention, comparison, results, strengths and weaknesses. Risk of bias will also be included.

A broader summary and tables of primary outcomes:

1. Development of a positive Kleihauer Betke test (a test that detects fetal cells in the maternal blood).
2. Development of RhD alloimmunisation in a subsequent pregnancy.

and secondary outcomes:

1. Detection of atypical blood group antibodies by positive indirect Coombs test after six months of exposure (non?prespecified outcome).
2. Need for increased surveillance for suspected fetal blood sampling and fetal transfusions in subsequent pregnancies.
3. Neonatal morbidity such as neonatal anaemia, jaundice, bilirubin encephalopathy, erythroblastosis, prematurity, hypoglycaemia (low blood sugar) in subsequent pregnancies.
4. Maternal adverse events of anti?D administration including anaphylactic reaction

and secondary outcomes (will also be included.

Analysis of subgroups or subsets

Not applicable

Contact details for further information

Michelle C Chan
michelle.c.chan@gmail.com

Organisational affiliation of the review

University of British Columbia/ World Health Organization, Department of Reproductive Health & Research

Review team members and their organisational affiliations

Dr Michelle Chan. University of British Columbia/ World Health Organization, Department of Reproductive Health & Research

Dr Caron Kim. World Health Organization, Department of Reproductive Health & Research

Dr Roopan Gill. World Health Organization, Department of Reproductive Health & Research

Collaborators

Ms Robyn Paynter. Cochrane Fertility Regulation Group Information Specialist, MLS

Type and method of review

Systematic review

Anticipated or actual start date

23 July 2019

Anticipated completion date

29 February 2020

Funding sources/sponsors

Not applicable, no funders

Conflicts of interest

Language

English

Country

Canada, Switzerland, United States of America

Stage of review

Review Ongoing

Subject index terms status

Subject indexing assigned by CRD

Subject index terms

Abortion, Induced; Abortion, Spontaneous; Female; Humans; Pregnancy; Rh Isoimmunization

Date of registration in PROSPERO

28 April 2020

Date of first submission

30 August 2019

Stage of review at time of this submission

Stage	Started	Completed
Preliminary searches	Yes	No
Piloting of the study selection process	Yes	No
Formal screening of search results against eligibility criteria	Yes	No
Data extraction	Yes	No
Risk of bias (quality) assessment	No	No
Data analysis	No	No

The record owner confirms that the information they have supplied for this submission is accurate and complete and they understand that deliberate provision of inaccurate information or omission of data may be construed as scientific misconduct.

The record owner confirms that they will update the status of the review when it is completed and will add publication details in due course.

Versions

28 April 2020