






OPEN ACCESS

Desire to Avoid Pregnancy scale: clinical considerations and comparison with other questions about pregnancy preferences

Jennifer Anne Hall ¹, Geraldine Barrett ¹, Judith M Stephenson,¹
Natalie Lois Edelman ^{2,3}, Corinne Rocca⁴

¹Research Department of Reproductive Health, UCL Institute for Women's Health, London, UK

²School of Sport & Health Sciences, University of Brighton, Brighton, UK

³Primary Care & Public Health, Brighton and Sussex Medical School, Brighton, UK

⁴Advancing New Standards in Reproductive Health (ANSIRH), Department of Obstetrics, Gynecology and Reproductive Sciences, University of California, San Francisco (UCSF) School of Medicine, Oakland, San Francisco, California, USA

Correspondence to

Dr Jennifer Anne Hall, Reproductive Health Research Department, UCL Institute for Women's Health, London, WC1E 6HU, UK; jennifer.hall@ucl.ac.uk

Received 10 November 2022
Accepted 18 January 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

To cite: Hall JA, Barrett G, Stephenson JM, et al. *BMJ Sex Reprod Health* Published Online First: [please include Day Month Year]. doi:10.1136/bmjsex-2022-201750

ABSTRACT

Background Clinicians and women of reproductive age would benefit from a reliable way to identify who is likely to become pregnant in the next year, in order to direct health advice. The 14-item Desire to Avoid Pregnancy (DAP) scale is predictive of pregnancy; this paper compares it with other ways of assessing pregnancy preferences to shortlist options for clinical implementation.

Methods A cohort of 994 UK women of reproductive age completed the DAP and other questions about pregnancy preferences, including the Attitude towards Potential Pregnancy Scale (APPS), at baseline and reported on pregnancies quarterly for a year. For each question, DAP item and combinations of DAP items, we examined the predictive ability, sensitivity, specificity, area under the receiver operating curve (AUROC), and positive and negative predictive values.

Results The AUROCs and predictive ability of the APPS and DAP single items were weaker than the full DAP, though all except one had acceptable AUROCs (>0.7). The most predictive individual DAP item was 'It would be a good thing for me if I became pregnant in the next 3 months', where women who strongly agreed had a 66.7% chance of pregnancy within 12 months and the AUROC was acceptable (0.77).

Conclusion We recommend exploring the acceptability to women and healthcare professionals of asking a single DAP item ('It would be a good thing for me if I became pregnant in the next 3 months'), possibly in combination with additional DAP items. This will help to guide service provision to support reproductive preferences.

INTRODUCTION

While there are a multitude of measures and screening tools available to predict pregnancy-related conditions, such as pre-eclampsia¹ or gestational diabetes,² there

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Clinicians do not currently have a valid and reliable way of asking women about their pregnancy preferences.

WHAT THIS STUDY ADDS

⇒ A single item from the Desire to Avoid Pregnancy (DAP) scale is effective at identifying who is likely to become pregnant in the next year; other questions, including the Attitude towards Potential Pregnancy Scale, are less discriminative but less burdensome than the full DAP Scale.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ The acceptability to women and health professionals of different ways of asking women about their pregnancy preferences in healthcare settings should be explored.

are no valid and reliable clinical tools to identify women who are likely to become pregnant (where we refer to 'women' this should be taken to include people who do not identify as women but who have the capability to become pregnant).¹ Such a tool would have great utility for clinicians working with women of reproductive age as it could be used to guide discussions on either preparation for pregnancy or contraceptive options, depending on if or when a pregnancy might be desired. This would simultaneously help to prevent unintended pregnancies and improve pregnancy outcomes and long-term health.³ Research shows that most clinicians see the value of such discussions⁴⁻⁷

but highlights the need for further evidence on the ‘validity, appropriate use, and integration into healthcare systems’.⁴

The Desire to Avoid Pregnancy (DAP) scale is a psychometrically validated measure of pregnancy preferences that was developed in the USA in 2019,⁸ and validated in the UK in 2022.⁹ It was shown to be highly predictive of pregnancy, with women with the lowest DAP score having an 80% chance of becoming pregnant within 12 months.⁹ Subsequently, the sensitivity and specificity of the DAP have been described (0.78 and 0.81, respectively, at a cut-point <2)¹⁰ and further analysis has shown how DAP score is associated with sociodemographic factors,¹⁰ findings which are in keeping with the literature on pregnancy preferences.^{11–13}

While there may be ways of incorporating the DAP scale into clinical practice, for example, preconsultation digital use or self-completion, a tool with 14 items is likely to be impractically lengthy in many clinical consultations. Use of selected items from the DAP, or other questions, may provide a brief and viable alternative if they distinguish sufficiently between who will and who will not become pregnant. There are limited options in terms of pregnancy intention screening tools with evidence of their performance. The only other measure with any psychometric evaluation is the five-question Attitude toward Potential Pregnancy Scale (APPS),¹⁴ though there is only one published use of it.¹⁵ Alternatively, ‘One Key Question’ (OKQ), which is a question not a measure, asks ‘Would you like to become pregnant in the next year?’ and is packaged with training and protocols.¹⁶ There is mixed evidence on the acceptability and effectiveness of the OKQ.^{17–20} There are no data on the performance or predictive ability of these alternatives and, until the DAP scale, there had not been a ‘gold standard’ to compare them to.

The aims of this paper are to: (1) evaluate the predictive ability, sensitivity, specificity, area under the receiver operating curve (AUROC) and positive and negative predictive values (PPV and NPV) of different methods of assessing pregnancy preferences and (2) examine the same criteria for individual DAP items and selected combinations of DAP items. Using the findings of these two aims, options are shortlisted for consideration for clinical implementation.

METHODS

Sample

We analysed data from a cohort of 994 non-pregnant women, aged 15 and over, who were recruited in the UK in October 2018 and followed up every 3 months for 1 year. The full details of recruitment and participation are described elsewhere,⁹ but in brief, people who self-reported as female, were premenopausal and not sterilised were recruited through advertising at a school, a university, a sexual health clinic and a

pregnancy termination clinic, as well as through online recruitment via both paid advertisements (Instagram and Facebook) and sharing through networks.⁹ The survey was programmed in RedCap^{21 22} and included the DAP scale, other questions about pregnancy preferences and sociodemographics. At each quarterly follow-up, participants were asked whether they were currently pregnant or had been pregnant since the last survey. Almost 90% (831/929) of eligible women completed follow-up at 12 months; those who were lost to follow-up were not significantly different by age, relationship status, ethnicity, baseline DAP score or number of children from those who did (suggesting that there is no selection bias in the lost to follow-up).¹⁰

Measures

Outcome

We created a binary variable of any incident pregnancy between baseline and 12 months. Given low attrition and to ease interpretation, we included participants in pregnancy denominators until they were lost to follow-up and report percentages rather than rates.

Attitude towards Potential Pregnancy Scale

The APPS is a five-item measure of a woman’s emotional outlook regarding a potential pregnancy (table 1).¹⁴ Each item is scored one-to-five on a visual anchored scale and summed to give an overall score from five to 25; higher scores represent a more positive attitude towards pregnancy. The APPS had a Cronbach’s alpha of 0.86 among 130 participants in the USA but has not been examined in the UK.

Other pregnancy preference questions

We compared the DAP score with three other single questions asking about pregnancy preferences that were available when the study was designed in 2018 (table 1). The ‘feelings question’ was used in the ADAPT study²³; the ‘trying’ and ‘thinking’ questions are ones that clinicians in the UK have told us are the kind of questions they currently use. We did not use OKQ as it is a proprietary tool that is unlikely to be available for widespread implementation in the UK, but the ‘thinking’ question is similar.

DAP Scale

The DAP scale is a psychometrically validated measure covering three conceptual domains: (1) cognitive desires and preferences; (2) affective feelings and attitudes; and (3) anticipated practical consequences (table 1).⁸ Each item uses a Likert scale, scored 0–4, to ask women how much they agree or disagree with a statement about either becoming pregnant in the next 3 months or having a baby in the next year. Responses are averaged, producing a total score between 0 and 4; higher scores represent a higher DAP. It was found to be highly acceptable in the UK evaluation.⁹

Table 1 Wording of questions about preferences regarding future pregnancy

| Question wording | Response options | Domain |
|--|---|----------------|
| DAP (with label for reference in text) | | |
| I wouldn't mind if I became pregnant in the next 3 months Pregnant: wouldn't mind | 0 Strongly agree—4 strongly disagree | 1 |
| It would be a good thing for me if I became pregnant in the next 3 months Pregnant: good thing for me | 0 Strongly agree—4 strongly disagree | 1 |
| Thinking about becoming pregnant in the next 3 months makes me feel unhappy. Pregnant: unhappy | 4 Strongly agree—0 strongly disagree | 2 |
| Thinking about becoming pregnant in the next 3 months makes me feel excited. Pregnant: excited | 0 Strongly agree—4 strongly disagree | 2 |
| Becoming pregnant in the next 3 months would bring me closer to my main partner Pregnant: closer to partner | 0 Strongly agree—4 strongly disagree | 3 |
| I want to have a baby within the next year. Baby: want | 0 Strongly agree—4 strongly disagree | 1 |
| If I had a baby in the next year, it would be bad for my life. Baby: bad for life | 4 Strongly agree—0 strongly disagree | 1 |
| It would be a positive addition to my life to have a baby in the next year. Baby: positive addition to life | 0 Strongly agree—4 strongly disagree | 1 |
| It would be the end of the world for me to have a baby in the next year Baby: end of the world for me | 4 Strongly agree—0 strongly disagree | 1 |
| Thinking about having a baby within the next year makes me smile Baby: makes me smile | 0 Strongly agree—4 strongly disagree | 2 |
| Thinking about having a baby within the next year make me feel stressed out. Baby: stressed out | 4 Strongly agree—0 strongly disagree | 2 |
| I would feel a loss of freedom if I had a baby in the next year. Baby: loss of freedom | 4 Strongly agree—0 strongly disagree | 3 |
| If I had a baby in the next year, it would be hard for me to manage raising the child. Baby: hard for me to manage | 4 Strongly agree—0 strongly disagree | 3 |
| I would worry that having a baby in the next year would make it harder for me to achieve other things in my life. Baby: hard to achieve other things | 4 Strongly agree—0 strongly disagree | 3 |
| AAPS | | Not applicable |
| How much do you want to be pregnant now? | 1 Not at all—5 Very much | |
| How important is it to you to avoid become pregnant now? | 5 Not at all important—1 Very important | |
| How worried would you be if you were pregnant now? | 5 Not worried at all—1 Very worried | |
| How upset would you be if you were pregnant now? | 5 Not at all upset—1 Very upset | |
| How happy would you be if you were pregnant now? | 1 Very unhappy—5 Very happy | |
| Feeling question | | |
| Overall, when I think about become pregnant in the next 3 months, I feel: | 1 Mostly positive 2 More positive than negative 3 In the middle between positive and negative 4 I don't feel strongly one way or the other 5 More negative than positive 6 Most negative | |
| Trying question | | |
| Are you currently trying to get pregnant? | 1 Yes | |
| Thinking question | | |
| Are you thinking about trying to get pregnant in the next year? | 2 Yes | |
| Domains: 1 = Cognitive desires and preferences, 2 = Affective feelings and attitudes, 3 = Anticipated practical consequences. Further information on the DAP wording and instructions for use are available here: https://www.ansirh.org/sites/default/files/publications/files/desire_to_avoid_pregnancy_scale_english_6_22_20.pdf . AAPS, Attitude towards Potential Pregnancy Scale; DAP, Desire to Avoid Pregnancy. | | |

Plan of analysis

Aim 1: evaluating the performance of other methods of assessing pregnancy preferences

We assessed the APPS's reliability (internal consistency) with Cronbach's alpha (>0.7 considered acceptable)²⁴ and checked that all item-rest correlations were >0.2 .²⁵ We examined construct (structural) validity via principal components analysis and considered the scale structurally valid if all items loaded on to one component with an Eigenvalue >1 .²⁶

To compare the assessment of pregnancy preferences by the DAP versus the APPS and the three individual questions ('feeling', 'trying' and 'thinking'), we examined the relationships among them. As the DAP and the APPS yield continuous scores, we used the Pearson correlation coefficient to examine the strength of association, expecting a negative correlation (ie, as DAP score increases, APPS score decreases), considering a strong correlation to be lower than -0.7 . This also served as a test of the APPS's concurrent validity. The range of DAP scores within, and distribution of DAP scores across, the response options of each question was examined. The Kruskal-Wallis test was used to assess whether differences in median DAP score across response options within each question were significant.

To investigate how well each approach predicted actual pregnancy, we modelled the probability of pregnancy using logistic regression models and examined the sensitivity, specificity, AUROC, PPV and NPV, using the Youden index suggested cut-point.²⁷ The AUROC represents the tool's ability to identify who will become pregnant. An AUROC of $0.7-0.8$ was considered acceptable, $0.8-0.9$ excellent and >0.9 outstanding.²⁸

Aim 2: examination of the performance of individual and combinations of DAP items

We explored the predictive ability of each individual DAP item, as well as selected combinations of items, using logistic regression, with incident pregnancy as the outcome. The item combinations were designed to ensure coverage of the three conceptual domains, positively and negatively framed items, different time frames and use of 'pregnancy' and 'baby', as well as the predictive ability of the items. The combinations were cross-validated by developing them on one half of the data (created with a random split) and tested on the other half to avoid overfitting and give a more accurate estimate of how the questions perform outside the data they were developed from.^{29 30} The results from the testing data are presented. The predictive ability, sensitivity, specificity, AUROC, PPV and NPV of the selected items/combinations were calculated (using the Youden index suggested cut-point)²⁷ and compared with the total DAP scale.

To inform which question(s) provided the best balance between brevity and predictive ability, the performance of the single questions, the APPS and

selected DAP items/combinations were compared using the predictive ability, AUROC and number of items/questions, to make recommendations on which item/question(s) should be taken forward for further consideration.

Patient and public involvement

We undertook public involvement in the development of our overall programme of research on pregnancy planning, the P3 Study, by discussing and refining ideas during the grant application stage with an existing patient and public involvement (PPI) group within our team. Findings were frequently discussed with the P3 Study's PPI group to inform the next steps and the group is currently planning our wider dissemination.

RESULTS

Sample

As previously described,⁹ the baseline cohort of 994 women were aged 15–50 years (median=31, IQR=23–36, mean=29.7). The sample was fairly evenly distributed by age, with 28% aged 15–24 years, 36% aged 25–34 and 31% aged over 35 (4% did not give an age). Most (82%) were in a relationship and 82% described themselves as heterosexual. The majority (84%) identified as white; 6.9% as Asian, 2.6% as Mixed, 2.4% as Black and 0.9% as Other (3.2% missing/preferred not to say). Around one in 10 women did not speak English as their first language. The sample was quite highly educated with 39% having an undergraduate degree and 31% a postgraduate or professional qualification. Fifty-seven per cent had one or more children in their household. The dataset is available in the UCL Research Data Repository.³¹

Aim 1: evaluating the performance of other methods of assessing pregnancy preferences

The full range of APPS scores^{5–25} was reported in our cohort. The Cronbach's alpha for the APPS was 0.93, all item-rest correlations were >0.2 and positive, and all items loaded on to one component with an Eigenvalue of 3.87. The Pearson correlation coefficient between the APPS and DAP was -0.893 ($p<0.001$) showing a strong negative correlation (ie, as a woman's attitude towards pregnancy becomes more positive, her DAP score reduces.) The median DAP score was statistically significantly different across the response options for the APPS and each of the 'feeling', 'trying' and 'thinking' questions (p values in [table 2](#)). For example, women who felt 'mostly positive' about pregnancy had a median DAP score of 0.64 (low DAP) whereas women who felt 'mostly negative' about pregnancy had a median DAP score of 3.50 (high DAP).

The AUROCs and predictive ability of the APPS and the single questions were weaker than the full DAP (AUROC 0.87, predictive ability 79.4%), though all

Table 2 Relationship with DAP and predicted probability, sensitivity, specificity, AuROC, PPV and NPV of other questions

| APPS score* | n | Median DAP score | IQR | P value | Predicted probability of pregnancy | 95%CI | Sensitivity | Specificity | AuROC | PPV | NPV |
|---|-----|------------------|--------------|---------|------------------------------------|----------------|-------------|-------------|-------|-----|-----|
| 5 | 265 | 3.64 | 3.29 to 3.86 | <0.001 | 0.03 | 0.019 to 0.044 | 78% | 74% | 0.76 | 36% | 95% |
| 10 | 59 | 2.50 | 2.29–2.71 | | 0.09 | 0.063 to 0.107 | | | | | |
| 15 | 35 | 1.71 | 1.50–2.07 | | 0.21 | 0.173 to 0.243 | | | | | |
| 20 | 13 | 1.07 | 0.86–1.50 | | 0.47 | 0.362 to 0.490 | | | | | |
| 25 | 42 | 0.57 | 0.36–0.64 | | 0.68 | 0.594 to 0.761 | | | | | |
| Overall, when I think about becoming pregnant in the next 3 months, I feel: | | | | | | | | | | | |
| Mostly positive | 117 | 0.64 | 0.43–0.93 | <0.001 | 0.56 | 0.47 to 0.65 | 75% | 76% | 0.75 | 37% | 94% |
| More positive than negative | 84 | 1.29 | 1.07–1.64 | | 0.37 | 0.26 to 0.48 | | | | | |
| In the middle | 107 | 2.00 | 1.57–2.36 | | 0.14 | 0.07 to 0.21 | | | | | |
| Don't feel strongly | 21 | 2.14 | 2.00–2.43 | | 0.07 | –0.06 to 0.19 | | | | | |
| More negative than positive | 264 | 2.64 | 2.36–3.00 | | 0.09 | 0.06 to 0.13 | | | | | |
| Mostly negative | 395 | 3.50 | 3.14–3.79 | | 0.03 | 0.02 to 0.05 | | | | | |
| Are you currently trying to get pregnant? | | | | | | | | | | | |
| Yes | 80 | 0.64 | 0.36–0.86 | >0.001 | 0.69 | 0.58 to 0.50 | 37% | 97% | 0.67 | 69% | 89% |
| No | 907 | 2.79 | 2.07–3.43 | | 0.11 | 0.09 to 0.13 | | | | | |
| Are you thinking about trying to get pregnant in the next year? | | | | | | | | | | | |
| Yes | 96 | 1.21 | 0.79–1.71 | <0.001 | 0.53 | 0.42 to 0.63 | 71% | 84% | 0.77 | 34% | 96% |
| Maybe | 97 | 1.86 | 1.36–2.21 | | 0.15 | 0.07 to 0.22 | | | | | |
| No | 707 | 3.07 | 2.57–3.57 | | 0.04 | 0.20 to 0.06 | | | | | |

*APPS score is continuous from 5 to 25, 5 points shown for illustration, therefore, n does not add up to the full sample.

APPS, Attitude towards Potential Pregnancy Scale; AUROC, area under the receiver operating curve; DAP, Desire to Avoid Pregnancy; NPV, negative predictive value; PPV, positive predictive value.

except the 'trying' question had acceptable AUROCs of >0.7.

Aim 2: evaluating the performance of individual and combinations of DAP items

All DAP items and domains were associated with pregnancy. The best performing individual DAP item in terms of pregnancy prediction was 'Pregnancy: good thing for me', where there was a 66.7% chance of pregnancy within 12 months among women who strongly agreed (table 3). At a cut point of 2.5 this item also had an acceptable AUROC (0.77).

Adding a second item of 'Baby: end of the world for me' improved the specificity and PPV, without affecting the AUROC. The highest AUROC (0.79) was achieved with the combination of three items 'Pregnancy: good thing for me', 'Baby: want' and 'Baby: end of the world for me', which also had the highest PPV (57.1%). The item combinations had AUROCs between 0.77 and 0.79, in line with that of the individual item 'Pregnancy: good thing for me' suggesting little additional gain.

DISCUSSION

This is the first examination of the predictive ability of two measures (DAP and APPS) and three single questions ('feeling', 'trying' and 'thinking') about pregnancy preferences on a sample that is broadly representative of women of reproductive age in the UK. In terms of the AUROC and the predicted probability of pregnancy in the next year, the performance of the APPS and the other questions was weaker than the complete DAP, though to varying degrees. The APPS and the other questions are all shorter than the DAP and therefore potentially less burdensome for clinical use. Arguably, their poorer performance is offset by their brevity. Our data show that the single question 'Are you currently trying to get pregnant?' had low sensitivity (37%) and the lowest AUROC (0.67). While women answering yes to this question were highly likely to become pregnant (PPV 69%) there were more pregnancies among women who answered no. Questions like this, which force women to answer yes-or-no, fail to recognise the complexity of the concept of pregnancy preferences, leading to misclassification.³²

When considering which question(s) might be best for clinical use, the trade-off between sensitivity and specificity, and the PPV and NPV is important. In addition, the number of questions that need to be asked, and the complexity of combining them to achieve a score, could be a barrier in consultations, therefore, having fewer questions will likely increase uptake. As our analysis shows, single questions generally have lower AUROCs than a set of questions, however, selected items from the DAP, which have been developed based on rigorous theoretical groundwork, do have higher AUROCs than the less carefully constructed questions. Neither the DAP nor the APPS were designed to be

spoken, whereas the other questions lend themselves more easily to being asked verbally.

While we did not use the OKQ, good correlation was previously seen between the DAP score and the OKQ in the USA.³³ As that study noted, DAP scores ranged widely within each OKQ response option; women who responded to the OKQ with 'want to get pregnant in future but not in next year' had DAP scores ranging from 1 to 4. Our data demonstrated the same patterns; women who reported feeling 'mostly negative' towards pregnancy had DAP scores ranging from 0.286 to 4, this shows the nuance that the DAP can capture and demonstrates the heterogeneity missed by a single question. In addition, the OKQ is frequently the least-favoured option in studies asking women and healthcare professionals' preference from a range of questions.^{5 34 35} There have been no comparisons made to the APPS.

There are well-known barriers to implementation of preconception care,³⁶ and it is likely that these apply to pregnancy intention screening as the gateway to identifying who requires preconception care. Lack of time is a commonly cited barrier,³⁷ as is the lack of clarity over whose role it should be.³⁶ However, there is growing evidence of the feasibility and acceptability, to women and healthcare professionals, of pregnancy intention screening,^{5-7 38} and of the effectiveness of preconception interventions in primary care.³⁹

Strengths and limitations

We used a large, broadly representative dataset and a validated measure (the DAP) to assess the performance of a range of questions for assessing pregnancy preferences, and have provided preliminary evidence that the APPS is valid in the UK. Given the intellectual property limitations on the use of the OKQ, which affect the feasibility of national rollout in the UK, we did not include it in our research. While we used a split dataset to conduct cross-validation within our sample, which is a strength, confirmation of our findings in other populations, particularly those that are more diverse in terms of ethnicity, education and socioeconomic status, would provide further confidence in selecting the most appropriate question(s). The selected DAP items/combinations were based on their potential as screening tools for pregnancy, as well as theoretical considerations; item-response theory-based analysis of the psychometric properties of these combinations could be conducted.

CONCLUSION

Discussions about pregnancy preferences are important, regardless of whether the woman wants to become pregnant in the future or not. Ensuring that those who do not wish to become pregnant have the right support to avoid pregnancy is just as important as identifying those who might benefit from prepregnancy health advice or those who may benefit from both. Equally,

Table 3 Exploration of the predicted probability, sensitivity, specificity, AuROC, PPV and NPV of selected individual and combined DAP items

| Description | Predicted probability of pregnancy at DAP score | | | | Empirical cut-point | Sensitivity | Specificity | AuROC | PPV | NPV | Domains covered | Positive and negatively worded items |
|---|---|-------|------|-------|---------------------|-------------|-------------|-------|------|---------------|-----------------|--------------------------------------|
| | 0 | 4 | 4 | 4 | | | | | | | | |
| Total DAP: All 14 items | 79.4% | 0.9% | 1.96 | 78.0% | 81.0% | 0.85 | 43.0% | 95.0% | All | Yes | | |
| Single DAP item: Pregnant: good thing for me. | 66.7% | 2.1% | 2.5 | 68.1% | 86.3% | 0.77 | 49.0% | 93.3% | 1 | Positive only | | |
| Single DAP item: Baby: end of the world for me. | 27.4% | – | 0.5 | 65.3% | 77.7% | 0.72 | 36.2% | 92.0% | 1 | Negative only | | |
| Two DAP items: Pregnant: good thing for me and Baby: end of the world for me. | 71% | 0.7% | 1.25 | 65.3% | 90.1% | 0.78 | 56.0% | 93.1% | 1 | Yes | | |
| Three DAP items: Pregnant: good thing for me, Pregnant: excited, Baby: hard to achieve other things. | 83% | 2.8% | 2.15 | 62.5% | 87.1% | 0.75 | 48.4% | 92.3% | All | Yes | | |
| Three DAP items: Pregnant: good thing for me, Pregnant: excited, Baby: end of the world for me. | 73% | 1.2% | 1.49 | 66.7% | 87.4% | 0.77 | 50.5% | 93.1% | 1, 2 | Yes | | |
| Three DAP items: Pregnant: good thing for me, Baby: end of the world for me, Baby: makes me smile. | 68% | 1.0% | 1.49 | 69.4% | 84.4% | 0.77 | 46.3% | 93.5% | 1, 2 | Yes | | |
| Three DAP items: Pregnant: good thing for me, Baby: want, Baby: end of the world for me. | 72% | 13.9% | 1.49 | 66.7% | 90.3% | 0.79 | 57.1% | 93.3% | 1 | Yes | | |
| Three DAP items: Pregnant: good thing for me, Baby: end of the world for me, Baby: hard for me to manage. | 76% | 8.8% | 1.82 | 75.0% | 77.2% | 0.76 | 38.8% | 94.1% | 1, 3 | Yes | | |
| Four DAP items: Pregnant: good thing for me, Pregnant: excited, Baby: want, Baby: end of the world for me. | 73% | 1.4% | 1.63 | 66.7% | 89.2% | 0.78 | 54.5% | 93.2% | 1, 2 | Yes | | |
| Four DAP items: Pregnant: good thing for me, Pregnant: excited, Baby: end of the world for me, Baby: hard to achieve other things. | 81% | 1.2% | 1.88 | 66.7% | 83.9% | 0.75 | 44.4% | 92.9% | All | Yes | | |
| Six DAP items: Pregnant: wouldn't mind, Pregnant: unhappy, Baby: want, Baby: positive addition to life, Baby: makes me smile, Baby: hard for me to manage. | 73% | 6.1% | 1.93 | 75.0% | 80.9% | 0.78 | 43.2% | 94.4% | All | Yes | | |
| AUROC, area under the receiver operating curve; DAP, Desire to Avoid Pregnancy; NPV, negative predictive value; PPV, positive predictive value . | | | | | | | | | | | | |

for those who have never formally considered their preferences, it provides an opportunity to empower them and increase their agency by highlighting that people do have choices about pregnancy and parenthood (recognising the effect of external factors) and encouraging them to explore their aspirations.

In the context of a face-to-face clinical encounter, where the full DAP is less likely to be suitable, a tool to assess people's preferences regarding a future pregnancy needs to be both practical (short) and discriminative, that is, identify who is and is not likely to become pregnant in the short term so that the appropriate advice can be given. Based on our findings, we recommend exploring the acceptability to women and healthcare professionals of a single item from the DAP ('It would be a good thing for me if I became pregnant in the next 3 months') or a combination of this with two additional DAP items (either 'Baby: want' and 'Baby: end of the world for me' or 'Pregnant: excited' and 'Baby: hard to achieve other things') adapted from the written format to a spoken one.

Twitter Jennifer Anne Hall @jennyhall33 and Natalie Lois Edelman @natalieedelman

Acknowledgements We would like to thank all the women who took part in the P3 study, the P3 Study steering committee and Catherine Stewart for her support in preparing the manuscript for submission.

Contributors JAH: guarantor, conceptualisation, funding acquisition, methodology, investigation, data curation, formal analysis, writing-original draft, visualisation, writing-review and editing, project administration. GB: conceptualisation, methodology, writing-review and editing. JMS: conceptualisation, writing-review and editing. CR: conceptualisation, methodology, writing-review and editing. NLE: methodology, writing-review and editing.

Funding The study was funded by an NIHR Advanced Fellowship held by JH (PDF-2017-10-021).

Disclaimer The funder had no role in the study design; in the collection, analysis and interpretation of the data; in the writing of the report; or in the decision to submit the paper for publication.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Consent obtained directly from patient(s).

Ethics approval Ethical approval was granted by the UCL Research Ethics Committee (ref 3974.003). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. The dataset is available in the UCL Research Data Repository, doi 10.5522/04/21263007.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link

to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Jennifer Anne Hall <http://orcid.org/0000-0002-2084-9568>

Geraldine Barrett <http://orcid.org/0000-0002-9738-1051>

Natalie Lois Edelman <http://orcid.org/0000-0001-6935-8877>

REFERENCES

- 1 Tan MY, Wright D, Syngelaki A, *et al*. Comparison of diagnostic accuracy of early screening for pre-eclampsia by NICE guidelines and a method combining maternal factors and biomarkers: results of spree. *Ultrasound Obstet Gynecol* 2018;51:743–50.
- 2 Artzi NS, Shilo S, Hadar E, *et al*. Prediction of gestational diabetes based on nationwide electronic health records. *Nat Med* 2020;26:71–6.
- 3 Stephenson J, Heslehurst N, Hall J, *et al*. Before the beginning: nutrition and lifestyle in the preconception period and its importance for future health. *Lancet* 2018;391:1830–41.
- 4 Romero D, Manze M, Roberts L, *et al*. Stakeholder perspectives in new york state on reproductive health and pregnancy intention screening in primary care: a qualitative study. *Contraception* 2022;112:93–9.
- 5 Manze MG, Calixte C, Romero DR, *et al*. Physician perspectives on routine pregnancy intention screening and counseling in primary care. *Contraception* 2020;101:91–6.
- 6 Shah SD, Prine L, Waltermaurer E, *et al*. Feasibility study of family planning services screening as clinical decision support at an urban federally qualified health center network. *Contraception* 2019;99:27–31.
- 7 Bello JK, Adkins K, Stulberg DB, *et al*. Perceptions of a reproductive health self-assessment tool (RH-SAT) in an urban community health center. *Patient Educ Couns* 2013;93:655–63.
- 8 Rocca CH, Ralph LJ, Wilson M, *et al*. Psychometric evaluation of an instrument to measure prospective pregnancy preferences: the desire to avoid pregnancy scale. *Med Care* 2019;57:152–8.
- 9 Hall J, Barrett G, Rocca C. Evaluation of the desire to avoid pregnancy scale in the UK: a psychometric analysis including predictive validity. *BMJ Open* 2022;12:e060287.
- 10 Hall JA, Barrett G, Stephenson J, *et al*. Predictive ability of the desire to avoid pregnancy scale. *Sexual and Reproductive Health* 2022.
- 11 Hill B, Ling M, Mishra G, *et al*. Lifestyle and psychological factors associated with pregnancy intentions: findings from a longitudinal cohort study of Australian women. *Int J Environ Res Public Health* 2019;16:24.
- 12 Lang AY, Harrison CL, Barrett G, *et al*. Opportunities for enhancing pregnancy planning and preconception health behaviours of Australian women. *Women Birth* 2021;34:e153–61.
- 13 Geist C, Everett BG, Simmons RG, *et al*. Changing lives, dynamic plans: prospective assessment of 12-month changes in pregnancy timing intentions and personal circumstances using data from her salt lake. *PLoS One* 2021;16:e0257411.
- 14 Paterno MT, Han H-R. Development and psychometric testing of the attitude toward potential pregnancy scale. *J Obstet Gynecol Neonatal Nurs* 2014;43:710–8.
- 15 Paterno MT, Hayat MJ, Wenzel J, *et al*. A mixed methods study of contraceptive effectiveness in a relationship context among young adult, primarily low-income African American women. *J Racial Ethn Health Disparities* 2017;4:184–94.

- 16 Allen D, Hunter MS, Wood S, *et al.* One key question®: first things first in reproductive health. *Matern Child Health J* 2017;21:387–92.
- 17 Ferketa M, Schueler K, Song B, *et al.* Facilitators of and barriers to successful implementation of the one key question® pregnancy intention screening tool. *Womens Health Rep (New Rochelle)* 2022;3:326–34.
- 18 Fitch J, Dorney E, Tracy M, *et al.* Acceptability and usability of “one key question” in Australian primary health care. *Aust J Prim Health* 9, 2022.
- 19 Song B, White VanGompel E, Wang C, *et al.* Effects of clinic-level implementation of one key question® on reproductive health counseling and patient satisfaction. *Contraception* 2021;103:6–12.
- 20 Thorman A, Engle A, Brintz B, *et al.* Quantitative and qualitative impact of one key question on primary care providers’ contraceptive counseling at routine preventive health visits. *Contraception* 2022;109:73–9.
- 21 Harris PA, Taylor R, Minor BL, *et al.* The redcap consortium: building an international community of software platform partners. *J Biomed Inform* 2019;95:103208.
- 22 Harris PA, Taylor R, Thielke R, *et al.* Research electronic data capture (redcap) -- a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81.
- 23 Rocca CH, Parra M, Muñoz I, *et al.* Comparison of pregnancy preferences preceding vs year 1 of the COVID-19 pandemic. *JAMA Netw Open* 2022;5:e2220093.
- 24 Cronbach LJ. *Essentials of psychologic testing*. 5th ed. New York: Harper & Row, 1990.
- 25 Streiner DL, Norman GR. *Health measurement scales: a practical guide to their development and use [program]*. Oxford: Oxford: Oxford University Press, 2008.
- 26 Kline P. *The new psychometrics: science, psychology and measurement*. London: Routledge, 1998.
- 27 Youden WJ. Index for rating diagnostic tests. *Cancer* 1950;3:32–5.
- 28 Hosmer DW, Lemeshow S. *Applied logistic regression*. 2nd ed. Hoboken, NJ, USA: John Wiley and Sons, 2000: 160–4.
- 29 Koul A, Becchio C, Cavallo A. Cross-validation approaches for replicability in psychology. *Front Psychol* 2018;9:1117.
- 30 Yarkoni T, Westfall J. Choosing prediction over explanation in psychology: lessons from machine learning. *Perspect Psychol Sci* 2017;12:1100–22.
- 31 Hall J. *Data from: data for DAP predictive ability and comparison*. UCL Research Data Repository, 2022.
- 32 Barrett G, Wellings K. What is a “planned” pregnancy? Empirical data from a british study. *Soc Sci Med* 2002;55:545–57.
- 33 Stulberg DB, Datta A, White VanGompel E, *et al.* One key question® and the desire to avoid pregnancy scale: a comparison of two approaches to asking about pregnancy preferences. *Contraception* 2020;101:231–6.
- 34 Jones HE, Calixte C, Manze M, *et al.* Primary care patients’ preferences for reproductive health service needs assessment and service availability in new york federally qualified health centers. *Contraception* 2020;101:226–30.
- 35 Manze MG, Romero DR, Sumberg A, *et al.* Women’s perspectives on reproductive health services in primary care. *Fam Med* 2020;52:112–9.
- 36 Steel A, Lucke J, Reid R, *et al.* A systematic review of women’s and health professional’s attitudes and experience of preconception care service delivery. *Fam Pract* 2016;33:588–95.
- 37 Heyes T, Long S, Mathers N. Preconception care: practice and beliefs of primary care workers. *Fam Pract* 2004;21:22–7.
- 38 Hall J, Chawla M, Watson D, *et al.* Addressing reproductive health needs across the life course: an integrated, community-based model combining contraception and preconception care. *Lancet Public Health* 2023;8:e76–84.
- 39 Withanage NN, Botfield JR, Srinivasan S, *et al.* Effectiveness of preconception care interventions in primary care: a systematic review protocol. *BJGP Open* 2022;6:BJGPO.2021.0191.