

SUPPLEMENTAL FIGURES AND TABLES

Table 1S. PMV report of abortion outcomes for clients obtaining misoprostol for induced abortion from their shop by reported gestational age of client and misoprostol dose dispensed.

	Complete abortion	Abortion with complications managed by PMV	Abortion with complications referred to another provider
PMV clients (n=3619)*	3489 (96.4)	25 (0.7)	105 (2.9)
Gestational age			
≤ 9 weeks (n=2884)	2791(96.8)	16 (0.6)	77 (2.7)
10-12 weeks (n=620)	593 (95.6)	9 (1.5)	18 (2.9)
13 weeks+ (n=115)	105 (91.3)	0	10 (8.7)
Misoprostol dose dispensed (mcg)			
Less than 800 (n=504)	464 (92.1)	7(1.4)	33 (6.5)
800 (n=2248)	2186 (97.2)	10 (0.4)	55 (2.3)
1000-1400 (n=98)	86 (87.8)	6 (6.1)	6 (6.1)
1600-2400 (n=755)	739 (97.9)	2 (0.3)	14 (1.9)
2600+ (n=14)	14 (100)	0	0

*Analysis excludes women who were missing data on gestational age (n=499), misoprostol dose dispensed (n=1140) or abortion outcome (n=493).




Figure 1S. Sample of PMVs trained, collection of data from PMVs, and recruitment of women during the intervention period.